

Strang splitting for a semilinear Schrödinger equation with damping and forcing

Tobias Jahnke, Marcel Mikl, Roland Schnaubelt

CRC Preprint 2016/4, February 2016

KARLSRUHE INSTITUTE OF TECHNOLOGY

CRC 1173



Wave
phenomena

Participating universities



Universität Stuttgart

EBERHARD KARLS
UNIVERSITÄT
TÜBINGEN



Funded by

DFG

ISSN 2365-662X

Strang splitting for a semilinear Schrödinger equation with damping and forcing ^{*}

TOBIAS JAHNKE[†], MARCEL MIKL[‡] AND ROLAND SCHNAUBELT[§]

Karlsruhe Institute of Technology, Department of Mathematics,
D-76128 Karlsruhe, Germany

February 23, 2016

Abstract

We propose and analyze a Strang splitting method for a cubic semilinear Schrödinger equation with forcing and damping terms. The nonlinear part is solved analytically, whereas the linear part – space derivatives, damping and forcing – is approximated by the exponential trapezoidal rule. The necessary operator exponentials and ϕ -functions can be computed efficiently by fast Fourier transforms if space is discretized by spectral collocation. We show wellposedness of the problem and $H^4(\mathbb{T})$ regularity of the solution for initial data in $H^4(\mathbb{T})$ and sufficiently smooth forcing. Under these regularity assumptions, we prove a first-order error bound in $H^1(\mathbb{T})$ and a second-order error bound in $L^2(\mathbb{T})$ on bounded time-intervals. Nonlinear Schrödinger equation; Strang splitting; error analysis; stability; wellposedness; regularity.

1 Introduction

Nonlinear Schrödinger equations (NLS) occur in many different forms and describe a multitude of different phenomena, such as Bose-Einstein condensates, small-amplitude surface water waves, Langmuir waves in hot plasmas, or signal processing through optical fibers, to name but a few. The intriguing properties – for example conservation of norm, energy, and momentum, near-conservation of actions over long times, existence of solitary waves, or possible blow-up – have inspired and challenged mathematicians for a long time. Surveys about these topics can be found, e.g., in the monographs of [Caz03] and [SS99].

^{*}We gratefully acknowledge financial support by the Deutsche Forschungsgemeinschaft (DFG) through RTG 1294 and CRC 1173.

[†]Corresponding author. Email: tobias.jahnke@kit.edu

[‡]Email: marcel.mikl@kit.edu

[§]Email: schnaubelt@kit.edu

In most applications, the solution of the NLS has to be approximated by a numerical scheme. For problems on the d -dimensional torus \mathbb{T}^d , splitting methods with spectral collocation in space are particularly popular. These integrators are based on the observation that the linear and the nonlinear part of the NLS can be solved at low computational costs in the absence of the other part. The splitting approach can also be applied to the Gross-Pitaevskii equation, a NLS on \mathbb{R}^d with constraining polynomial potential, by using the basis of Hermite functions for the space discretization. The accuracy of such integrators has been analyzed, e.g., by [BBD02], [DT13], [Fao12], [Gau11], [HKT14], [KNT13], [Lub08], [ML15], and [Tha12]. The long-time behavior of numerical solutions, in particular the (near-)conservation of invariants over long times and the stability of plane waves, has been investigated by [Fao12], [FGL14], [GL10], and for exponential integrators by [CG12].

In this article we consider the *Lugiato-Lefever equation*, a cubic, focusing NLS which, in contrast to the “classical” NLS, contains a damping and a forcing term; cf. [LL87]. This equation has been proposed as a model for the formation of Kerr-frequency combs in microresonators coupled to optical waveguides and driven by an external pump tuned to a resonance wavelength; see [CY10] and [HHR⁺12]. The frequency combs generated by such a device can be used as optical sources for high-speed data transmission. In the mathematical model, the forcing term represents the external pump, whereas the radiation into the waveguide is modeled by the damping term. In practice it is typically not clear which parametrization generates a suitable frequency comb. As a consequence, the Lugiato-Lefever equation has to be solved many times with different parametrizations, which requires a reliable and efficient simulation method. So far, however, both the properties of the exact solution and the performance of numerical integrators for its approximation are only poorly understood.

Our contribution in this article is twofold. First, we show global wellposedness of the Lugiato-Lefever equation and $H^4(\mathbb{T})$ regularity of the solution for initial data in $H^4(\mathbb{T})$ and a forcing function in $C^{2-j}([0, T]; H^j(\mathbb{T}))$ for $j = 0, 1, 2$. In contrast to the NLS without damping and forcing where wellposedness has been investigated in detail (cf. [Caz03]) such an analysis has been missing for the Lugiato-Lefever equation so far. Our result is derived by means of a modified energy functional and regularity properties of several differentiated versions of the equation. Second, we extend the Strang splitting approach to the Lugiato-Lefever equation and provide an error analysis for this method. The linear inhomogeneous part (including the space derivatives and the forcing/damping terms) is propagated by an exponential integrator whereas the nonlinear part is solved exactly as for the standard NLS. We prove that under the above regularity conditions on the initial data and the forcing the method converges on bounded time-intervals with the classical order 2 in $L^2(\mathbb{T})$, and with order 1 in $H^1(\mathbb{T})$; see Theorem 2 below. The proof consists of several steps which are formulated as self-contained results. As in [Lub08], the classical argument “consistency plus stability yields convergence” must be suitably adapted, because the stability result (Theorem 5 below) assumes the numerical solution to be bounded in $H^1(\mathbb{T})$. This a-priori bound is needed to control the nonlinearity.

In the proof of Theorem 2 this bound is verified by means of an error bound of second order for the local error in $H^1(\mathbb{T})$, in addition to the local (third order) error bound in $L^2(\mathbb{T})$ required for consistency (see Theorems 3 and 4, respectively).

Following [Lub08], many authors have used the calculus of Lie derivatives and commutator bounds in their error analysis, e.g. [DT13], [Fao12], [Gau11], [HKT14], [KNT13], [ML15], and [Tha12]. In contrast to these works, we avoid the notationally rather involved Lie derivatives, because in case of the Lugiato-Lefever equation the iterated commutators between the linear and nonlinear part are not the only source of error: An additional difficulty arising in our situation is the fact that the forcing term is coupled to the space derivatives and to the nonlinear part in a complicated way. Instead of Lie derivatives, our error formulas are based on iterated variation-of-constants formulas, the calculus of ϕ -functions (see Section 3) and the exponential trapezoidal rule (see formula (15)). The resulting error formulas are quite involved, and it requires some effort to keep track of the necessary regularity of the exact and the numerical solutions.

In the next section, we introduce the Lugiato-Lefever equation and present the result on wellposedness and regularity of solutions (Theorem 1). It is the cornerstone of the numerical analysis in later chapters. Since the proof is rather long, it is postponed to Section 8. The splitting method for the Lugiato-Lefever equation is introduced in Section 3, and we formulate the error bounds for the global error (Theorem 2) along with the results required for its proof (bounds of the local error in $L^2(\mathbb{T})$ and $H^1(\mathbb{T})$ and stability of the scheme). All following sections are devoted to the proofs of these assertions. In Section 4, we prove stability of the numerical scheme, and we compile a number of auxiliary results. The bounds of the local errors are shown in Sections 5 and 6, respectively, and the proof of the global error bound follows in Section 7.

2 The Lugiato-Lefever equation

The cubic semilinear Schrödinger equation

$$\partial_t u(t, x) = -u(t, x) + i\partial_x^2 u(t, x) + i|u(t, x)|^2 u(t, x) + g(t, x), \quad t > 0 \quad (1a)$$

$$u(0, x) = u_0(x) \quad (1b)$$

on the one-dimensional torus $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$ is known as the Lugiato-Lefever equation in physics and electronic engineering. The terms $-u(t, x)$ and $g(t, x)$ model damping and external forcing, respectively, and do not appear in the “classical” NLS. Clearly, these terms destroy the Hamiltonian structure, and in general the energy, momentum and norm of the solution do not remain constant in time. Solely the one-dimensional torus is considered, because this is the relevant setting for modeling frequency comb generation; cf. [CY10] and [HHR⁺12].

The evolution equation (1) is considered on $L^2(\mathbb{T})$, i.e. on the Hilbert space

of square integrable functions with the inner product

$$\langle v, w \rangle = \int_{\mathbb{T}} v(x) \overline{w(x)} \, dx, \quad v, w \in L^2(\mathbb{T})$$

and induced norm $\|v\|_{L^2} = \sqrt{\langle v, v \rangle}$. The Sobolev space of all functions $v : \mathbb{T} \rightarrow \mathbb{C}$ with partial derivatives up to order $k \in \mathbb{N}_0$ in $L^2(\mathbb{T})$ is denoted by $H^k(\mathbb{T})$. For every k , $H^k(\mathbb{T})$ is a Hilbert space with norm

$$\|v\|_{H^k}^2 = \sum_{j=0}^k \|\partial_x^j v\|_{L^2}^2.$$

In particular, we identify $H^0(\mathbb{T}) = L^2(\mathbb{T})$. We further define the Sobolev spaces of negative order by duality, i.e., we set

$$H^{-k}(\mathbb{T}) := H^k(\mathbb{T})^* \quad \text{with norm} \quad \|\varphi\|_{H^{-k}} = \sup\{|\varphi(v)| \mid \|v\|_{H^k} \leq 1\}$$

for $k \in \mathbb{N}$. We assume the regularity

$$u_0 \in H^4(\mathbb{T}) \quad \text{and} \quad g \in C^{2-j}([0, T]; H^j(\mathbb{T})), \quad j = 0, 1, 2, \quad (2)$$

for the initial data u_0 and the forcing g , respectively, where $T > 0$ is fixed. It will be shown in Theorem 1 that these assumptions guarantee the global existence and uniqueness of a sufficiently smooth solution.

Henceforth, we will usually omit the space variable and write $u(t)$ instead of $u(t, x)$, and so on. Throughout the paper, $C > 0$ and $C(\cdot) > 0$ denote universal constants, possibly taking different values at various appearances. The notation $C(\cdot)$ means that the constant depends only on the values specified in the brackets.

2.1 Analytic setting

For $v \in H^1(\mathbb{T})$ and $w \in L^2(\mathbb{T})$, the Sobolev embedding $H^{j+1}(\mathbb{T}) \hookrightarrow C^j(\mathbb{T})$ for $j \in \mathbb{N}_0$ implies the inequality

$$\|vw\|_{L^2} \leq \|v\|_{L^\infty} \|w\|_{L^2} \leq C \|v\|_{H^1} \|w\|_{L^2}. \quad (3)$$

For every $k \in \mathbb{N}_0$ we define

$$k^* = \max\{1, k\}. \quad (4)$$

Then, for $v \in H^{k^*}(\mathbb{T})$ and $w \in H^k(\mathbb{T})$ the bound

$$\|vw\|_{H^k} \leq C \|v\|_{H^{k^*}} \|w\|_{H^k}, \quad k \geq 0 \quad (5)$$

follows from (3). For every $k \in \mathbb{Z}$, the operator

$$A := i\partial_x^2 - I \quad \text{with domain} \quad H^{k+2}(\mathbb{T})$$

generates a strongly continuous group e^{tA} in $H^k(\mathbb{T})$. The level k of regularity is not expressed in our notation since the respective operators are restrictions of each other.

2.2 Existence and uniqueness of solutions

In the analysis of the numerical method presented below, it is assumed that a unique solution of (1) exists in $H^4(\mathbb{T})$ on $[0, T]$, under assumption (2). In the next theorem we construct the desired (global) solution, and we also establish additional results on the longtime behavior (on the time interval \mathbb{R}_+). We refer to e.g. [Caz03] for local wellposedness in such regularity classes and for related results on global wellposedness without forcing.

Theorem 1. *Let assumption (2) hold. Then, (1) has a unique solution*

$$u \in C(\mathbb{R}_+; H^4(\mathbb{T})) \cap C^1(\mathbb{R}_+; H^2(\mathbb{T})) \cap C^2(\mathbb{R}_+; L^2(\mathbb{T})). \quad (6)$$

Under additional assumptions on g , the following bounds hold.

- (a) *If $g \in L^\infty(\mathbb{R}_+; L^2(\mathbb{T}))$, then $\|u(t)\|_{L^2} \leq C$ for $t \geq 0$.*
- (b) *If $g, \partial_t g \in L^\infty(\mathbb{R}_+; L^2(\mathbb{T}))$, then $\|u(t)\|_{H^1} \leq C\sqrt{1+t}$ for $t \geq 0$.*
- (c) *If $g \in L^2(\mathbb{R}_+; L^2(\mathbb{T}))$, then $u \in L^2(\mathbb{R}_+; L^2(\mathbb{T}))$.*
- (d) *If $g \in H^1(\mathbb{R}_+; L^2(\mathbb{T}))$, then $\|u(t)\|_{H^1} \leq C$ for $t \geq 0$.*

Here C only depends on $\|u_0\|_{L^2}$ in (a) and on $\|u_0\|_{H^1}$ in (b) and (d), as well as on the respective norms of g .

The proof of this theorem is postponed to Section 8.

3 Strang splitting for the Lugiato-Lefever equation

In order to formulate a numerical method for (1), it is convenient to define the nonlinear mapping

$$B : L^2(\mathbb{T}) \longrightarrow L^1(\mathbb{T}), \quad B(w) = i|w|^2.$$

If $w \in H^1(\mathbb{T})$, then $B(w) \in L^\infty(\mathbb{T})$ due to the Sobolev embedding $H^1(\mathbb{T}) \hookrightarrow L^\infty(\mathbb{T})$. For a fixed $w \in H^1(\mathbb{T})$, the function $x \mapsto B(w)(x) = i|w(x)|^2$ will be identified with the multiplication operator

$$B(w) : L^2(\mathbb{T}) \rightarrow L^2(\mathbb{T}), \quad B(w)v = i|w|^2v$$

which generates a unitary group $(e^{tB(w)})_{t \in \mathbb{R}}$ on $L^2(\mathbb{T})$. For $v \in H^{k^*}(\mathbb{T})$ and $w \in H^k(\mathbb{T})$ it follows from (5) that

$$\|B(w)v\|_{H^k} \leq C\|v\|_{H^{k^*}}^2 \|w\|_{H^k}, \quad k \geq 0. \quad (7)$$

As in 2.1 we let $A := i\partial_x^2 - I$. Then, the Lugiato-Lefever equation (1) reads

$$\partial_t u = Au + B(u)u + g, \quad (8a)$$

$$u(0) = u_0. \quad (8b)$$

The solution is supposed to be approximated on the time-interval $[0, T]$ for $T > 0$.

3.1 ϕ -functions

In the construction and analysis of the splitting method for (1) we use the operator-valued functions $\phi_j(tA)$ defined by

$$\phi_j(tA)v = \int_0^1 \frac{\theta^{j-1}}{(j-1)!} e^{(1-\theta)tA} v \, d\theta, \quad j \in \mathbb{N}, \quad \phi_0(tA)v = e^{tA}v, \quad (9)$$

cf. [HO10]. For every $j, k \in \mathbb{N}_0$ and $t \geq 0$, the operator $\phi_j(tA) : H^k(\mathbb{T}) \rightarrow H^k(\mathbb{T})$ is bounded, and

$$\|\phi_j(tA)v\|_{H^k(\mathbb{T})} \leq \frac{1}{j!} \|v\|_{H^k(\mathbb{T})} \quad \text{for all } v \in H^k(\mathbb{T}).$$

For every $v \in L^2(\mathbb{T})$ and $t > 0$, the recurrence relation

$$\phi_{j+1}(tA)v = (tA)^{-1} \left(\phi_j(tA)v - \frac{1}{j!} v \right), \quad j \in \mathbb{N}_0 \quad (10)$$

follows from (9) via integration by parts. This recursion yields the Taylor expansions

$$e^{tA}v = \phi_0(tA)v = \sum_{k=0}^{m-1} \frac{t^k}{k!} A^k v + (tA)^m \phi_m(tA)v \quad (11)$$

for $m \in \mathbb{N}$ and $v \in D(A^m)$. Similar to (9) we define for $j \in \mathbb{N}$, $w \in H^1(\mathbb{T})$ and $v \in L^2(\mathbb{T})$

$$\phi_j(tB(w))v = \int_0^1 \frac{\theta^{j-1}}{(j-1)!} e^{(1-\theta)tB(w)} v \, d\theta, \quad \phi_0(tB(w))v = e^{tB(w)}v. \quad (12)$$

Equation (11) still holds if A is replaced by $B(w)$.

3.2 Time-integration scheme

Strang splitting methods for (8) are based on the observation that solving each of the two sub-problems

$$\partial_t v(t) = Av(t) + g(t), \quad (13)$$

$$\partial_t w(t) = B(w(t))w(t), \quad (14)$$

is much easier than solving (8a). Let $t_n = n\tau$ with step-size $\tau > 0$. Applying the variation-of-constants formula to (13) yields

$$v(t_{n+1}) = e^{\tau A}v(t_n) + \int_0^\tau e^{(\tau-s)A}g(t_n + s) \, ds.$$

After $s \mapsto g(t_n + s)$ has been approximated by the linear interpolation

$$s \mapsto g(t_n) + s \frac{g(t_{n+1}) - g(t_n)}{\tau},$$

the integral can be computed analytically via integration by parts, and we obtain the exponential trapezoidal rule

$$v_{n+1} = e^{\tau A} v_n + \tau \left(\phi_1(\tau A) g(t_n) + \phi_2(\tau A) (g(t_{n+1}) - g(t_n)) \right) \quad (15)$$

which yields approximations $v_n \approx v(t_n)$ to the solution of (13); cf. [HO10].

The sub-problem (14) can even be solved exactly: Since

$$\partial_t (|w(t)|^2) = 2\operatorname{Re}(\bar{w}(t)\partial_t w(t)) = 2\operatorname{Re}(i|w(t)|^4) = 0,$$

it follows that $|w(t)| = |w(0)|$ is time invariant, and hence the solution of (14) is given explicitly by

$$w(t) = e^{tB(w(0))} w(0).$$

This is a well-known fact; see [Fao12]. Approximations $u_n \approx u(t_n)$ to the solution of the *full* problem (8) can now be computed recursively with the Strang splitting

$$u_n^+ = e^{\tau B(u_n)/2} u_n, \quad (16a)$$

$$u_n^* = e^{\tau A} u_n^+ + \tau \left(\phi_1(\tau A) g(t_n) + \phi_2(\tau A) (g(t_{n+1}) - g(t_n)) \right), \quad (16b)$$

$$u_{n+1} = e^{\tau B(u_n^*)/2} u_n^*. \quad (16c)$$

Every time-step $u_n \mapsto u_{n+1}$ of the Strang splitting consists of three sub-steps. First, (14) is solved over the interval $[t_n, t_n + \frac{\tau}{2}]$ with initial data $w(t_n) = u_n$, which yields an update $u_n^+ = w(t_n + \frac{\tau}{2})$. Then, one step of the exponential trapezoidal rule (15) with step-size τ and $v_n = u_n^+$ is carried out, which turns u_n^+ into u_n^* . Finally, (14) is propagated over the interval $[t_n + \frac{\tau}{2}, t_{n+1}]$, which gives the new approximation $u_{n+1} \approx u(t_{n+1})$. Note that for $A = i\Delta$ and $g(t) \equiv 0$, (16) reduces to the method considered in [Lub08] for solving the NLS in absence of damping and forcing.

For every $\theta \geq 0$, the result after $n \in \mathbb{N}_0$ steps of the Strang splitting (16) with step-size $\tau > 0$ starting at time θ with initial data z will be denoted by

$$\Phi_{\tau, \theta}^n(z).$$

If $n = 1$, then we simply write $\Phi_{\tau, \theta}(z)$ instead of $\Phi_{\tau, \theta}^1(z)$. For any $\tau > 0$ and $n \in \mathbb{N}$, the relations

$$\Phi_{\tau, \theta}^0(z) = z, \quad \Phi_{\tau, 0}^n(z) = \Phi_{\tau, t_{n-1}}(\Phi_{\tau, 0}^{n-1}(z)) = \Phi_{\tau, t_1}^{n-1}(\Phi_{\tau, 0}(z))$$

follow directly from the definition. In addition to the numerical flow $\Phi_{\tau, \theta}^n(z)$, we also consider the exact flow given by the exact solution of (8a)

$$t \mapsto \Psi_{t, \theta}(z)$$

with initial data $u(\theta) = z$ at time θ .

For the discretization of space the spectral collocation method can be used, i.e. the solution $u(t) = u(t, x)$ is approximated by a trigonometric polynomial which satisfies (1a) in $m \in \mathbb{N}$ equidistant collocation points $x_k = 2\pi k/m$; see [Fao12] for details. If p is such a trigonometric polynomial, then $e^{\tau A}p$ can be easily computed by means of the fast Fourier transform. Terms like $e^{\tau B(p)/2}p$ are approximated with a trigonometric polynomial which interpolates the values $e^{i\tau|p(x_k)|^2/2}p(x_k)$ in the collocation points. Hence, all terms in (16) can be evaluated quickly at low computational costs. In this paper, however, only the semidiscretization in time with the Strang splitting (16) and without any approximation in space will be analyzed.

3.3 Error analysis: main results

Our goal is to prove that the Strang splitting converges with order 1 in $H^1(\mathbb{T})$ and with order 2 in $L^2(\mathbb{T})$ on bounded time-intervals. In order to state our results, we define the abbreviations

$$\begin{aligned} m_u^k &:= \sup_{t \in [0, T]} \|u(t)\|_{H^k}, & m_g^k &:= \sup_{t \in [0, T]} \|g(t)\|_{H^k}, \\ m_{g'}^k &:= \sup_{t \in [0, T]} \|\partial_t g(t)\|_{H^k}, & m_{g''}^k &:= \sup_{t \in [0, T]} \|\partial_t^2 g(t)\|_{H^k}. \end{aligned}$$

Observe that for $k \leq 4$ the number m_u^k is finite by Theorem 1 and assumption (2). An inspection of the proof of Theorem 1 shows that m_u^4 only depends on the norms of u_0 and g in the spaces involved in (2). For a solution $u(t) \in H^4(\mathbb{T})$ of (8) we immediately obtain the estimates

$$\sup_{t \in [0, T]} \|\partial_t u(t)\|_{H^k} \leq C(m_u^{k+2}, m_g^k), \quad \text{for } 0 \leq k \leq 2, \quad (17)$$

$$\sup_{t \in [0, T]} \|\partial_t^2 u(t)\|_{L^2} \leq C(m_u^4, m_g^2, m_{g'}^0). \quad (18)$$

The following theorem is the main result of the error analysis.

Theorem 2. *Let $u(t) = \Psi_{t,0}(u_0)$ be the exact solution of (8) and assume that the initial data u_0 and the forcing g have the regularity (2). Then, the global error of the splitting method (16) is bounded by*

$$\|\Phi_{\tau,0}^n(u_0) - u(t_n)\|_{H^1} \leq \tau C(T, m_u^3, m_g^1, m_{g'}^1), \quad (19)$$

$$\|\Phi_{\tau,0}^n(u_0) - u(t_n)\|_{L^2} \leq \tau^2 C(T, m_u^4, m_g^2, m_{g'}^1, m_{g''}^0) \quad (20)$$

for all $n \in \mathbb{N}$ with $t_n = n\tau \leq T$ and sufficiently small $\tau > 0$.

Theorem 2 is shown in Section 7. In (51) and (53) we give upper bounds for the step-size τ , but we remark that this step-size restriction is typically too pessimistic in practice. The outline of the proof is taken from [Lub08]. The first two ingredients are bounds for the local error of (16) in $H^1(\mathbb{T})$ and in $L^2(\mathbb{T})$, respectively. They are established in Section 5 and 6.

Theorem 3 (Local error in $H^1(\mathbb{T})$). *Let $n \in \mathbb{N}$ with $t_{n+1} = t_n + \tau \leq T$. If $u(t_n) \in H^3(\mathbb{T})$ and if $g \in C^1([0, T]; H^1(\mathbb{T}))$, then the error after one step of the splitting method (16) is bounded by*

$$\|\Phi_{\tau, t_n}(u(t_n)) - \Psi_{\tau, t_n}(u(t_n))\|_{H^1} \leq \tau^2 C(m_u^3, m_g^1, m_{g'}^1).$$

Theorem 4 (Local error in $L^2(\mathbb{T})$). *Let $n \in \mathbb{N}$ with $t_{n+1} = t_n + \tau \leq T$. Under assumption (2) the error after one step of the splitting method (16) is bounded by*

$$\|\Phi_{\tau, t_n}(u(t_n)) - \Psi_{\tau, t_n}(u(t_n))\|_{L^2} \leq \tau^3 C(m_u^4, m_g^2, m_{g'}^1, m_{g''}^0).$$

The error bound for the global error of (16) is obtained by combining the bounds for the local error with the following stability result, proved in Section 4.

Theorem 5 (Stability). *Let $n \in \mathbb{N}$ with $t_{n+1} = t_n + \tau \leq T$. For $v, w \in H^1(\mathbb{T})$ with $\|v\|_{H^1} \leq M$ and $\|w\|_{H^1} \leq M$, the splitting method (16) satisfies*

$$\|\Phi_{\tau, t_n}(v) - \Phi_{\tau, t_n}(w)\|_{H^k} \leq e^{C \cdot (M_*^2 + M^2 - 1)\tau} \|v - w\|_{H^k}, \quad k = 0, 1 \quad (21)$$

with constant

$$M_* = e^{(CM^2 - 1)\tau} M + \tau C m_g^1. \quad (22)$$

4 Stability and auxiliary results

Now we state three lemmas which will be used frequently throughout the paper. The first lemma asserts a stability result for the mapping $v \mapsto e^{tB(v)}v$. As before in (4), we let $k^* = \max\{1, k\}$.

Lemma 1. *If $v, w \in H^{k^*}(\mathbb{T})$ with $\|v\|_{H^{k^*}} \leq M$ and $\|w\|_{H^{k^*}} \leq M$ for some $k \in \mathbb{N}_0$, then*

$$\|e^{tB(v)}v - e^{tB(w)}w\|_{H^k} \leq e^{CM^2t} \|v - w\|_{H^k}, \quad t \geq 0, \quad (23a)$$

$$\|e^{tB(v)}v\|_{H^k} \leq M e^{CM^2t}, \quad t \geq 0. \quad (23b)$$

Note that for the stability in $L^2(\mathbb{T})$ (i.e. $k = 0$ and $k^* = 1$) the functions v and w have to belong to $H^1(\mathbb{T})$.

Proof. The proof uses ideas of [Lub08]. Let $k \in \mathbb{N}_0$ and let $v, w \in H^{k^*}(\mathbb{T})$ with $\|v\|_{H^{k^*}} \leq M$ and $\|w\|_{H^{k^*}} \leq M$. Then, the functions $x(t) = e^{tB(v)}v$ and $y(t) = e^{tB(w)}w$ are the solutions of the initial value problems

$$\begin{aligned} x'(t) &= B(v)x(t), & x(0) &= v, & t &\geq 0, \\ y'(t) &= B(w)y(t), & y(0) &= w, & t &\geq 0, \end{aligned} \quad (24)$$

respectively, cf. Section 3.2. The inequality (5) implies that $B(v) \in H^k(\mathbb{T})$ and hence $e^{tB(v)} \in H^k(\mathbb{T})$, and applying (5) once again shows that $x(t) = e^{tB(v)}v \in H^k(\mathbb{T})$ for every $t \in [0, T]$. The same arguments yield that $y(t) \in H^k(\mathbb{T})$ for every $t \in [0, T]$.

First, we examine $\|x(t)\|_{H^k}$. From (7) we derive the estimate

$$\|B(v)x(t)\|_{H^k} \leq CM^2\|x(t)\|_{H^k}, \quad t \geq 0, \quad (25)$$

and hence

$$\|x(t)\|_{H^k} \leq \|x(0)\|_{H^k} + \int_0^t \|B(v)x(s)\|_{H^k} \, ds \leq M + CM^2 \int_0^t \|x(s)\|_{H^k} \, ds.$$

Gronwall's lemma now yields

$$\|x(t)\|_{H^k} \leq Me^{CM^2t}, \quad (26)$$

which proves (23b). In order to show (23a), we consider the difference

$$\begin{aligned} B(v)x(t) - B(w)y(t) &= i|v|^2x(t) - i|w|^2y(t) \\ &= i(v-w)\bar{v}x(t) + iw(\bar{v}-\bar{w})x(t) + iw\bar{w}(x(t)-y(t)). \end{aligned}$$

Using also (5) and (26), we derive

$$\begin{aligned} \|B(v)x(t) - B(w)y(t)\|_{H^k} &\leq C[2M\|x(t)\|_{H^k} \|v-w\|_{H^k} + M^2\|x(t)-y(t)\|_{H^k}] \\ &\leq C[2M^2e^{CM^2t}\|v-w\|_{H^k} + M^2\|x(t)-y(t)\|_{H^k}]. \end{aligned}$$

The equations (24) thus imply

$$\begin{aligned} \|x(t) - y(t)\|_{H^k} &\leq \|v-w\|_{H^k} + \int_0^t \|B(v)x(s) - B(w)y(s)\|_{H^k} \, ds \\ &\leq \left(1 + 2CM^2 \int_0^t e^{CM^2s} \, ds\right) \|v-w\|_{H^k} \\ &\quad + CM^2 \int_0^t \|x(s) - y(s)\|_{H^k} \, ds \end{aligned}$$

for $t \geq 0$. Since $0 \leq (e^{CM^2t} - 1)^2$ yields $2e^{CM^2t} - 1 \leq e^{2CM^2t}$, it follows that

$$1 + 2CM^2 \int_0^t e^{CM^2s} \, ds = 1 + 2(e^{CM^2t} - 1) \leq e^{2CM^2t}.$$

Applying Gronwall's lemma once again, we arrive at

$$\|e^{tB(v)}v - e^{tB(w)}w\|_{H^k} = \|x(t) - y(t)\|_{H^k} \leq e^{\widehat{C}M^2t}\|v-w\|_{H^k}$$

with $\widehat{C} = 3C$. □

The next lemma concerns technical estimates regarding the quantity u^* in the splitting method (16).

Lemma 2. For $n \in \mathbb{N}_0$, a given v and $\tau \in [0, T]$ we define

$$v^*(\tau) = e^{\tau A} e^{\tau B(v)/2} v + \tau \left(\phi_1(\tau A) g(t_n) + \phi_2(\tau A) (g(t_n + \tau) - g(t_n)) \right). \quad (27)$$

(i) If $v, u_n \in H^1(\mathbb{T})$ with $\|v\|_{H^1}, \|u_n\|_{H^1} \leq M$, then

$$\|v^*(\tau)\|_{H^k} \leq e^{(CM^2-1)\tau} M + \tau C m_g^k \quad \text{for } k = 0, 1.$$

(ii) If $k \in \{0, 1\}$ and $v, u_n \in H^{k+2}(\mathbb{T})$ with $\|v\|_{H^{k+2}}, \|u_n\|_{H^{k+2}} \leq M$, then

$$\|\partial_\tau v^*(\tau)\|_{H^k} \leq C(T, M, m_g^k, m_{g'}^k).$$

(iii) If $v, u_n \in H^4(\mathbb{T})$ with $\|v\|_{H^4}, \|u_n\|_{H^4} \leq M$, then

$$\|\partial_\tau^2 v^*(\tau)\|_{L^2} \leq C(T, M, m_g^2, m_{g'}^0, m_{g''}^0).$$

Remark. If $v = u_n$, then $v^*(\tau) = u_n^*$ defined in (16b). In the error analysis below, however, Lemma 2 will sometimes also be applied with $v = u(t_n)$.

Proof. The first assertion follows from Lemma 1 and the boundedness of the operators $\phi_j(\tau A)$. For the proof of (ii) and (iii), it is useful to represent the derivative $\partial_\tau \phi_j(\tau A)$ in terms of $\phi_{j-1}(\tau A)$ and $\phi_j(\tau A)$: If $v \in D(A)$, then (9) yields

$$\begin{aligned} \partial_\tau \phi_j(\tau A) v &= \int_0^1 \frac{\theta^{j-1}}{(j-1)!} (1-\theta) A e^{(1-\theta)\tau A} v \, d\theta \\ &= \int_0^1 \frac{\theta^{j-1}}{(j-1)!} A e^{(1-\theta)\tau A} v \, d\theta - j \int_0^1 \frac{\theta^j}{j!} A e^{(1-\theta)\tau A} v \, d\theta \\ &= (\phi_j(\tau A) - j \phi_{j+1}(\tau A)) A v, \end{aligned} \quad (28)$$

and with (10) we obtain

$$\begin{aligned} \partial_\tau \phi_0(\tau A) v &= e^{\tau A} A v, & v \in D(A), \quad \tau \geq 0, \\ \partial_\tau \phi_j(\tau A) v &= \frac{1}{\tau} (\phi_{j-1}(\tau A) - j \phi_j(\tau A)) v, & v \in L^2(\mathbb{T}), \quad j > 0, \quad \tau > 0. \end{aligned}$$

Now (ii) and (iii) can be shown with straightforward calculations using (7), Lemma 1, the boundedness of $\phi_j(\tau A)$, and the fact that $\tau \leq T$. \square

After these preparations we are ready to prove stability of the Strang splitting scheme (16). In order to simplify notation (in particular in Sections 5 and 6), we define

$$B_{1/2}(u) = \frac{i}{2} |u|^2 = \frac{1}{2} B(u). \quad (29)$$

Proof of Theorem 5. Let $v, w \in H^1(\mathbb{T})$ with $\|v\|_{H^1} \leq M$ and $\|w\|_{H^1} \leq M$. As in (16), we define

$$\begin{aligned} v^* &= e^{\tau A} e^{\tau B_{1/2}(v)} v + \tau \left(\phi_1(\tau A) g(t_n) + \phi_2(\tau A) (g(t_n + \tau) - g(t_n)) \right), \\ w^* &= e^{\tau A} e^{\tau B_{1/2}(w)} w + \tau \left(\phi_1(\tau A) g(t_n) + \phi_2(\tau A) (g(t_n + \tau) - g(t_n)) \right). \end{aligned}$$

According to Lemma 2, we have $\|v^*\|_{H^1} \leq M_*$ and $\|w^*\|_{H^1} \leq M_*$ with M_* defined in (22). Applying Lemma 1 twice results in the estimates

$$\begin{aligned} \|\Phi_{\tau, t_n}(v) - \Phi_{\tau, t_n}(w)\|_{H^k} &= \|e^{\tau B_{1/2}(v^*)} v^* - e^{\tau B_{1/2}(w^*)} w^*\|_{H^k} \\ &\leq e^{CM_*^2 \tau} \|v^* - w^*\|_{H^k} \\ &= e^{CM_*^2 \tau} \|e^{\tau A} e^{\tau B_{1/2}(v)} v - e^{\tau A} e^{\tau B_{1/2}(w)} w\|_{H^k} \\ &\leq e^{(CM_*^2 - 1)\tau} \|e^{\tau B_{1/2}(v)} v - e^{\tau B_{1/2}(w)} w\|_{H^k} \\ &\leq e^{C \cdot (M_*^2 + M^2 - 1)\tau} \|v - w\|_{H^k} \end{aligned}$$

for $k \in \{0, 1\}$. □

The last lemma in this subsection will be useful in the proofs of Theorems 3 and 4.

Lemma 3. For a given $n \in \mathbb{N}$ and $\tau \in [0, T]$ let

$$b(\tau) = B(u(t_n + \tau)) - B(v^*(\tau)),$$

where $v^*(\tau)$ is defined by (27) with $v = u(t_n)$. Under the assumption (2), we have

$$b(\tau) = \int_0^\tau \partial_\tau b(s) ds = \int_0^\tau \int_0^s \partial_\tau^2 b(r) dr ds.$$

Proof. The fundamental theorem of calculus gives

$$b(\tau) = b(0) + \int_0^\tau \partial_\tau b(s) ds = b(0) + \tau \partial_\tau b(0) + \int_0^\tau \int_0^s \partial_\tau^2 b(r) dr ds.$$

As $v^*(0) = v = u(t_n)$ by assumption, it is clear that $b(0) = 0$. Hence, we only have to show that $\partial_\tau b(0) = 0$. By definition, we have

$$\partial_\tau b(\tau) = 2i \operatorname{Re} (\bar{u}(t_n + \tau) \partial_\tau u(t_n + \tau)) - 2i \operatorname{Re} (\bar{v}^*(\tau) \partial_\tau v^*(\tau)).$$

Since $u(t_n) = v = v^*(0)$,

$$\begin{aligned} \partial_\tau u(t_n + \tau) \Big|_{\tau=0} &= Au(t_n) + B(u(t_n))u(t_n) + g(t_n), \\ \partial_\tau v^*(\tau) \Big|_{\tau=0} &= Av + B_{1/2}(v)v + g(t_n), \end{aligned}$$

we deduce

$$\begin{aligned}\partial_\tau b(0) &= 2i\operatorname{Re}(\bar{u}(t_n)\partial_\tau u(t_n)) - 2i\operatorname{Re}(\bar{u}(t_n)\partial_\tau v^*(0)) \\ &= 2i\operatorname{Re}(\bar{u}(t_n)B(u(t_n))u(t_n)) - 2i\operatorname{Re}(\bar{u}(t_n)B_{1/2}(u(t_n))u(t_n)) = 0\end{aligned}$$

because $\bar{u}(t_n)B(u(t_n))u(t_n) = i|u(t_n)|^4$ and $\bar{u}(t_n)B_{1/2}(u(t_n))u(t_n) = \frac{i}{2}|u(t_n)|^4$ are both purely imaginary functions; cf. [CKT07]. \square

5 Local error in $H^1(\mathbb{T})$: Proof of Theorem 3

Without loss of generality we assume that $n = 0$, i.e. $u(t_n) = u(0) = u_0$ and $\Psi_{\tau, t_n}(u(t_n)) = \Psi_{\tau, 0}(u_0) = u(\tau)$.

Step 1. The variation-of-constants formula yields the representation

$$u(\tau) = e^{\tau A}u_0 + \int_0^\tau e^{(\tau-s)A} \left(B(u(s))u(s) + g(s) \right) ds$$

of the exact solution of (8). Substituting the formula a second time for $u(s)$, we obtain

$$u(\tau) = e^{\tau A}u_0 + I_1 + I_2 + R_1, \quad (30)$$

where we set

$$I_1 = \int_0^\tau e^{(\tau-s)A} B(u(s))e^{sA}u_0 ds, \quad (31)$$

$$I_2 = \int_0^\tau e^{(\tau-s)A} g(s) ds, \quad (32)$$

$$R_1 = \int_0^\tau \int_0^s e^{(\tau-s)A} B(u(s))e^{(s-\sigma)A} [B(u(\sigma))u(\sigma) + g(\sigma)] d\sigma ds.$$

Using (7), it can be shown that

$$\|R_1\|_{H^1} \leq \tau^2 C(m_u^1, m_g^1). \quad (33)$$

The approximation $u_1 = \Phi_{\tau, 0}(u_0)$ of the numerical method after one step reads

$$\begin{aligned}u_1 &= e^{\tau B_{1/2}(u_0^*)} e^{\tau A} e^{\tau B_{1/2}(u_0)} u_0 \\ &\quad + \tau e^{\tau B_{1/2}(u_0^*)} \left(\phi_1(\tau A)g(0) + \phi_2(\tau A)(g(\tau) - g(0)) \right)\end{aligned} \quad (34)$$

with $B_{1/2}(\cdot)$ defined in (29) and

$$u_0^* = e^{\tau A} e^{\tau B_{1/2}(u_0)} u_0 + \tau \left(\phi_1(\tau A)g(0) + \phi_2(\tau A)(g(\tau) - g(0)) \right).$$

We consider u_0^* as a function of $\tau > 0$. Using further the expansion

$$e^{\tau B_{1/2}(u_0)} v = \sum_{k=0}^{m-1} \frac{\tau^k}{k!} B_{1/2}^k(u_0)v + \tau^m B_{1/2}^m(u_0)\phi_m(\tau B_{1/2}(u_0))v$$

with $m \in \{1, 2\}$, see (11), we derive

$$u_1 = e^{\tau A} u_0 + T_1 + T_2 + R_2, \quad (35)$$

with

$$T_1 = \tau \left(B_{1/2}(u_0^*) e^{\tau A} + e^{\tau A} B_{1/2}(u_0) \right) u_0, \quad (36)$$

$$T_2 = \tau \left(\phi_1(\tau A) g(0) + \phi_2(\tau A) (g(\tau) - g(0)) \right), \quad (37)$$

$$\begin{aligned} R_2 = \tau^2 & \left[e^{\tau A} B_{1/2}^2(u_0) \phi_2(\tau B_{1/2}(u_0)) u_0 + B_{1/2}(u_0^*) e^{\tau A} B_{1/2}(u_0) \phi_1(\tau B_{1/2}(u_0)) u_0 \right. \\ & + B_{1/2}^2(u_0^*) \phi_2(\tau B_{1/2}(u_0^*)) e^{\tau A} e^{\tau B_{1/2}(u_0)} u_0 \\ & \left. + B_{1/2}(u_0^*) \phi_1(\tau B_{1/2}(u_0^*)) \left(\phi_1(\tau A) g(0) + \phi_2(\tau A) (g(\tau) - g(0)) \right) \right]. \end{aligned}$$

Estimate (7) and Lemmas 1 and 2 imply

$$\|R_2\|_{H^1} \leq \tau^2 C(T, m_u^1, m_g^1). \quad (38)$$

Step 2. We compare the exact solution (30) with the numerical solution (35). Using (33) and (38), we infer

$$\|u(\tau) - u_1\|_{H^1} \leq \|I_1 - T_1\|_{H^1} + \|I_2 - T_2\|_{H^1} + \tau^2 C(m_u^1, m_g^1).$$

Our goal is now to bound the terms $\|I_1 - T_1\|_{H^1}$ and $\|I_2 - T_2\|_{H^1}$. With the abbreviations

$$h_1(s) = e^{(\tau-s)A} B(u(s)) e^{sA} u_0, \quad (39)$$

$$b(\tau) = B(u(\tau)) - B(u_0^*(\tau)), \quad (40)$$

the first term can be represented as $I_1 - T_1 = Q_1 + E_1$, where

$$Q_1 = \int_0^\tau h_1(s) ds - \frac{\tau}{2} (h_1(0) + h_1(\tau)) \quad (41)$$

is the local quadrature error of the trapezoidal rule and

$$E_1 = \frac{\tau}{2} b(\tau) e^{\tau A} u_0 \quad (42)$$

is a remainder term. The order of the trapezoidal rule is two, and hence its local error scales like $O(\tau^3)$ if the integrand is smooth enough. For the proof of Theorem 3, however, the bound

$$\|Q_1\|_{H^1} \leq \tau^2 C \sup_{s \in [0, \tau]} \|\partial_s h_1(s)\|_{H^1} \quad (43)$$

is sufficient. Applying Lemma 3 with $n = 0$ and $t_n = 0$, the remainder term E_1 can be bounded by

$$\|E_1\|_{H^1} \leq \tau^2 C m_u^1 \sup_{s \in [0, \tau]} \|\partial_s b(s)\|_{H^1}.$$

The difference

$$I_2 - T_2 = \int_0^\tau e^{(\tau-s)A} g(s) \, ds - \tau \left(\phi_1(\tau A) g(0) + \phi_2(\tau A) (g(\tau) - g(0)) \right)$$

is the local error of the exponential trapezoidal rule so that

$$\|I_2 - T_2\|_{H^1} \leq \tau^2 C m_{g'}^1,$$

see Theorem 2.7 in [HO10].

Step 3. To complete the proof of Theorem 3, it remains to show that the terms

$$\sup_{s \in [0, \tau]} \|\partial_s h_1(s)\|_{H^1} \quad \text{and} \quad \sup_{s \in [0, \tau]} \|\partial_s b(s)\|_{H^1}$$

are bounded. The equations (7) and (17) yield

$$\sup_{s \in [0, \tau]} \|\partial_s h_1(s)\|_{H^1} \leq C (m_u^3, m_g^1).$$

Finally, in view of (17) and Lemma 2, the term

$$\partial_s b(s) = 2i \left(\operatorname{Re}(\bar{u}(s) \partial_s u(s)) - \operatorname{Re}(\bar{u}_0^*(s) \partial_s u_0^*(s)) \right)$$

can be estimated

$$\sup_{s \in [0, \tau]} \|\partial_s b(s)\|_{H^1} \leq C (T, m_u^3, m_g^1, m_{g'}^1),$$

which completes the proof of Theorem 3. \square

6 Local error in $L^2(\mathbb{T})$: Proof of Theorem 4

To prove the third-order local error bound in $L^2(\mathbb{T})$, we mimic the proof for the second-order bound of the local error in $H^1(\mathbb{T})$. However, we have to expand the analytical solution and the numerical scheme to a higher order. As before, we assume without any loss of generality that $n = 0$ and let $u(\tau) = \Psi_{\tau,0}(u_0)$.

Step 1. We expand the exact solution further by inserting the variation-of-constants formula for $u(\sigma)$ into (30). It follows that

$$u(\tau) = e^{\tau A} u_0 + I_1 + I_2 + I_3 + I_4 + \hat{R}_1,$$

where I_1 and I_2 have been defined in (31) and (32), respectively, and we introduce

$$\begin{aligned} I_3 &= \int_0^\tau \int_0^s e^{(\tau-s)A} B(u(s)) e^{(s-\sigma)A} B(u(\sigma)) e^{\sigma A} u_0 \, d\sigma \, ds, \\ I_4 &= \int_0^\tau \int_0^s e^{(\tau-s)A} B(u(s)) e^{(s-\sigma)A} g(\sigma) \, d\sigma \, ds, \\ \hat{R}_1 &= \int_0^\tau \int_0^s \int_0^\sigma e^{(\tau-s)A} B(u(s)) e^{(s-\sigma)A} \\ &\quad \times B(u(\sigma)) e^{(\sigma-\xi)A} [B(u(\xi))u(\xi) + g(\xi)] \, d\xi \, d\sigma \, ds. \end{aligned}$$

The estimate (7) yields

$$\|\hat{R}_1\|_{L^2} \leq \tau^3 C(m_u^1, m_g^0). \quad (44)$$

Substituting the expansion

$$e^{\tau B_{1/2}(\cdot)} = I + \tau B_{1/2}(\cdot) + \frac{\tau^2}{2} B_{1/2}^2(\cdot) + \tau^3 B_{1/2}^3(\cdot) \phi_3(\tau B_{1/2}(\cdot))$$

into the splitting method (34), we derive

$$u_1 = e^{\tau A} u_0 + T_1 + T_2 + T_3 + T_4 + \hat{R}_2$$

with T_1, T_2 from (36), (37) and

$$\begin{aligned} T_3 &= \frac{\tau^2}{2} \left(B_{1/2}^2(u_0^*) e^{\tau A} + 2B_{1/2}(u_0^*) e^{\tau A} B_{1/2}(u_0) + e^{\tau A} B_{1/2}^2(u_0) \right) u_0 \\ T_4 &= \tau^2 B_{1/2}(u_0^*) \left(\phi_1(\tau A) g(0) + \phi_2(\tau A) (g(\tau) - g(0)) \right) \\ \hat{R}_2 &= \tau^3 \left[e^{\tau A} B_{1/2}^3(u_0) \phi_3(\tau B_{1/2}(u_0)) u_0 + B_{1/2}(u_0^*) e^{\tau A} B_{1/2}^2(u_0) \phi_2(\tau B_{1/2}(u_0)) u_0 \right. \\ &\quad + \frac{1}{2} B_{1/2}^2(u_0^*) e^{\tau A} B_{1/2}(u_0) \phi_1(\tau B_{1/2}(u_0)) u_0 \\ &\quad + B_{1/2}^3(u_0^*) \phi_3(\tau B_{1/2}(u_0^*)) e^{\tau A} e^{\tau B_{1/2}(u_0)} u_0 \\ &\quad \left. + B_{1/2}^2(u_0^*) \phi_2(\tau B_{1/2}(u_0^*)) \left(\phi_1(\tau A) g(0) + \phi_2(\tau A) (g(\tau) - g(0)) \right) \right]. \end{aligned}$$

Inequality (7) and Lemma 1 imply

$$\|\hat{R}_2\|_{L^2} \leq \tau^3 C(m_u^1, m_g^0). \quad (45)$$

Step 2. Comparing the exact solution with the numerical solution and using (44) and (45), we estimate

$$\begin{aligned} \|u(\tau) - u_1\|_{L^2} &\leq \|I_1 - T_1\|_{L^2} + \|I_2 - T_2\|_{L^2} + \|I_3 - T_3\|_{L^2} \\ &\quad + \|I_4 - T_4\|_{L^2} + \tau^3 C(m_u^1, m_g^0). \end{aligned}$$

As before, the terms of the numerical solution are splitted into a suitable quadrature formula and a remainder term. In addition to $h_1(s)$ defined in (39) and $b(\tau)$ defined in (40), we employ the abbreviations

$$\begin{aligned} h_2(s, \sigma) &= e^{(\tau-s)A} B(u(s)) e^{(s-\sigma)A} B(u(\sigma)) e^{\sigma A} u_0, \\ h_3(s) &= e^{(\tau-s)A} B(u(s)) \phi_1(sA) g(0). \end{aligned}$$

We still use the decomposition $I_1 - T_1 = Q_1 + E_1$ with the quadrature error Q_1 from (41) and the remainder E_1 from (42). Since now we aim at a local error in $L^2(\mathbb{T})$ of third order, we replace the error bound (43) by

$$\|Q_1\|_{L^2} \leq \tau^3 C \sup_{s \in [0, \tau]} \|\partial_s^2 h_1(s)\|_{L^2},$$

see [KI66]. Lemma 3 implies

$$\|E_1\|_{L^2} \leq \tau^3 C m_u^1 \sup_{s \in [0, \tau]} \|\partial_s^2 b(s)\|_{L^2}.$$

The difference

$$I_2 - T_2 = \int_0^\tau e^{(\tau-s)A} g(s) \, ds - \tau \left(\phi_1(\tau A) g(0) + \phi_2(\tau A) (g(\tau) - g(0)) \right)$$

is the local error of the exponential trapezoidal rule. We thus acquire

$$\|I_2 - T_2\|_{L^2} \leq \tau^3 C m_{g''}^0,$$

see [HO10]. For the third error term we use the partition $I_3 - T_3 = Q_3 + E_3$ with

$$Q_3 = \int_0^\tau \int_0^s h_2(s, \sigma) \, d\sigma \, ds - \frac{\tau^2}{8} (h_2(0, 0) + 2h_2(\tau, 0) + h_2(\tau, \tau)),$$

$$E_3 = \frac{\tau^2}{8} b(\tau) \left(2e^{\tau A} B(u_0) + [B(u(\tau)) + B(u_0^*(\tau))] e^{\tau A} \right) u_0,$$

and $b(\tau)$ defined in (40). We identify Q_3 as the error of a cubature formula which integrates constant functions exactly. It follows

$$\|Q_3\|_{L^2} \leq C \tau^3 \left(\sup_{\Delta} \|\partial_s h_2(s, \sigma)\|_{L^2} + \sup_{\Delta} \|\partial_\sigma h_2(s, \sigma)\|_{L^2} \right),$$

where Δ is the triangle $0 \leq s \leq \tau, 0 \leq \sigma \leq s$, see p. 362 in [KI66]. From Lemma 3 we infer

$$\|E_3\|_{L^2} \leq \tau^3 C m_u^1 \sup_{s \in [0, \tau]} \|\partial_s b(s)\|_{L^2}.$$

The fourth term is decomposed into three parts

$$I_4 - T_4 = E_4^1 + E_4^2 + E_4^3$$

given by

$$E_4^1 = \int_0^\tau e^{(\tau-s)A} B(u(s)) F(s) \, ds,$$

$$F(s) = \int_0^s e^{(s-\sigma)A} g(\sigma) \, d\sigma - s \phi_1(sA) g(0),$$

$$E_4^2 = \int_0^\tau s h_3(s) \, ds - \tau^2 B_{1/2}(u_0^*) \phi_1(\tau A) g(0),$$

$$E_4^3 = -\tau^2 B_{1/2}(u_0^*) \phi_2(\tau A) (g(\tau) - g(0)).$$

Since $F(s)$ is the local error of the exponential Euler rule, we can estimate

$$\sup_{s \in [0, \tau]} \|F(s)\|_{L^2} \leq \tau^2 C m_{g'}^0,$$

see [HO10]. This inequality and (7) lead to

$$\|E_4^1\|_{L^2} \leq \tau^3 C(m_u^1, m_{g'}^0).$$

Integrating by parts, we calculate

$$\int_0^\tau s h_3(s) \, ds = \frac{\tau^2}{2} h_3(\tau) - \frac{1}{2} \int_0^\tau s^2 \partial_s h_3(s) \, ds,$$

so that

$$E_4^2 = \frac{\tau^2}{2} b(\tau) \phi_1(\tau A) g(0) - \frac{1}{2} \int_0^\tau s^2 \partial_s h_3(s) \, ds.$$

Lemma 3 then yields

$$\|E_4^2\|_{L^2} \leq C \tau^3 \left(m_g^0 \sup_{s \in [0, \tau]} \|\partial_s b(s)\|_{L^2} + \sup_{s \in [0, \tau]} \|\partial_s h_3(s)\|_{L^2} \right).$$

Exploiting the regularity of g , we obtain

$$\|g(\tau) - g(0)\|_{L^2} \leq \tau m_{g'}^0.$$

Estimate (7), Lemma 2 and the boundedness of $\phi_j(\tau A)$ finally imply

$$\|E_4^3\|_{L^2} \leq \tau^3 C(m_u^1, m_g^0, m_{g'}^0).$$

Step 3. To complete the proof of Theorem 4, it remains to show that the terms

$$\begin{aligned} & \sup_{s \in [0, \tau]} \|\partial_s^2 h_1(s)\|_{L^2}, \quad \sup_{\Delta} \|\partial_s h_2(s, \sigma)\|_{L^2}, \quad \sup_{\Delta} \|\partial_\sigma h_2(s, \sigma)\|_{L^2}, \\ & \sup_{s \in [0, \tau]} \|\partial_s h_3(s)\|_{L^2} \quad \text{and} \quad \sup_{s \in [0, \tau]} \|\partial_s^2 b(s)\|_{L^2} \end{aligned}$$

are bounded. Formulas (7), (2) and (28) yield

$$\begin{aligned} & \sup_{s \in [0, \tau]} \|\partial_s^2 h_1(s)\|_{L^2} \leq C(m_u^4, m_g^2, m_{g'}^0), \\ & \sup_{\Delta} \|\partial_s h_2(s, \sigma)\|_{L^2} \leq C(m_u^2, m_g^0), \\ & \sup_{\Delta} \|\partial_\sigma h_2(s, \sigma)\|_{L^2} \leq C(m_u^2, m_g^0), \\ & \sup_{s \in [0, \tau]} \|\partial_s h_3(s)\|_{L^2} \leq C(m_u^2, m_g^2). \end{aligned}$$

We then apply (17), (18) and Lemma 2 to

$$\partial_s^2 b(s) = 2i \left(|\partial_s u(s)|^2 + \operatorname{Re}(\bar{u}(s) \partial_s^2 u(s)) - |\partial_s u_0^*(s)|^2 - \operatorname{Re}(\bar{u}_0^*(s) \partial_s^2 u_0^*(s)) \right),$$

and conclude the last bound

$$\sup_{s \in [0, \tau]} \|\partial_s^2 b(s)\|_{L^2} \leq C(T, m_u^4, m_g^2, m_{g'}^1, m_{g''}^0). \quad \square$$

7 Global error: Proof of Theorem 2

In order to prove the global error estimates in Theorem 2, the local error bounds from Theorems 3 and 4 are combined with the stability result from Theorem 5 in the classical construction known as Lady Windermere's fan, see [HNW08]. However, the stability result (21) can only be applied if the numerical solution $\Phi_{\tau,0}^n(u_0)$ stays bounded in $H^1(\mathbb{T})$ for all $n \in \mathbb{N}$ with $\tau n \leq T$. This condition can be shown by the following induction argument. Let $u_0 \in H^3(\mathbb{T})$ and assume that there is a constant $\widehat{M} > m_u^1$ such that

$$\|\Phi_{\tau,t_\ell}^k(u(t_\ell))\|_{H^1} \leq \widehat{M} \quad \text{for all } \ell \in \mathbb{N}_0, \quad k = 0, \dots, n-1, \quad t_{\ell+k} \leq T. \quad (46)$$

We will prove that

$$\|\Phi_{\tau,t_\ell}^n(u(t_\ell))\|_{H^1} \leq \widehat{M} \quad \text{for all } \ell \in \mathbb{N}_0, \quad t_{\ell+n} \leq T \quad (47)$$

provided that the step-size τ is sufficiently small. Since the argument is the same for all ℓ , we assume that $\ell = 0$ with no loss of generality. Representing $\Phi_{\tau,0}^n(u_0)$ by the telescoping sum

$$\Phi_{\tau,0}^n(u_0) = u(t_n) + \sum_{j=0}^{n-1} \Phi_{\tau,t_j}^{n-j}(u(t_j)) - \Phi_{\tau,t_{j+1}}^{n-j-1}(u(t_{j+1})) \quad (48)$$

with $u(t_n) = \Psi_{t_n,0}(u_0)$ and $u(t_0) = u_0$ gives

$$\|\Phi_{\tau,0}^n(u_0)\|_{H^1} \leq \|u(t_n)\|_{H^1} + \sum_{j=0}^{n-1} \|\Phi_{\tau,t_j}^{n-j}(u(t_j)) - \Phi_{\tau,t_{j+1}}^{n-j-1}(u(t_{j+1}))\|_{H^1}. \quad (49)$$

According to (46), Theorem 5 can be applied and yields for $n-j-1 \geq 1$ that

$$\begin{aligned} & \left\| \Phi_{\tau,t_j}^{n-j}(u(t_j)) - \Phi_{\tau,t_{j+1}}^{n-j-1}(u(t_{j+1})) \right\|_{H^1} \\ &= \left\| \Phi_{\tau,t_{n-1}} \left(\Phi_{\tau,t_j}^{n-j-1}(u(t_j)) \right) - \Phi_{\tau,t_{n-1}} \left(\Phi_{\tau,t_{j+1}}^{n-j-2}(u(t_{j+1})) \right) \right\|_{H^1} \\ &\leq e^{C \cdot (\widehat{M}_*^2 + \widehat{M}^2 - 1)} \left\| \Phi_{\tau,t_j}^{n-j-1}(u(t_j)) - \Phi_{\tau,t_{j+1}}^{n-j-2}(u(t_{j+1})) \right\|_{H^1} \end{aligned} \quad (50)$$

with constant

$$\widehat{M}_* = e^{(C\widehat{M}^2 - 1)\tau} \widehat{M} + \tau C m_g^1,$$

cf. (22). If τ is sufficiently small, then $\widehat{M}_* \leq C\widehat{M}$ so that $e^{C \cdot (\widehat{M}_*^2 + \widehat{M}^2 - 1)} \leq e^{C\widehat{M}^2}$. Applying (50) recursively, we then obtain

$$\begin{aligned} \left\| \Phi_{\tau,t_j}^{n-j}(u(t_j)) - \Phi_{\tau,t_{j+1}}^{n-j-1}(u(t_{j+1})) \right\|_{H^1} &\leq e^{C\widehat{M}^2(n-j-1)\tau} \left\| \Phi_{\tau,t_j}(u(t_j)) - u(t_{j+1}) \right\|_{H^1} \\ &\leq e^{CT\widehat{M}^2} C_{loc} \tau^2 \end{aligned}$$

due to Theorem 3, with the constant C_{loc} from the local error bound. So (49) yields

$$\|\Phi_{\tau,0}^n(u_0)\|_{H^1} \leq \|u(t_n)\|_{H^1} + ne^{CT\widehat{M}^2} C_{loc}\tau^2 \leq m_u^1 + e^{CT\widehat{M}^2} C_{loc}T\tau.$$

If τ is so small that

$$\tau \leq \frac{\widehat{M} - m_u^1}{C_{loc}T} e^{-CT\widehat{M}^2}, \quad (51)$$

then $\|\Phi_{\tau,0}^n(u_0)\|_{H^1} \leq \widehat{M}$, as required.

It is now easy to show the bound for the global error in $L^2(\mathbb{T})$. The telescoping sum (48) yields

$$\|\Phi_{\tau,0}^n(u_0) - u(t_n)\|_{L^2} \leq \sum_{j=0}^{n-1} \|\Phi_{\tau,t_j}^{n-j}(u(t_j)) - \Phi_{\tau,t_j}^{n-j-1}(u(t_{j+1}))\|_{L^2},$$

and with Theorem 5 and Theorem 4 we obtain similar as before

$$\|\Phi_{\tau,0}^n(u_0) - u(t_n)\|_{L^2} \leq ne^{CT\widehat{M}^2} \tilde{C}_{loc}\tau^3 \leq e^{CT\widehat{M}^2} \tilde{C}_{loc}T\tau^2 \quad (52)$$

with \tilde{C}_{loc} denoting the constant from the local error bound in Theorem 4. The bound for the global error in $H^1(\mathbb{T})$ is obtained upon replacing $\|\cdot\|_{L^2}$ by $\|\cdot\|_{H^1}$, τ^p by τ^{p-1} , and \tilde{C}_{loc} by C_{loc} . \square

Remark. According to (52) the global error is small if

$$\tau^2 \ll \frac{1}{\tilde{C}_{loc}T} e^{-CT\widehat{M}^2}. \quad (53)$$

Hence, even if one could avoid the step-size restriction (51) imposed by *stability*, there is still a similar step-size restriction imposed by *accuracy*. Of course, both (51) and (53) are usually too pessimistic in practice. These step-size restrictions are *not* a characteristic property of the equation (1) nor of the splitting method (16). For example, the error bound for the global error of Runge-Kutta methods for solving *ordinary* differential equations is similar to (52); cf. Theorem 3.6 in chapter II in [HNW08].

8 Wellposedness and regularity: Proof of Theorem 1

Before we start with the proof of Theorem 1, we make a few preparations. First, we set

$$F(t, v) = i|v|^2v + g(t)$$

for $t \in \mathbb{R}$ and $v \in H^1(\mathbb{T})$ and define for $a \in (0, \infty]$ and $k \in \mathbb{Z}$ the space

$$E_k := C([0, a]; H^k(\mathbb{T})) \cap C^1([0, a]; H^{k-2}(\mathbb{T})).$$

Second, we state and prove the following lemma which leads to the existence of solutions to (1) with values in $H^1(\mathbb{T})$.

Lemma 4. *a) Let $g \in C^1(\mathbb{R}_+; H^1(\mathbb{T}))$. Then the map $F : \mathbb{R}_+ \times H^1(\mathbb{T}) \rightarrow H^1(\mathbb{T})$ is continuously differentiable. Its derivative with respect to $v \in H^1(\mathbb{T})$ is given by*

$$\partial_v F(t, v)w = i|v|^2 w + 2i \operatorname{Re}(\bar{v}w)v$$

for $t \in \mathbb{R}$ and $v, w \in H^1(\mathbb{T})$. Moreover, F is Lipschitz on bounded subsets.

b) Let $u \in E_1$ and $f(t) = |u(t)|^2 u(t)$ for $t \in [0, a)$. Then $f \in E_1$ and

$$f' = |u|^2 \partial_t u + 2 \operatorname{Re}(\bar{u} \partial_t u)u =: h.$$

Proof of Lemma 4. We omit the well known and straightforward proof of part a). For a given $v \in H^1(\mathbb{T})$ the map $\phi \mapsto v\phi$ is bounded in $H^{-1}(\mathbb{T})$ by duality and (5). In view of (5), we obtain $f \in C([0, a); H^1(\mathbb{T}))$ and $h \in C([0, a); H^{-1}(\mathbb{T}))$. Observe that

$$\begin{aligned} f(t+s) - f(t) &= (u(t+s) - u(t))|u(t+s)|^2 + (u(t+s) - u(t))\bar{u}(t+s)u(t) \\ &\quad + (\bar{u}(t+s) - \bar{u}(t))u(t)^2. \end{aligned}$$

Using $u \in E_1$ and (5), one can differentiate f in $H^{-1}(\mathbb{T})$ with $f' = h$. \square

We are now in a position to prove Theorem 1.

Proof of Theorem 1. Step 1. We first look for a local solution of (1) in E_1 , starting with the integral equation

$$u(t) = e^{tA}u_0 + \int_0^t e^{(t-s)A}F(s, u(s)) \, ds. \quad (54)$$

Theorem 6.1.4 in [Paz83] and Lemma 4 yield a unique maximal solution u in $C([0, a); H^1(\mathbb{T}))$ of (54) for some $a > 0$. Moreover, if a is finite, then $\|u(t)\|_{H^1}$ becomes unbounded as $t \rightarrow a$. Differentiating (54) in $H^{-1}(\mathbb{T})$, one obtains a solution $u \in E_1$ of (1). It is unique since every solution of (1) in E_1 satisfies (54). In the following steps we improve the regularity of u iteratively.

Step 2. Lemma 4 allows us to differentiate the right hand side of (1) in $H^{-3}(\mathbb{T})$ with respect to t , and so there exists $\partial_t^2 u$ in $C([0, a); H^{-3}(\mathbb{T}))$. We set $v = \partial_t u \in E_{-1}$ and $v_0 := i\partial_x^2 u_0 - u_0 + i|u_0|^2 u_0 + g(0)$. Note that $v_0 \in H^2(\mathbb{T})$ by assumption (2). These functions then satisfy

$$\begin{aligned} \partial_t v &= i\partial_x^2 v - v + i|u|^2 v + 2i \operatorname{Re}(\bar{u}v)u + \partial_t g, \\ v(0) &= v_0. \end{aligned} \quad (55)$$

Since u is bounded on $[0, a - \delta] \times \mathbb{T}$ for each $\delta \in (0, a)$, there is a unique function w in $C([0, a]; L^2(\mathbb{T}))$ solving

$$w(t) = e^{tA}v_0 + \int_0^t e^{(t-s)A} \left(i|u(s)|^2w(s) + 2i \operatorname{Re}(\bar{u}(s)w(s))u(s) + \partial_t g(s) \right) ds, \quad (56)$$

see Theorem 6.1.2 in [Paz83]. Because $v \in E_{-1}$ solves (55), it also satisfies (56). As noted above, the multiplication by u or \bar{u} is a bounded operator on $H^{-1}(\mathbb{T})$. If we now subtract the two integral equations for v and w , we derive

$$\|v(t) - w(t)\|_{H^{-1}} \leq C(\delta) \int_0^t \|v(s) - w(s)\|_{H^{-1}} ds \quad \text{for } 0 \leq t \leq a - \delta.$$

Gronwall's inequality implies that $v = w \in C([0, a]; L^2(\mathbb{T}))$; i.e., u is contained in $C^1([0, a]; L^2(\mathbb{T}))$. Equation (1) then yields $u \in C([0, a]; H^2(\mathbb{T}))$ since $(i\partial_x^2 - I)^{-1}$ maps $L^2(\mathbb{T})$ continuously into $H^2(\mathbb{T})$. Moreover, u belongs to $C^2([0, a]; H^{-2}(\mathbb{T}))$ by (55).

Step 3. We next differentiate (55) in $H^{-3}(\mathbb{T})$ with respect to x . For instance, we have $\partial_x(|u|^2v) = |u|^2\partial_x v + 2 \operatorname{Re}(\bar{u}\partial_x u)v$ since

$$\begin{aligned} \langle \partial_x(|u|^2v), \varphi \rangle &= -\langle v, |u|^2\partial_x \varphi \rangle = -\langle v, \partial_x(|u|^2\varphi) \rangle + \langle v, 2 \operatorname{Re}(\bar{u}\partial_x u)\varphi \rangle \\ &= \langle |u|^2\partial_x v, \varphi \rangle + \langle 2 \operatorname{Re}(\bar{u}\partial_x u)v, \varphi \rangle \end{aligned}$$

for $u, \varphi \in H^1(\mathbb{T})$, $v \in L^2(\mathbb{T})$, and the duality between $H^{-1}(\mathbb{T})$ and $H^1(\mathbb{T})$. The function $w := \partial_x v = \partial_t \partial_x u$ then fulfills

$$\begin{aligned} \partial_t w &= i\partial_x^2 w - w + i|u|^2 w + 2i \operatorname{Re}(\bar{u}w)u + \partial_t \partial_x g + h, \\ w(0) &= \partial_x v_0 \in H^1(\mathbb{T}), \end{aligned} \quad (57)$$

where $h := 2i(\operatorname{Re}(u\partial_x \bar{u})v + \operatorname{Re}(v\partial_x \bar{u})u + \operatorname{Re}(v\bar{u})\partial_x u)$ and $\partial_t \partial_x g$ belong to $C([0, a]; L^2(\mathbb{T}))$ by step 2) and assumption (2). As in step 2), we see that the integrated version of (57) has a solution $\varphi \in C([0, a]; L^2(\mathbb{T}))$ and that $w = \varphi$. Hence, $u \in C^1([0, a]; H^1(\mathbb{T}))$.

Step 4. In (55) we have a non-autonomous perturbation given by

$$P(t)\psi = i|u(t)|^2\psi + 2i \operatorname{Re}(\bar{u}(t)\psi)u(t), \quad \psi \in L^2(\mathbb{T}).$$

Thanks to step 3), the linear maps $P(t)$ are strongly continuously differentiable with respect to $t \in [0, a]$ as bounded operators on $L^2(\mathbb{T})$. Also, v_0 is contained in $H^2(\mathbb{T})$ and $\partial_t g$ in $C^1(\mathbb{R}_+; L^2(\mathbb{T}))$ by assumption (2). So (55) possesses a unique solution φ in $C^1([0, a]; L^2(\mathbb{T})) \cap C([0, a]; H^2(\mathbb{T}))$ by Theorem 6.3 in [Paz83]. The uniqueness of the corresponding integral equation (56) then shows that $\varphi = v = \partial_t u$, and so u belongs to $C^2([0, a]; L^2(\mathbb{T})) \cap C^1([0, a]; H^2(\mathbb{T}))$.

Step 5. We finally differentiate (1) twice with respect to x and obtain

$$\partial_t \partial_x^2 u = i\partial_x^4 u - \partial_x^2 u + P(t)\partial_x^2 u + 4i \operatorname{Re}(\bar{u}\partial_x u)\partial_x u + 2i|\partial_x u|^2 u + \partial_x^2 g$$

at first in $H^{-2}(\mathbb{T})$. This equation, assumption (2) and the previous steps imply that $(i\partial_x^2 - I)\partial_x^2 u$ is contained in $C([0, a]; L^2(\mathbb{T}))$; i.e., $u \in C([0, a]; H^4(\mathbb{T}))$.

Step 6. It remains to show global existence, where we employ the regularity of u established above. Using (1) and integrating by parts, we first compute

$$\begin{aligned}
\partial_t \|u(t)\|_{L^2}^2 &= 2 \operatorname{Re} \int_{\mathbb{T}} \bar{u}(t) \partial_t u(t) \, dx \\
&= 2 \operatorname{Re} \int_{\mathbb{T}} \bar{u}(t) \left(i\partial_x^2 u(t) - u(t) + i|u(t)|^2 u(t) + g(t) \right) \, dx \\
&= -2 \|u(t)\|_{L^2}^2 + 2 \operatorname{Re} \int_{\mathbb{T}} \bar{u}(t) g(t) \, dx \\
&\leq -2 \|u(t)\|_{L^2}^2 + 2 \|u(t)\|_{L^2} \|g(t)\|_{L^2} \\
&\leq -\|u(t)\|_{L^2}^2 + \|g(t)\|_{L^2}^2
\end{aligned} \tag{58}$$

for $t \in [0, a)$. Hence, $\partial_t (e^t \|u(t)\|_{L^2}^2) \leq e^t \|g(t)\|_{L^2}^2$ and integration yields

$$\begin{aligned}
\|u(t)\|_{L^2}^2 &\leq e^{-t} \|u_0\|_{L^2}^2 + \int_0^t e^{s-t} \|g(s)\|_{L^2}^2 \, ds \\
&\leq \|u_0\|_{L^2}^2 + \sup_{0 \leq s \leq b} \|g(s)\|_{L^2}^2 =: C_0(b)
\end{aligned} \tag{59}$$

for $0 \leq t \leq b < a$. We further need a modified energy of (1) given by

$$\mathcal{E}(t, v) = \frac{1}{2} \|\partial_x v\|_{L^2}^2 - \frac{1}{4} \|v\|_{L^4}^4 + \operatorname{Re} \int_{\mathbb{T}} i g(t) \bar{v} \, dx \tag{60}$$

for $v \in H^1(\mathbb{T})$ and $t \geq 0$. Proceeding as above, we obtain

$$\begin{aligned}
\partial_t \mathcal{E}(t, u(t)) &= \operatorname{Re} \int_{\mathbb{T}} [\partial_x u(t) \partial_{tx} \bar{u}(t) - (|u(t)|^2 u(t) - i g(t)) \partial_t \bar{u}(t) + i \bar{u}(t) \partial_t g(t)] \, dx \\
&= \operatorname{Re} \int_{\mathbb{T}} [(-\partial_x^2 u(t) - |u(t)|^2 u(t) + i g(t)) \partial_t \bar{u}(t) + i \bar{u}(t) \partial_t g(t)] \, dx \\
&= \operatorname{Re} \int_{\mathbb{T}} i [(\partial_t u(t) + u(t)) \partial_t \bar{u}(t) + \bar{u}(t) \partial_t g(t)] \, dx \\
&= \operatorname{Re} \int_{\mathbb{T}} i \left[u(t) (\bar{g}(t) - i \partial_x^2 \bar{u}(t) - \bar{u}(t) - i |u(t)|^2 \bar{u}(t)) + \bar{u}(t) \partial_t g(t) \right] \, dx \\
&= \operatorname{Re} \int_{\mathbb{T}} [i (u(t) \bar{g}(t) + \bar{u}(t) \partial_t g(t)) + |u(t)|^4 - |\partial_x u(t)|^2] \, dx
\end{aligned} \tag{61}$$

for $0 \leq t \leq b < a$. On the other hand, Sobolev's embedding theorem and complex interpolation (see Sect. 7.4.2 and 7.4.5 in [Tri92]) yield

$$\begin{aligned}
\|v\|_{L^4} &\leq C \|v\|_{H^{1/4}} \leq C \|v\|_{L^2}^{\frac{3}{4}} \|v\|_{H^1}^{\frac{1}{4}}, \\
\|v\|_{L^4}^4 &\leq \|\partial_x v\|_{L^2}^2 + \|v\|_{L^2}^2 + C \|v\|_{L^2}^6.
\end{aligned} \tag{62}$$

We thus deduce

$$\begin{aligned}
\partial_t \mathcal{E}(t, u(t)) &\leq \operatorname{Re} \int_{\mathbb{T}} i(u(t)\bar{g}(t) + \bar{u}(t)\partial_t g(t)) \, dx + \|u(t)\|_{L^2}^2 + C \|u(t)\|_{L^2}^6 \\
&\leq 2 \|u(t)\|_{L^2}^2 + \frac{1}{2} \|g(t)\|_{L^2}^2 + \frac{1}{2} \|\partial_t g(t)\|_{L^2}^2 + C \|u(t)\|_{L^2}^6 \\
&\leq 2C_0(b) + CC_0(b)^3 + C_1(b)
\end{aligned} \tag{63}$$

with $C_1(b) := \frac{1}{2} \sup_{0 \leq t \leq b} (\|g(t)\|_{L^2}^2 + \|\partial_t g(t)\|_{L^2}^2)$, where we also used (59). The estimate (62) further leads to the lower bound

$$\mathcal{E}(t, u(t)) \geq \frac{1}{4} \|\partial_x u(t)\|_{L^2}^2 - \frac{3}{4} C_0(b) - CC_0(b)^3 - \frac{1}{2} \|g(t)\|_{L^2}^2. \tag{64}$$

Combining (59), (63) and (64), we arrive at

$$\|u(t)\|_{H^1}^2 \leq 4\mathcal{E}(0, u_0) + C(1+b)(C_0(b) + C_0(b)^3 + C_1(b)) \tag{65}$$

for $0 \leq t \leq b < a$. If a was finite, we could take here $b = a$ and obtain a contradiction to the blow-up condition stated in Step 1). Hence, $a = \infty$.

Step 7. If g is bounded in $L^2(\mathbb{T})$, then (59) shows that $u(t)$ is bounded in $L^2(\mathbb{T})$ for $t \geq 0$. In particular, we can replace $C_0(b)$ by C in this case.

If also $\partial_t g$ is bounded in $L^2(\mathbb{T})$, then (65) implies that $u(t)$ grows at most as $\sqrt{1+t}$ in $H^1(\mathbb{T})$.

Next, if g is contained in $L^2(\mathbb{R}_+; L^2(\mathbb{T}))$, then we infer from the line before (59) and Young's convolution inequality that $u \in L^2(\mathbb{R}_+; L^2(\mathbb{T}))$.

If even $g \in H^1(\mathbb{R}_+; L^2(\mathbb{T}))$, then g is also bounded in $L^2(\mathbb{T})$. Thus u belongs $L^6(\mathbb{R}_+; L^2(\mathbb{T}))$. Integrating in t , we now deduce from the line before (63) that $\mathcal{E}(t, u(t))$ is uniformly bounded. Hence, the boundedness of $u(t)$ in $H^1(\mathbb{T})$ follows from (64). \square

Acknowledgements. The authors thank Sina Busch, Johannes Eilinghoff, Marlis Hochbruck, Christian Koos, Christian Lubich, Jörg Pfeifle, Wolfgang Reichel, and Katharina Schratz for helpful discussions on various topics related to this work.

References

- [BBD02] Christophe Besse, Brigitte Bidégaray, and Stéphane Descombes. Order estimates in time of splitting methods for the nonlinear Schrödinger equation. *SIAM J. Numer. Anal.*, 40(1):26–40, 2002.
- [Caz03] Thierry Cazenave. *Semilinear Schrödinger equations*, volume 10 of *Courant Lecture Notes in Mathematics*. New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2003.
- [CG12] David Cohen and Ludwig Gauckler. One-stage exponential integrators for nonlinear Schrödinger equations over long times. *BIT*, 52(4):877–903, 2012.

- [CKT07] M. Caliari, G. Kirchner, and M. Thalhammer. Convergence and energy conservation of the Strang time-splitting Hermite spectral method for nonlinear Schrödinger equations. Technical report, Universität Innsbruck, 2007.
- [CY10] Y. K. Chembo and Nan Yu. Modal expansion approach to optical-frequency-comb generation with monolithic whispering-gallery-mode resonators. *Physical Review A*, 82:033801, 2010.
- [DT13] Stéphane Descombes and Mechthild Thalhammer. The Lie-Trotter splitting for nonlinear evolutionary problems with critical parameters: a compact local error representation and application to nonlinear Schrödinger equations in the semiclassical regime. *IMA J. Numer. Anal.*, 33(2):722–745, 2013.
- [Fao12] Erwan Faou. *Geometric numerical integration and Schrödinger equations*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zurich, 2012.
- [FGL14] Erwan Faou, Ludwig Gauckler, and Christian Lubich. Plane wave stability of the split-step Fourier method for the nonlinear Schrödinger equation. *Forum Math. Sigma*, 2:e5, 45, 2014.
- [Gau11] Ludwig Gauckler. Convergence of a split-step Hermite method for the Gross-Pitaevskii equation. *IMA J. Numer. Anal.*, 31(2):396–415, 2011.
- [GL10] Ludwig Gauckler and Christian Lubich. Splitting integrators for nonlinear Schrödinger equations over long times. *Found. Comput. Math.*, 10(3):275–302, 2010.
- [HHR⁺12] T. Herr, K. Hartinger, J. Riemensberger, C.Y. Wang, E. Gavartin, R. Holzwarth, M.L. Gorodetsky, and T.J. Kippenberg. Universal formation dynamics and noise of Kerr-frequency combs in microresonators. *Nature Photonics*, 6:480–487, 2012.
- [HKT14] Harald Hofstätter, Othmar Koch, and Mechthild Thalhammer. Convergence analysis of high-order time-splitting pseudo-spectral methods for rotational Gross-Pitaevskii equations. *Numer. Math.*, 127(2):315–364, 2014.
- [HNW08] E. Hairer, S. P. Nørsett, and G. Wanner. *Solving ordinary differential equations I: Nonstiff Problems*, volume 8 of *Springer Series in Computational Mathematics*. Springer, Berlin, second edition, 2008.
- [HO10] Marlis Hochbruck and Alexander Ostermann. Exponential integrators. *Acta Numerica*, 19:209–286, 2010.
- [KI66] Herbert Bishop Keller and Eugene Isaacson. Analysis of numerical methods. *J. Wiley & Sons, New York*, 1966.

- [KNT13] Othmar Koch, Christof Neuhauser, and Mechthild Thalhammer. Error analysis of high-order splitting methods for nonlinear evolutionary Schrödinger equations and application to the MCTDHF equations in electron dynamics. *ESAIM, Math. Model. Numer. Anal.*, 47(5):1265–1286, 2013.
- [LL87] L. A. Lugiato and R. Lefever. Spatial dissipative structures in passive optical systems. *Phys. Rev. Lett.*, 58:2209–2211, 1987.
- [Lub08] Christian Lubich. On splitting methods for Schrödinger-Poisson and cubic nonlinear Schrödinger equations. *Math. Comp.*, 77(264):2141–2153, 2008.
- [ML15] Jeremy L. Marzuola and Jianfeng Lu. Strang splitting methods for a quasilinear Schrödinger equation: convergence, instability, and dynamics. *Commun. Math. Sci.*, 13(5):1051–1074, 2015.
- [Paz83] A Pazy. *Semigroups of linear operators and applications to partial differential equations*. Springer, New York, 1983.
- [SS99] Catherine Sulem and Pierre-Louis Sulem. *The nonlinear Schrödinger equation. Self-focusing and wave collapse*, volume 139 of *Applied Mathematical Sciences*. Springer, New York, 1999.
- [Tha12] Mechthild Thalhammer. Convergence analysis of high-order time-splitting pseudospectral methods for nonlinear Schrödinger equations. *SIAM J. Numer. Anal.*, 50(6):3231–3258, 2012.
- [Tri92] Hans Triebel. *Theory of function spaces II*, volume 84 of *Monographs in Mathematics*. Birkhäuser, Basel, 1992.