

Polynomial stability for a system of coupled strings

Łukasz Rzepnicki, Roland Schnaubelt

CRC Preprint 2017/35, December 2017

KARLSRUHE INSTITUTE OF TECHNOLOGY

CRC 1173



Participating universities



Universität Stuttgart

EBERHARD KARLS
UNIVERSITÄT
TÜBINGEN



Funded by

DFG

ISSN 2365-662X

POLYNOMIAL STABILITY FOR A SYSTEM OF COUPLED STRINGS

ŁUKASZ RZEPNICKI AND ROLAND SCHNAUBELT

ABSTRACT. We study the long-time behavior of two vibrating strings which are coupled at a common boundary point by a damper. We show that the classical solutions converge polynomially with a uniform rate, where the decay exponent depends on number theoretic properties of the quotient of the wave speeds of the two strings. The proof is based on a resolvent characterization of polynomial stability due to Borichev–Tomilov and Batty–Duyckaerts.

1. INTRODUCTION

In this paper we investigate the long-term behavior of two string equations which are coupled by a boundary damper, see (1.1) and (1.8). Their asymptotic properties heavily depend on the ratio $d = c_2/c_1$ of the wave speeds c_k of the two strings. The case of rational d was studied in [19] and [12]. In our main result Theorem 4.2 we show the polynomial convergence of all classical solutions to 0, to a constant or to an at most linearly growing solution, respectively, depending on the chosen boundary and interface conditions. The convergence rate is determined by a *irrationality measure* of d , which describes how fast d can be approximated by (appropriate) rational numbers, cf. Section 3. We also see that the obtained decay rate is almost sharp. Such diophantine properties of a parameter have been used before in the stability theory of evolution equations in e.g. [1], [14], [15] or [26], where other problems were treated by different methods. However, our system seems to be the first one exhibiting an explicit dependence of the (polynomial) decay on such number theoretic quantities.

In recent years there has been an important progress in the understanding of stability properties of linear evolution equations beyond the well studied case of exponential decay. In a Hilbert space, Gearhart’s theorem characterizes the exponential convergence to 0 of all (mild) solutions by the boundedness of the resolvent $R(\lambda, A)$ for $\operatorname{Re} \lambda \geq 0$, where A is the generator governing the problem. But damping mechanism in wave type equations often lead to weaker decay properties, see e.g. the discussion in [2].

1991 *Mathematics Subject Classification*. Primary: 35B40. Secondary: 35L53, 47D06.

Key words and phrases. Polynomial decay, boundary damping, resolvent bound, irrationality measure.

We thank Przemysław Berk, Yann Bugeaud, Stefan Kühnlein and Andrzej Schinzel for their help concerning questions about number theory. R.S. gratefully acknowledges financial support by the Deutsche Forschungsgemeinschaft (DFG) through CRC 1173. Ł. Rz. was partially supported by the National Science Centre (Poland) grant DEC-2014/13/B/ST1/03153.

Logarithmic decay of all classical solutions u (i.e., $u(0) \in D(A)$) was characterized in terms of resolvent estimates in the pioneering work [18]. For a bounded semigroup $T(\cdot)$ with generator A on a Hilbert space X , one has the polynomial decay $\|T(t)(I - A)^{-1}\| \leq ct^{-1/\alpha}$ for $t \geq 1$ and some $\alpha > 0$ if and only if the spectrum of A belongs to the open left half plane and $\|(is - A)^{-1}\| \leq c|s|^\alpha$ for real s with $|s| \geq 1$. The sufficiency of the conditions on A was shown in [8], whereas their necessity is true even in a Banach space X by [6]. (See [3] and [20] for earlier contributions.) The sufficiency fails in L^1 -spaces in general, see [8]. Still in Banach spaces the resolvent condition implies the corresponding decay up to a logarithmic correction, as shown in [6] also for more general rates. In the recent papers [4] and [5] these results have been refined in various directions. In these papers one finds plenty of references concerning applications to PDEs, see also [2], [3], [6], [8], [20].

In this work we establish the polynomial decay of the system of two strings

$$\begin{aligned} \partial_{tt}u(t, x) &= c_1^2 \partial_{xx}u(t, x), & x \in (-1, 0), \quad t \geq 0, \\ \partial_{tt}v(t, x) &= c_2^2 \partial_{xx}v(t, x), & x \in (0, 1), \quad t \geq 0, \\ u(0, x) &= u_0(x), \quad \partial_t u(0, x) = u_1(x), & x \in (-1, 0), \\ v(0, x) &= v_0(x), \quad \partial_t v(0, x) = v_1(x), & x \in (0, 1), \end{aligned} \quad (1.1)$$

with the wave speeds $c_k = \sqrt{T_k/m_k}$ and initial functions u_0, u_1, v_0, v_1 . The mass and tension densities $m_1, m_2 > 0$ and $T_1, T_2 > 0$ are given constants. In (1.8) we add to (1.1) some of the boundary and interface conditions

$$u(t, -1) = 0, \quad (1.2)$$

$$\partial_x u(t, -1) = 0, \quad (1.3)$$

$$v(t, 1) = 0, \quad (1.4)$$

$$\partial_x v(t, 1) = 0, \quad (1.5)$$

$$T_1 \partial_x u(t, 0) = T_2 \partial_x v(t, 0), \quad \partial_t u(t, 0) - \partial_t v(t, 0) = -kT_1 \partial_x u(t, 0), \quad (1.6)$$

$$u(t, 0) = v(t, 0), \quad T_1 \partial_x u(t, 0) - T_2 \partial_x v(t, 0) = -k \partial_t u(t, 0), \quad (1.7)$$

for $t \geq 0$ and a damping constant $k > 0$.

This system was introduced in [11], where a mechanical interpretation of the interface damping was given and in particular the case $c_1 = c_2$ was studied. The long-time behavior of this system was further investigated in [19]. There it was seen that the quotient $d = c_2/c_1$ plays a crucial role in the analysis of (1.1). If it is a rational number of the right type, see (2.19), then the energy of all solutions decays exponentially. For other rational numbers there are time periodic solutions and thus no decay. For irrational d , convergence of the solutions was shown without providing rates. Actually, the paper [19] contains some errors in details, partly fixed in [12], see Section 2.

Let d be irrational. As in [19], we complement (1.1) by the combinations

$$\begin{aligned} \text{Case I:} & \quad (1.2) \quad \text{and} \quad (1.4) \quad \text{and} \quad (1.6), \\ \text{Case II:} & \quad (1.2) \quad \text{and} \quad (1.5) \quad \text{and} \quad (1.6), \\ \text{Case III:} & \quad (1.3) \quad \text{and} \quad (1.5) \quad \text{and} \quad (1.6), \\ \text{Case IV:} & \quad (1.2) \quad \text{and} \quad (1.4) \quad \text{and} \quad (1.7), \\ \text{Case V:} & \quad (1.2) \quad \text{and} \quad (1.5) \quad \text{and} \quad (1.7), \\ \text{Case VI:} & \quad (1.3) \quad \text{and} \quad (1.5) \quad \text{and} \quad (1.7). \end{aligned} \quad (1.8)$$

We note that the long-time behavior in case III was not studied in [19] and [12]. In the next section we see that in each case the system is governed by a generator A_j on a suitable Hilbert space \mathcal{H}_j . In the cases I, IV and V the spectrum of A_j belongs to the open left half plane, whereas in the other cases 0 is an isolated eigenvalue of A_j . More precisely, in II and VI we only have a nontrivial kernel of A_j , but A_{III} also has a proper generalized eigenvector for $\lambda = 0$. Accordingly, we show polynomial decay to 0, respectively to a constant, for $j \neq III$, and to an at most linearly growing solution if $j = III$, see Theorems 4.2 and 4.3. It is crucial for our approach to obtain the boundedness of the solution semigroup, respectively of its restriction to the kernel of the spectral projection for the spectral set $\{0\}$. This fact is fairly standard for $j \in \{I, IV, V\}$. For $j \in \{II, VI\}$ one can use ideas from the recent paper [7] about a certain parabolic-hyperbolic system in one space dimension. The case III, however, requires a detailed analysis of the spectral projection, cf. Proposition 2.2.

The main results heavily depend on the decay behavior of a complex function $\Delta_j(is)$ as $s \rightarrow \pm\infty$, see (2.12), since the resolvent $(is - A_j)^{-1}$ is bounded by $c|\Delta_j(is)|^{-1}$ thanks to Proposition 2.2. The map $\Delta_j(is)$ is a linear combination of products of $\sin(c_k s)$ and $\cos(c_k s)$ for $k \in \{1, 2\}$. Therefore its decay is determined by the rate with which one can approximate d by fractions p/q of the type odd/odd, even/odd or odd/even, depending of the case j , see (3.2). We call the optimal rate $\mu_j(d)$, and it is closely related to the irrationality measure on d , which has thoroughly been studied in number theory as discussed in Section 3. One has $\mu_j(d) \geq 2$ for every irrational d and equality holds if d algebraic by Roth's theorem [22].

Depending on $\mu_j(d)$, we can then compute the decay rate of $\Delta_j(is)$ to 0 in Lemma 3.1 in a rather delicate way where we crucially use the structure of $\Delta_j(is)$. It is further shown that the obtained rate is almost optimal. Astonishingly one can discuss the six cases more or less in parallel. In the last section we then prove our main results, also using the results of [6] and [8]. For classical solutions, Theorem 4.2 gives almost optimal polynomial decay rates in terms of the irrationality measure of d . For instance, if $\mu_j(d) = 2$, we have the decay $c_\epsilon t^{-\frac{1}{2}-\epsilon}$ for all $\epsilon > 0$ and solutions starting in the unit ball of $D(A_j)$, and we know that this exponent is almost sharp. Based on somewhat different number theoretic results, in Theorem 4.3 we finally show decay by $t^{-1/2}$ times a logarithmic correction, for a.e. d .

2. BASIC PROPERTIES OF THE SOLUTION SEMIGROUP

We rewrite the system (1.1) with the boundary conditions in (1.8) as an evolution equation of first order in time for the states $(u, \partial_t u, v, \partial_t v) = w = (w_1, w_2, w_3, w_4)$ in the product space

$$\mathcal{H} = H^1(-1, 0) \times L_2(-1, 0) \times H^1(0, 1) \times L_2(0, 1)$$

which is equipped with its canonical norm $\|\cdot\|_{\mathcal{H}}$. We mostly work in smaller state spaces which include the Dirichlet conditions from (1.8); namely in

$$\begin{aligned} \mathcal{H}_I &= \{w \in \mathcal{H} \mid w_1(-1) = 0 = w_3(1)\}, \\ \mathcal{H}_{II} &= \{w \in \mathcal{H} \mid w_1(-1) = 0\}, \end{aligned}$$

$$\begin{aligned}
\mathcal{H}_{III} &= \mathcal{H}, \\
\mathcal{H}_{IV} &= \{w \in \mathcal{H} \mid w_1(-1) = 0 = w_3(1), w_1(0) = w_3(0)\}, \\
\mathcal{H}_V &= \{w \in \mathcal{H} \mid w_1(-1) = 0, w_1(0) = w_3(0)\}, \\
\mathcal{H}_{VI} &= \{w \in \mathcal{H} \mid w_1(0) = w_3(0)\}
\end{aligned} \tag{2.1}$$

for the respective cases $j \in J := \{I, II, III, IV, V, VI\}$, noting that $H^1(a, b)$ embeds into $C([a, b])$. We endow \mathcal{H}_I , \mathcal{H}_{IV} , and \mathcal{H}_V with the scalar product

$$\langle w, z \rangle_E = \frac{1}{2} \int_{-1}^0 (T_1 w_1' \bar{z}_1' + m_1 w_2 \bar{z}_2) dx + \frac{1}{2} \int_0^1 (T_2 w_3' \bar{z}_3' + m_2 w_4 \bar{z}_4) dx.$$

They are Hilbert spaces because of the Dirichlet conditions imposed in (2.1). Note that the square of the induced norm

$$\|w\|_E^2 = \langle w, w \rangle_E = \frac{T_1}{2} \|w_1'\|_2^2 + \frac{m_1}{2} \|w_2\|_2^2 + \frac{T_2}{2} \|w_3'\|_2^2 + \frac{m_2}{2} \|w_4\|_2^2$$

is equal to the system's energy given by

$$\mathcal{E}(t) = \frac{1}{2} \int_{-1}^0 (m_1 (\partial_t u)^2 + T_1 (\partial_x u)^2) dx + \frac{1}{2} \int_0^1 (m_2 (\partial_t v)^2 + T_2 (\partial_x v)^2) dx.$$

For $\|\cdot\|_E$ we can check dissipativity of the generators below. The spaces \mathcal{H}_{II} , \mathcal{H}_{III} , and \mathcal{H}_{VI} are equipped with the modified scalar product

$$\begin{aligned}
\langle\langle w, z \rangle\rangle_E &= \frac{1}{2} \int_{-1}^0 (T_1 w_1' \bar{z}_1' + m_1 w_1 \bar{z}_1 + m_1 w_2 \bar{z}_2) dx \\
&\quad + \frac{1}{2} \int_0^1 (T_2 w_3' \bar{z}_3' + m_2 w_3 \bar{z}_3 + m_2 w_4 \bar{z}_4) dx,
\end{aligned}$$

whose induced norm is denoted by $\|\|\cdot\|\|_E$. Also these spaces are Hilbertian. For each $j \in J$, on \mathcal{H}_j the norms $\|\cdot\|_E$ resp. $\|\|\cdot\|\|_E$, are equivalent to $\|\cdot\|_{\mathcal{H}}$.

In every case the generator corresponding to (1.1) is a restriction of the operator matrix

$$\mathcal{A} = \begin{pmatrix} 0 & I & 0 & 0 \\ c_1^2 D_{xx} & 0 & 0 & 0 \\ 0 & 0 & 0 & I \\ 0 & 0 & c_2^2 D_{xx} & 0 \end{pmatrix} \quad \text{with domain}$$

$$\mathcal{D} = H^2(-1, 0) \times H^1(-1, 0) \times H^2(0, 1) \times H^1(0, 1)$$

in \mathcal{H} . In view of the boundary and interface conditions in (1.8), we define

$$\begin{aligned}
D(A_I) &= \{w \in \mathcal{D} \cap \mathcal{H}_I \mid w_2(-1) = 0 = w_4(-1), T_1 w_1'(0) = T_2 w_3'(0), \\
&\quad w_2(0) - w_4(0) = -k T_1 w_1'(0)\},
\end{aligned}$$

$$\begin{aligned}
D(A_{II}) &= \{w \in \mathcal{D} \cap \mathcal{H}_{II} \mid w_2(-1) = 0, w_3'(1) = 0, T_1 w_1'(0) = T_2 w_3'(0), \\
&\quad w_2(0) - w_4(0) = -k T_1 w_1'(0)\},
\end{aligned}$$

$$\begin{aligned}
D(A_{III}) &= \{w \in \mathcal{D} \mid w_1'(-1) = 0, w_3'(1) = 0, T_1 w_1'(0) = T_2 w_3'(0), \\
&\quad w_2(0) - w_4(0) = -k T_1 w_1'(0)\},
\end{aligned}$$

$$\begin{aligned}
D(A_{IV}) &= \{w \in \mathcal{D} \cap \mathcal{H}_{IV} \mid w_2(-1) = 0 = w_4(1), \\
&\quad T_1 w_1'(0) - T_2 w_3'(0) = -k w_2(0) = -k w_4(0)\},
\end{aligned}$$

$$\begin{aligned}
D(A_V) &= \{w \in \mathcal{D} \cap \mathcal{H}_V \mid w_2(-1) = 0, w_3'(1) = 0, \\
&\quad T_1 w_1'(0) - T_2 w_3'(0) = -k w_2(0) = -k w_4(0)\},
\end{aligned}$$

$$D(A_{VI}) = \{w \in \mathcal{D} \cap \mathcal{H}_{VI} \mid w_1'(-1) = 0 = w_3'(1), \\ T_1 w_1'(0) - T_2 w_3'(0) = -k w_2(0) = -k w_4(0)\},$$

and set $A_j w = \mathcal{A}w$ for $w \in D(A_j)$. Some of the above equations for w_2 and w_4 are not explicitly stated in (1.8). They arise from differentiating the Dirichlet conditions on w_1 and w_3 in (1.8) with respect to time t , and are thus implicitly contained in the system. Observe that the restriction $A_j : D(A_j) \rightarrow \mathcal{H}_j$ actually maps into \mathcal{H}_j for each $j \in J$.

We look for solutions $(u, v) \in C^2(\mathbb{R}_+, L^2(0, 1))^2 \cap C(\mathbb{R}_+, H^2(0, 1))^2$ of the problem (1.1) with the boundary conditions given by (1.8). Existence and uniqueness of such solutions are established in Proposition 2.2 below.

This and most of the other results in this section are contained in [19] and [12] in a somewhat different form since there the wave equations (1.1) were rewritten as a first order system. However, the precise expression for the resolvent of A_j is needed later on. Moreover, the transformation to the first order system has a one- or two-dimensional kernel in Cases II, III, and VI which was overlooked in [19] and [12]. In addition Case III was treated only partly there. We thus give most of the proofs within our framework.

We first show dissipativity and deal with the kernels of A_{II} , A_{III} and A_{VI} . For our main results we have to describe the spectral projection for the set $\{0\}$ of these operators, which will be achieved for $j = III$ only in Proposition 2.2. In the cases II and VI we argue as in [7].

Lemma 2.1. *a) The operators A_j are injective and dissipative on \mathcal{H}_j for $j \in \{I, IV, V\}$.*

b) Let $j \in \{II, VI\}$. Then the operator $A_j - I$ is dissipative on \mathcal{H}_j . Moreover, A_j has a one-dimensional kernel and a closed range with $\mathcal{H}_j = \text{ran } A_j \oplus \ker A_j$. The projections onto $\ker A_j$ along $\text{ran } A_j$ are given by $P_{II}g = (0, 0, \phi_{II}(g)\mathbb{1}, 0)$ and $P_{VI}g = (\phi_{VI}(g)\mathbb{1}, 0, \phi_{VI}(g)\mathbb{1}, 0)$ for the functionals ϕ_j defined in (2.5) respectively (2.7). On $\text{ran } A_j$ the norm of \mathcal{H}_j is equivalent to $\|\cdot\|_E$, and the part of A_j in $\text{ran } A_j$ is dissipative for $\langle \cdot, \cdot \rangle_E$.

c) The operator $A_{III} - I$ is dissipative on \mathcal{H}_{III} . The space of generalized eigenfunctions of A_{III} for $\lambda = 0$ is spanned by $(\mathbb{1}, 0, 0, 0)$, $(0, 0, \mathbb{1}, 0)$ and $(0, \mathbb{1}, 0, \mathbb{1})$, where the first two functions form a basis for $\ker A_{III}$.

Proof. 1) We first treat the cases $j \in \{I, IV, V\}$. Let $w \in D(A_j)$. Integrating by parts and using the boundary conditions at ± 1 we compute

$$\begin{aligned} 2 \langle A_j w, w \rangle_E &= T_1 \int_{-1}^0 (w_2' \bar{w}_1' + w_1'' \bar{w}_2) dx + T_2 \int_0^1 (w_4' \bar{w}_3' + w_3'' \bar{w}_4) dx \\ &= T_1 \int_{-1}^0 (w_2' \bar{w}_1' - w_1' \bar{w}_2') dx + T_2 \int_0^1 (w_4' \bar{w}_3' - w_3' \bar{w}_4') dx \\ &\quad + T_1 w_1' \bar{w}_2 \Big|_{-1}^0 + T_2 w_3' \bar{w}_4 \Big|_0^1, \\ 2 \text{Re} \langle A_j w, w \rangle_E &= T_1 w_1'(0) \bar{w}_2(0) - T_2 w_3'(0) \bar{w}_4(0). \end{aligned} \quad (2.2)$$

In case I, the interface conditions in $D(A_I)$ then yield

$$2 \text{Re} \langle A_I w, w \rangle_E = -T_1^2 k |w_1'(0)|^2. \quad (2.3)$$

Similarly, the conditions corresponding to (1.7) imply

$$2 \text{Re} \langle A_j w, w \rangle_E = -k |w_2(0)|^2 \quad (2.4)$$

for $j \in \{IV, V\}$. So these operators are dissipative. If $A_j w = 0$, then the functions w_1 and w_3 are affine, whereas w_2 and w_4 vanish. The interface equations further yield $T_1 w_1'(0) = T_2 w_3'(0)$, and this number equals 0 if $j \in \{I, V\}$. Finally, the relations in \mathcal{H}_j show that also $w_1 = w_3 = 0$, and hence A_j is injective for $j \in \{I, IV, V\}$.

2) We next look at the cases $j \in \{II, III, VI\}$. Let $w \in D(A_j)$. As in (2.2)–(2.4) we compute

$$\begin{aligned} & 2 \operatorname{Re} \langle A_j w, w \rangle_E \\ &= T_1 w_1'(0) \bar{w}_2(0) - T_2 w_3'(0) \bar{w}_4(0) + m_1 \operatorname{Re} \int_{-1}^0 w_2 \bar{w}_1 \, dx + m_2 \operatorname{Re} \int_0^1 w_4 \bar{w}_3 \, dx \\ &\leq T_1 w_1'(0) \bar{w}_2(0) - T_2 w_3'(0) \bar{w}_4(0) + \frac{m_1}{2} (\|w_1\|_2^2 + \|w_2\|_2^2) + \frac{m_2}{2} (\|w_3\|_2^2 + \|w_4\|_2^2) \\ &\leq \frac{m_1}{2} (\|w_1\|_2^2 + \|w_2\|_2^2) + \frac{m_2}{2} (\|w_3\|_2^2 + \|w_4\|_2^2). \end{aligned}$$

Consequently, $A_j - I$ is dissipative.

Let $A_j w = 0$. As above, then w_1 and w_3 are affine, whereas w_2 and w_4 are zero. The Neumann conditions in the domains imply that w_1 and w_3 are constant. The Dirichlet conditions in \mathcal{H}_j now yield that $\ker A_{II}$ is spanned by $(0, 0, \mathbb{1}, 0)$ and $\ker A_{VI}$ by $(\mathbb{1}, 0, \mathbb{1}, 0)$, whereas $\ker A_{III}$ has the basis $\{(\mathbb{1}, 0, 0, 0), (0, 0, \mathbb{1}, 0)\}$.

Moreover, a function $w \in D(A_{III})$ satisfies $A_{III} w = (a\mathbb{1}, 0, b\mathbb{1}, 0)$ for some numbers $a, b \in \mathbb{C}$ if and only if all components w_j are constant and $w_2 = a\mathbb{1}$ is equal to $w_4 = b\mathbb{1}$.

3) To establish the asserted decompositions in b) for $j \in \{II, VI\}$, we first introduce the bounded linear map

$$\phi_{II} : \mathcal{H}_{II} \rightarrow \mathbb{C}; \quad \phi_{II}(g_1, g_2, g_3, g_4) = g_3(0) - g_1(0) + \frac{kT_2}{c_2^2} \int_0^1 g_4(x) \, dx. \quad (2.5)$$

First, let $f = A_{II} w$ belong to the range of A_{II} . We then deduce the equations $w_2 = f_1$, $w_4 = f_3$,

$$\begin{aligned} w_1(x) &= a(1+x) + \frac{1}{c_1^2} \int_{-1}^x (x-s) f_2(s) \, ds, & x \in [-1, 0], \\ w_3(x) &= b + \frac{1}{c_2^2} \int_x^1 (s-x) f_4(s) \, ds, & x \in [0, 1], \end{aligned} \quad (2.6)$$

for complex numbers a and b , taking into account the boundary conditions $w_1(-1) = 0$ and $w_3'(1) = 0$. Note that

$$w_1'(0) = a + \frac{1}{c_1^2} \int_{-1}^0 f_2(s) \, ds \quad \text{and} \quad w_3'(0) = -\frac{1}{c_2^2} \int_0^1 f_4(s) \, ds.$$

The interface conditions in $D(A_{II})$ then determine the parameter a and imply that the vector f belongs to $\ker \phi_{II}$.

Conversely, let $f \in \mathcal{H}_{II}$ be contained in the kernel of ϕ_{II} . We then define w by $w_2 = f_1$, $w_4 = f_3$, and (2.6) with $b = 0$ and

$$a = \frac{f_3(0) - f_1(0)}{kT_1} - \frac{1}{c_1^2} \int_{-1}^0 f_2(x) \, dx = -\frac{T_2}{T_1 c_2^2} \int_0^1 f_4(x) \, dx - \frac{1}{c_1^2} \int_{-1}^0 f_2(x) \, dx,$$

where we have used $\phi_{II}(f) = 0$ in the second equation. One can now check that w belongs to $D(A_{II})$, and we have $A_{II}w = f$ by construction. Therefore, $\text{ran } A_{II} = \ker \phi_{II}$ is closed and the map $P_{II}g = (0, 0, \phi_{II}(g)\mathbb{1}, 0)$ is the bounded projection onto $\ker A_{II}$ along $\text{ran } A_{II}$.

Let $g \in \text{ran } A_{II}$. We clearly have $\|g\|_E^2 \leq \|g\|$ and also $\|g_1\|_{H^1} \leq c\|g'_1\|_{L^2}$ since $g_1(-1) = 0$, for some constants $c > 0$. Due to $\phi_{II}(g) = 0$, we can then bound $|g_3(0)|$ by $c(\|g'_1\|_{L^2} + \|g_4\|_{L^2})$. Hence, the norm $\|\cdot\|$ is equivalent to $\|\cdot\|_E$ on $\text{ran } A_{II}$. The dissipativity of the part of A_{II} in $\text{ran } A_{II}$ is proved as in (2.3).

The remaining assertions for A_{VI} are similarly shown using the functional

$$\phi_{VI} : \mathcal{H}_{VI} \rightarrow \mathbb{C}; \quad \phi_{VI}(f) = \frac{T_1}{kc_1^2} \int_{-1}^0 f_2(s) ds + \frac{T_2}{kc_2^2} \int_0^1 f_4(s) ds + f_1(0) \quad (2.7)$$

and the projection $P_{VI}g = (\phi_{VI}(g)\mathbb{1}, 0, \phi_{VI}(g)\mathbb{1}, 0)$. \square

We next describe the spectrum of A_j and compute its resolvent. Take $j \in J$, $f \in \mathcal{H}_j$, and $\lambda \in \mathbb{C} \setminus \{0\}$. We introduce the numbers

$$a_l = \frac{T_l}{c_l}, \quad \omega_l = \frac{\lambda}{c_l}, \quad c_l = \frac{\sqrt{T_l}}{\sqrt{m_l}}, \quad d = \frac{c_2}{c_1}$$

for $\lambda \in \mathbb{C}$ and $l \in \{1, 2\}$, and the functions

$$g_j : \mathbb{C} \rightarrow \mathbb{C}; \quad g_j(z) = \begin{cases} \sinh(z + \omega_1), & j \in \{I, II, IV, V\}, \\ \cosh(z + \omega_1), & j \in \{III, VI\}, \end{cases}$$

$$h_j : \mathbb{C} \rightarrow \mathbb{C}; \quad h_j(z) = \begin{cases} \sinh(\omega_2 - z), & j \in \{I, IV\}, \\ \cosh(\omega_2 - z), & j \in \{II, III, V, VI\}. \end{cases}$$

The function \sinh corresponds to Dirichlet conditions at $x = -1$ for g_j and at $x = 1$ for h_j , whereas \cosh is used in the Neumann cases. We further set

$$U_\lambda(x) = \frac{-1}{c_1\lambda} \int_{-1}^x \sinh(\omega_1(x-r))(\lambda f_1(r) + f_2(r)) dr, \quad x \in [-1, 0],$$

$$W_\lambda(x) = \frac{-1}{c_2\lambda} \int_x^1 \sinh(\omega_2(r-x))(\lambda f_3(r) + f_4(r)) dr, \quad x \in [0, 1]. \quad (2.8)$$

All solutions $w \in \mathcal{D}$ to $\lambda w - \mathcal{A}w = f$ satisfying the boundary conditions at $x = \pm 1$ for Case j in (1.8) are given by

$$\begin{aligned} w_1 &= \alpha(\lambda)g_j(\omega_1 \cdot) + U_\lambda, & w_2 &= \lambda w_1 - f_1 & \text{on } [-1, 0], \\ w_3 &= \beta(\lambda)h_j(\omega_2 \cdot) + W_\lambda, & w_4 &= \lambda w_3 - f_3 & \text{on } [0, 1], \end{aligned} \quad (2.9)$$

for some numbers $a(\lambda), b(\lambda) \in \mathbb{C}$. Since f belongs to \mathcal{H}_j , these functions w_2 and w_4 satisfy the Dirichlet conditions at $x = \pm 1$ in $D(A_j)$.

We next impose the interface conditions induced by (1.6) for $j \in \{I, II, III\}$ and by (1.7) for $j \in \{IV, V, VI\}$. They are satisfied if and only if the solution w fulfill the equations

$$\begin{aligned} \lambda a_1 \alpha(\lambda) g'_j(0) + T_1 U'_\lambda(0) &= \lambda a_2 \beta(\lambda) h'_j(0) + T_2 W'_\lambda(0), \\ \lambda \alpha(\lambda) g_j(0) + \lambda U_\lambda(0) - f_1(0) - \lambda \beta(\lambda) h_j(0) - \lambda W_\lambda(0) + f_3(0) \\ &= -k(\lambda a_1 \alpha(\lambda) g'_j(0) + T_1 U'_\lambda(0)). \end{aligned}$$

if $j \in \{I, II, III\}$, as well as

$$\begin{aligned} \lambda\alpha(\lambda)g_j(0) + \lambda U_\lambda(0) &= \lambda\beta(\lambda)h_j(0) + \lambda W_\lambda(0), \\ \lambda a_1\alpha(\lambda)g_j'(0) + T_1U_\lambda'(0) - \lambda a_2\beta(\lambda)h_j'(0) - T_2W_\lambda'(0) \\ &= -k(\lambda\alpha(\lambda)g_j(0) + \lambda U_\lambda(0) - f_1(0)). \end{aligned}$$

if $j \in \{IV, V, VI\}$. (For w_2 and w_4 we employ (2.9) and $f \in \mathcal{H}_j$.)

We thus obtain a solution $w \in D(A_j)$ to $\lambda w - \mathcal{A}w = f$ if and only if the coefficients $(a(\lambda), b(\lambda))$ solve the system

$$\begin{aligned} M_j(\lambda) \begin{pmatrix} \alpha(\lambda) \\ \beta(\lambda) \end{pmatrix} &= \begin{pmatrix} -T_1U_\lambda'(0) + T_2W_\lambda'(0) \\ -\lambda U_\lambda(0) + \lambda W_\lambda(0) - kT_1U_\lambda'(0) + f_1(0) - f_3(0) \end{pmatrix}, \\ M_j(\lambda) &= \lambda \begin{pmatrix} a_1g_j'(0) & -a_2h_j'(0) \\ g_j(0) + a_1kg_j'(0) & -h_j(0) \end{pmatrix}, \end{aligned} \quad (2.10)$$

in the cases $j \in \{I, II, III\}$, respectively

$$\begin{aligned} M_j(\lambda) \begin{pmatrix} \alpha(\lambda) \\ \beta(\lambda) \end{pmatrix} &= \begin{pmatrix} -\lambda U_\lambda(0) + \lambda W_\lambda(0) \\ -T_1U_\lambda'(0) + T_2W_\lambda'(0) - k\lambda U_\lambda(0) + kf_1(0) \end{pmatrix}, \\ M_j(\lambda) &= \lambda \begin{pmatrix} g_j(0) & -h_j(0) \\ a_1g_j'(0) + kg_j(0) & -a_2h_j'(0) \end{pmatrix} \end{aligned} \quad (2.11)$$

for $j \in \{IV, V, VI\}$. The determinant of $M_j(\lambda)$ is given by

$$\begin{aligned} \det M_j(\lambda) &= -\lambda^2 \Delta_j(\lambda), \quad j \in \{I, II, III\}, \\ \det M_j(\lambda) &= \lambda^2 \Delta_j(\lambda), \quad j \in \{IV, V, VI\}, \end{aligned} \quad (2.12)$$

$$\begin{aligned} \Delta_I(\lambda) &= [\sinh(\omega_1) + a_1k \cosh(\omega_1)]a_2 \cosh(\omega_2) + a_1 \cosh(\omega_1) \sinh(\omega_2), \\ \Delta_{II}(\lambda) &= [\sinh(\omega_1) + a_1k \cosh(\omega_1)]a_2 \sinh(\omega_2) + a_1 \cosh(\omega_1) \cosh(\omega_2), \\ \Delta_{III}(\lambda) &= [\cosh(\omega_1) + a_1k \sinh(\omega_1)]a_2 \sinh(\omega_2) + a_1 \sinh(\omega_1) \cosh(\omega_2), \\ \Delta_{IV}(\lambda) &= [a_1 \cosh(\omega_1) + k \sinh(\omega_1)] \sinh(\omega_2) + a_2 \sinh(\omega_1) \cosh(\omega_2), \\ \Delta_V(\lambda) &= [a_1 \cosh(\omega_1) + k \sinh(\omega_1)] \cosh(\omega_2) + a_2 \sinh(\omega_1) \sinh(\omega_2), \\ \Delta_{VI}(\lambda) &= [a_1 \sinh(\omega_1) + k \cosh(\omega_1)] \cosh(\omega_2) + a_2 \cosh(\omega_1) \sinh(\omega_2). \end{aligned}$$

If $\Delta_j(\lambda) = 0$ for some $\lambda \neq 0$, the above observations yield a non-zero solution $w \in D(A_j)$ of $\lambda w - A_j w = 0$, and thus λ is an eigenvalue of A_j . Otherwise we have constructed a unique solution $w \in D(A_j)$ of the equation $\lambda w - A_j w = f$ for the given function $f \in \mathcal{H}_j$, where the map $f \mapsto w$ is continuous in \mathcal{H}_j . Hence, $\lambda \neq 0$ belongs to the resolvent set if $\Delta_j(\lambda) \neq 0$.

Observe that the functions Δ_j are positive on $(0, \infty)$, so that the resolvent of A_j exists for $\lambda > 0$, and that it is compact. Because of these properties and Lemma 2.1, A_j is invertible for $j \in \{I, IV, V\}$ and 0 is an isolated eigenvalue of A_j for $j \in \{II, III, VI\}$.

We state these facts and several consequences in the next basic proposition, using the following notation. The spectrum of a closed operator B is denoted by $\sigma(A)$ and its point spectrum by $\sigma_p(B)$. We write $\mathbb{C}_- = \{\lambda \in \mathbb{C} \mid \operatorname{Re} \lambda < 0\}$.

Proposition 2.2. *a) The resolvent of A_j is compact, and we have*

$$\sigma(A_j) = \sigma_p(A_j) = \{\lambda \in \mathbb{C} \setminus \{0\} \mid \Delta_j(\lambda) = 0\}, \quad j \in \{I, IV, V\},$$

$$\sigma(A_j) = \sigma_p(A_j) = \{0\} \cup \{\lambda \in \mathbb{C} \setminus \{0\} \mid \Delta_j(\lambda) = 0\}, \quad j \in \{II, III, VI\}.$$

b) For $j \in \{I, IV, V\}$ the operator A_j generates a contractive C_0 -semigroup $T_j(\cdot)$ on \mathcal{H}_j .

c) Let $j \in \{II, III, VI\}$. Then 0 is an isolated eigenvalue of A_j . Let P_j be the corresponding spectral projection and A_j^1 be the restriction of A_j to $D(A_j) \cap \ker P_j$. Then P_{II} and P_{VI} are given as in Lemma 2.1 and P_{III} by (2.14). We have $\sigma(A_j^1) = \sigma(A_j) \setminus \{0\}$.

For $j \in \{II, IV\}$ the operator A_j generates a bounded C_0 -semigroup $T_j(\cdot)$ on \mathcal{H}_j . The restriction of $T_j(\cdot)$ to $\text{ran } P_j = \ker A_j$ is constant and that to $\ker P_j = \text{ran } A_j$ is contractive for $\|\cdot\|_E$ and has the generator A_j^1 .

On $\ker P_{III}$ the norm of \mathcal{H}_{III} is equivalent to $\|\cdot\|_E$. The operator A_{III} generates a C_0 -semigroup $T_{III}(\cdot)$ on \mathcal{H}_{III} with linear growth. The restriction of $T_{III}(\cdot)$ to $\text{ran } P_j$ is affine in t and that to $\ker P_j$ is contractive for $\|\cdot\|_E$ and has the generator A_{III}^1 .

c) In each case, for every $w_0 = (u_0, u_1, v_0, v_1) \in D(A_j)$ the unique solution (u, v) of (1.1) with the boundary conditions of (1.8) is given by $w(t) = (u(t), \partial_t u(t), v(t), \partial_t v(t)) = T_j(t)w_0$ for $t \geq 0$.

d) Let $d = c_2/c_1 \notin \mathbb{Q}$. Then $\sigma(A_j) \subseteq \mathbb{C}_-$ for $j \in \{I, IV, V\}$ and $\sigma(A_j^1) \subseteq \mathbb{C}_-$ for $j \in \{II, III, VI\}$. The resolvent is bounded by

$$\|R(is, A_j)\| \leq \frac{c}{|\Delta_j(is)|} \quad (2.13)$$

for a constant $c > 0$, all $s \in \mathbb{R}$ with $|s| \geq 1$, and $j \in J$.

Proof. 1) Assertion a) was shown above, and we have seen that $(0, \infty)$ belongs to the resolvent set for each $j \in J$. Hence, Lemma 2.1 and the Lumer–Phillips theorem show that A_j generates a C_0 -semigroup on \mathcal{H}_j which is contractive for $j \in \{I, IV, V\}$. Part c) is deduced from the generation property by a standard calculation.

Let $j \in \{II, VI\}$. The operator A_j has no proper generalized eigenvector for $\lambda = 0$ since $\ker A_j \cap \text{ran } A_j = \{0\}$. Moreover, the projections P_j from Lemma 2.1 map onto $\ker A_j$ and commute with A_j . They thus coincide with the spectral projection of A_j for $\{0\}$ which implies the assertion $\sigma(A_j^1) = \sigma(A_j) \setminus \{0\}$. Clearly, $T_j(\cdot)$ leaves invariant the decomposition $\mathcal{H}_j = \text{ran } A_j \oplus \ker A_j$ and its restrictions to $\text{ran } A_j$ and $\ker A_j$ are generated by A_j^1 and 0, respectively. Since A_j^1 is dissipative for the equivalent norm $\|\cdot\|_E$ on $\text{ran } A_j$ by Lemma 2.1, the semigroup $T_j(\cdot)$ is bounded on $\text{ran } A_j$ and hence on \mathcal{H}_j .

2) We next prove part c) also for $j = III$. Take a number $\epsilon > 0$ such that Δ_{III} is non-zero on $B(0, 2\epsilon) \setminus \{0\}$. The spectral projection P_{III} is given by

$$P_{III}f = \frac{1}{2\pi i} \int_{|\lambda|=\epsilon} R(\lambda, A_{III})f \, d\lambda = \frac{1}{2\pi i} \int_{|\lambda|=\epsilon} \begin{pmatrix} \alpha(\lambda) \cosh(\lambda(\cdot+1)/c_1) \\ \lambda\alpha(\lambda) \cosh(\lambda(\cdot+1)/c_1) \\ \beta(\lambda) \cosh(\lambda(\cdot-1)/c_2) \\ \lambda\beta(\lambda) \cosh(\lambda(\cdot-1)/c_2) \end{pmatrix} d\lambda.$$

for $f \in \mathcal{H}_{III}$, where we use (2.8)–(2.10) and Cauchy’s integral theorem and formula. (Observe that the integrands of U_λ and W_λ vanish at $\lambda = 0$.)

Equation (2.10) next yields

$$\begin{pmatrix} \alpha(\lambda) \\ \beta(\lambda) \end{pmatrix} = \frac{1}{\lambda^2} \varphi(\lambda) z(\lambda),$$

where the factors are given by $\varphi(\lambda) = \lambda/\Delta_{III}(\lambda)$ for $\lambda \in B(0, 2\epsilon) \setminus \{0\}$ and

$$z_1(\lambda) = \cosh(\lambda/c_2) (-T_1 U'_\lambda(0) + T_2 W'_\lambda(0)) + a_2 \sinh(\lambda/c_2) (-\lambda U_\lambda(0) + \lambda W_\lambda(0) - kT_1 U'_\lambda(0) + f_1(0) - f_3(0))$$

$$z_2(\lambda) = (\cosh(\lambda/c_1) + a_1 k \sinh(\lambda/c_1)) (-T_1 U'_\lambda(0) + T_2 W'_\lambda(0)) - a_1 \sinh(\lambda/c_1) (-\lambda U_\lambda(0) + \lambda W_\lambda(0) - kT_1 U'_\lambda(0) + f_1(0) - f_3(0))$$

for $\lambda \in \mathbb{C}$. Note that z is holomorphic on \mathbb{C} . The power series of \sinh and \cosh imply the expansion

$$\Delta_{III}(\lambda) = \left(\frac{a_1}{c_1} + \frac{a_2}{c_2} \right) \lambda + \frac{a_1 a_2 k}{c_1 c_2} \lambda^2 + \mathcal{O}(\lambda^3) = (m_1 + m_2) \lambda + k m_1 m_2 \lambda^2 + \mathcal{O}(\lambda^3)$$

near $\lambda = 0$, so that φ has a holomorphic extension at 0 with $\varphi(0) = \frac{1}{m_1 + m_2}$. Moreover, its derivative is given by

$$\begin{aligned} \varphi'(\lambda) &= \frac{\Delta_{III}(\lambda) - \lambda \Delta'_{III}(\lambda)}{\Delta_{III}(\lambda)^2} = \frac{-k m_1 m_2 \lambda^2 + \mathcal{O}(\lambda^3)}{(m_1 + m_2)^2 \lambda^2 + \mathcal{O}(\lambda^3)}, \\ \varphi'(0) &= \frac{-k m_1 m_2}{(m_1 + m_2)^2}. \end{aligned}$$

By means of Cauchy's integral formula, we can now evaluate the above expression for $P_{III}f$ and derive

$$\begin{aligned} P_{III}f &= \left(\frac{d}{d\lambda} \right) \Big|_{\lambda=0} \begin{pmatrix} \varphi(\lambda) z_1(\lambda) \cosh(\lambda(\cdot+1)/c_1) \\ \lambda \varphi(\lambda) z_1(\lambda) \cosh(\lambda(\cdot+1)/c_1) \\ \varphi(\lambda) z_2(\lambda) \cosh(\lambda(\cdot-1)/c_2) \\ \lambda \varphi(\lambda) z_2(\lambda) \cosh(\lambda(\cdot-1)/c_2) \end{pmatrix} \\ &= \frac{-k m_1 m_2}{(m_1 + m_2)^2} \begin{pmatrix} z_1(0) \mathbb{1} \\ 0 \\ z_2(0) \mathbb{1} \\ 0 \end{pmatrix} + \frac{1}{m_1 + m_2} \begin{pmatrix} 0 \\ z_1(0) \mathbb{1} \\ 0 \\ z_2(0) \mathbb{1} \end{pmatrix} + \frac{1}{m_1 + m_2} \begin{pmatrix} z'_1(0) \mathbb{1} \\ 0 \\ z'_2(0) \mathbb{1} \\ 0 \end{pmatrix}. \end{aligned} \quad (2.14)$$

To compute the vectors $z(0)$ and $z'(0)$, we first infer from (2.8) the identities

$$\begin{aligned} U'_\lambda(0) &= -\frac{1}{c_1^2} \int_{-1}^0 \cosh(\lambda r/c_1) (\lambda f_1(r) + f_2(r)) dr, \\ W'_\lambda(0) &= \frac{1}{c_2^2} \int_0^1 \cosh(\lambda r/c_1) (\lambda f_3(r) + f_4(r)) dr, \end{aligned} \quad (2.15)$$

and hence

$$z(0) = \begin{pmatrix} \frac{T_1}{c_1^2} \int_{-1}^0 f_2 dr + \frac{T_2}{c_2^2} \int_0^1 f_4 dr \\ \frac{T_1}{c_1^2} \int_{-1}^0 f_2 dr + \frac{T_2}{c_2^2} \int_0^1 f_4 dr \end{pmatrix} = \begin{pmatrix} m_1 \int_{-1}^0 f_2 dr + m_2 \int_0^1 f_4 dr \\ m_1 \int_{-1}^0 f_2 dr + m_2 \int_0^1 f_4 dr \end{pmatrix}.$$

We further compute

$$\frac{d}{d\lambda} U'_\lambda(0) = -\frac{1}{c_1^2} \int_{-1}^0 \left(\frac{1}{c_1} \sinh\left(\frac{\lambda r}{c_1}\right) (\lambda f_1(r) + f_2(r)) + \cosh\left(\frac{\lambda r}{c_1}\right) f_1(r) \right) dr,$$

$$\begin{aligned}\frac{d}{d\lambda}U'_\lambda(0)|_{\lambda=0} &= -\frac{1}{c_1^2} \int_{-1}^0 f_1 dr, \\ \frac{d}{d\lambda}W'_\lambda(0)|_{\lambda=0} &= \frac{1}{c_2^2} \int_0^1 f_3 dr.\end{aligned}$$

As a result,

$$\begin{aligned}z'_1(0) &= m_1 \int_{-1}^0 f_1 dr + m_2 \int_0^1 f_3 dr + km_1 m_2 \int_{-1}^0 f_2 dr + m_2 f_1(0) - m_2 f_3(0), \\ z'_2(0) &= m_1 \int_{-1}^0 f_1 dr + m_2 \int_0^1 f_3 dr + km_1 m_2 \int_0^1 f_4 dr - m_1 f_1(0) + m_1 f_3(0).\end{aligned}$$

Therefore a function $f \in \mathcal{H}_{III}$ belongs to $\ker P_{III}$ if and only if $0 = z_1(0) = z'_1(0) = z'_2(0)$; i.e.,

$$\begin{aligned}m_1 \int_{-1}^0 f_2 dr + m_2 \int_0^1 f_4 dr &= 0, \\ \frac{m_1 + m_2}{m_2} \int_{-1}^0 f_1 dr + \frac{m_1 + m_2}{m_1} \int_0^1 f_3 dr &= 0, \\ f_1(0) - f_3(0) + km_1 \int_{-1}^0 f_2 dr &= 0.\end{aligned}\tag{2.16}$$

We can now show that the norm $\|\cdot\|_E$ is equivalent to $\|\|\cdot\|\|_E$ on $\ker P_{III}$. Let $f \in \ker P_{III}$. We use the equations

$$f_l(x) = f_l(0) + \int_0^x f'_l(r) dr \tag{2.17}$$

for $l \in \{1, 3\}$ and $x \in [-1, 0]$, resp. $x \in [0, 1]$. Formulas (2.16) then yield

$$\begin{aligned}\frac{m_1 + m_2}{m_2} f_1(0) + \frac{m_1 + m_2}{m_1} f_3(0) &= -\frac{m_1 + m_2}{m_2} \int_{-1}^0 \int_0^x f'_1(r) dr dx \\ &\quad - \frac{m_1 + m_2}{m_1} \int_0^1 \int_0^x f'_3(r) dr dx, \\ f_1(0) - f_3(0) &= -km_1 \int_{-1}^0 f_2 dr.\end{aligned}$$

Here the coefficient matrix in front of the vector $(f_1(0), f_3(0))$ is invertible. We can thus bound $|f_1(0)|$ and $|f_3(0)|$ by $c\|f\|_{\mathcal{H}}$ for a constant $c > 0$. The claimed equivalence then easily follows from (2.17).

As in (2.3) we see that A_{III} is dissipative for $\|\cdot\|_E$. Its restriction A_{III}^1 to $\ker P_{III}$ thus generates a bounded semigroup that coincides with the restriction of $T_{III}(\cdot)$ to $\ker P_{III}$. Moreover, for $w_0 \in \text{ran } P_{III}$ the solution $T_{III}(t)w_0$ is a linear combination of functions $(\mathbb{1}, 0, 0, 0)$, $(0, 0, \mathbb{1}, 0)$, and $t(0, \mathbb{1}, 0, \mathbb{1})$ by Lemma 2.1. Altogether we have shown assertion c).

3) We now look at the purely imaginary non-zero eigenvalues of A_{III} . By part a), $is \in \mathbb{R} \setminus \{0\}$ is an eigenvalue if and only if

$$\begin{aligned}0 = \Delta_{III}(is) &= -a_1 a_2 k \sin\left(\frac{s}{c_1}\right) \sin\left(\frac{s}{c_2}\right) \\ &\quad + i[a_2 \cos\left(\frac{s}{c_1}\right) \sin\left(\frac{s}{c_2}\right) + a_1 \sin\left(\frac{s}{c_1}\right) \cos\left(\frac{s}{c_2}\right)].\end{aligned}$$

Hence, s/c_1 or s/c_2 have to be a zero of \sin . In the first case, the second summand of $\text{Im } \Delta_{III}(is)$ vanishes and $\cos(s/c_1)$ has absolute value one. As a result, also $\sin(s/c_2)$ has to be zero. Here we can interchange the role of s/c_1 and s/c_2 , so that both numbers must belong to $\pi\mathbb{Z}$. This means that

$is \in \mathbb{R} \setminus \{0\}$ is an eigenvalue of A_{III} if and only if $s = c_1 k \pi = c_2 l \pi$ for some integers k and l , which is equivalent to $d \in \mathbb{Q}$.

4) Let d be irrational. The spectral assertions in d) follow from part a) as well as formula (4.4) of [19], Theorem 1 of [12] for $j = IV$, and step 3) for $j = III$. Let $f \in \mathcal{H}_j$, $s \in \mathbb{R} \setminus \{0\}$, and $w = R(is, A_j)f$. Observe that $\sinh(is) = i \sin s$ and $\cosh(is) = \cos s$. We first take $j \in \{I, IV, V\}$. Formulas (2.9)–(2.12) and (2.15) with $\lambda = is$ imply the inequalities

$$\begin{aligned} \|w\|_E &\leq c (\|w'_1\|_{L^2} + \|w_2\|_{L^2} + \|w'_3\|_{L^2} + \|w_4\|_{L^2}) \\ &\leq c (|s\alpha(is)| + \|U'_{is}\|_{L^2} + \|sU_{is}\|_{L^2} + \|f_1\|_{L^2} + |s\beta(is)| + \|W'_{is}\|_{L^2} \\ &\quad + \|sW_{is}\|_{L^2} + \|f_3\|_{L^2}) \\ &\leq \frac{c}{|\Delta(is)|} (\|U'_{is}\|_\infty + \|W'_{is}\|_\infty + \|sU_{is}\|_\infty + \|sW_{is}\|_\infty + \|f_1\|_\infty + \|f_3\|_\infty) \\ &\quad + (\|U'_{is}\|_{L^2} + \|sU_{is}\|_{L^2} + \|f_1\|_{L^2} + \|W'_{is}\|_{L^2} + \|sW_{is}\|_{L^2} + \|f_3\|_{L^2}). \end{aligned} \quad (2.18)$$

Here and below, the constants $c > 0$ only depend on the given constants. Combined with equation (2.8), Lemma 3.3 of [7] shows that

$$\|sU_{is}\|_\infty, \|U'_{is}\|_\infty \leq c (\|f_1\|_{H^1} + \|f_2\|_{L^2}).$$

The maps W_{is} are treated similarly. Hence, (2.13) is true for $j \in \{I, IV, V\}$. In Cases II, III, and VI one has to add the norms $\|w_1\|_{L^2}$ and $\|w_3\|_{L^2}$ in (2.18), which can be absorbed using $|s| \geq 1$. \square

For completeness we also state the results from [19] and [12] for *rational* $d = c_2/c_1$. We write $d = o/e$ if $d = p/q$ for coprime integers with odd p and even q , and analogously for the other two cases. Purely imaginary eigenvalues of A_j exist if and only if

$$\begin{array}{llll} \text{Case I:} & d = o/o, & \text{Case II:} & d = o/e, & \text{Case III:} & d \in \mathbb{Q}, \\ \text{Case IV:} & d \in \mathbb{Q}, & \text{Case V:} & d = e/o, & \text{Case VI:} & d = o/o. \end{array} \quad (2.19)$$

(Case III was treated in the proof of Proposition 2.2.)

3. THE CORE ESTIMATES AND IRRATIONALITY MEASURES

We now consider irrational $d = c_2/c_1 > 0$, so that $\Delta_j(is) \neq 0$ for $s \in \mathbb{R} \setminus \{0\}$ by Proposition 2.2. Because of (2.13), we need lower bounds for $|\Delta_j(is)|$ to control the growth of the resolvent of A_j along $i\mathbb{R}$. Such bounds depend on the rate of simultaneous convergence of sequences (s_n/c_1) and (s_n/c_2) to the set of zeros of sine or cosine as $|s_n| \rightarrow \infty$. We are thus led to certain number theoretic properties of the ratio $d = c_2/c_1$.

Let $r > 0$ be irrational. We define its *irrationality measure* $\mu(r)$ as the supremum of all $\theta > 0$ such that

$$\exists \text{ infinitely many } p, q \in \mathbb{N} \text{ with } \left| r - \frac{p}{q} \right| < q^{-\theta}, \quad (3.1)$$

see e.g. Appendix E in [10] or Chapter XI in [13]. Theorem 1 of [25] or Exercise E.1 in [10] give a formula for $\mu(r)$ in terms of the continued fraction of r . The larger $\mu(r)$ the better one can approximate r by rationals and, as we see below, the faster $\Delta_j(is_n)$ will tend to 0 for certain $|s_n| \rightarrow \infty$.

A famous result by Roth [22] says that $\mu(r) = 2$ for algebraic r . One always has $\mu(r) \geq 2$, see Theorem 185 in [13]. It is known that $\mu(e) = 2$,

$\mu(\pi) \leq 7.6063$, and $\mu(\ln 2) \leq 3.5746$, for instance; see [9], [23], resp. [21]. There are numbers with $\mu(r) = \infty$, see Example 1 of [25] or §11.7 in [13]. One can construct r with a prescribed irrationality measure due to Corollary 4 of [25]. We have $\mu(r) = 2$ for a.e. r by Theorem E.3 in [10].

However, the eigenvalue results (2.19) indicate that maybe one should not look for approximations by any rationals, but only by those in one of the classes odd/odd, odd/even, or even/odd. We thus define the restricted irrationality measures $\mu_{oo}(r)$, $\mu_{oe}(r)$, and $\mu_{eo}(r)$ as the supremum of all $\theta > 0$ satisfying (3.1) with $p/q = o/o$, $p/q = o/e$, respectively $p/q = e/o$. Clearly, these numbers are less or equal $\mu(r)$, and at least one of them has to be equal to $\mu(r)$. It is known that $\mu_{oo}(r), \mu_{oe}(r), \mu_{eo}(r) \geq 2$, and hence they are all equal to $\mu(r)$ if $\mu(r) = 2$, see [24] or Theorem II in [17].

The continued fraction $[a_0, a_1, \dots]$ of r allows to relate these numbers. We first recall that only the convergents of r can approach it faster than quadratically, see Theorem 184 in [13]. If all quotients a_n are even integers, then the convergents belong to only two of the classes o/o , o/e and e/o (determined by initial a_n), see Lemmas 1 and 2 of [17]. So, if e.g. o/o is left out and $\mu(r) > 2$, then $\mu_{oo}(r) = 2$ and one of the other two classes is equal to $\mu(r) > 2$. We set

$$\begin{aligned} \text{Case I: } \quad \mu_I &= \mu_{oo}(d), & \text{Case II: } \quad \mu_{II} &= \mu_{oe}(d), \\ \text{Case III: } \quad \mu_{III} &= \mu(d), & \text{Case IV: } \quad \mu_{IV} &= \mu(d), \\ \text{Case V: } \quad \mu_V &= \mu_{eo}(d), & \text{Case VI: } \quad \mu_{VI} &= \mu_{oo}(d). \end{aligned} \quad (3.2)$$

The next lemma contains the core estimate to treat the case of irrational ratios d . It also shows that this estimate is almost optimal.

Lemma 3.1. *Let $j \in J$, $d = c_2/c_1$ be irrational, and μ_j in (3.2) be finite. Take any $\eta > 0$. Then the following assertions are true.*

a) *There exists a constant $c_\eta > 0$ such that*

$$|\Delta_j(is)| \geq \frac{c_\eta}{|s|^{(2\mu_j-2)+\eta}} \quad (3.3)$$

for all $s \in \mathbb{R}$ with $|s| \geq 1$.

b) *There are numbers $\tilde{s}_n \in [1, \infty)$ with $\tilde{s}_n \rightarrow \infty$ and a constant $\tilde{c} > 0$ with*

$$|\Delta_j(i\tilde{s}_n)| \leq \frac{\tilde{c}}{|s|^{(2\mu_j-2)-\eta}} \quad (3.4)$$

for all $n \in \mathbb{N}$. In (3.4) we can take $\eta = 0$ if $\mu_j = 2$.

Proof. We do not relabel subsequences of (s_n) , and write κ or κ_l for positive constants only depending on the given numbers T_j , m_j , and k .

1a) For $\lambda = is$ and $s \in \mathbb{R}$, we can rewrite the maps Δ_j from (2.12) as

$$\begin{aligned} \Delta_I(is) &= a_1 a_2 k \cos\left(\frac{s}{c_1}\right) \cos\left(\frac{s}{c_2}\right) + i[a_2 \sin\left(\frac{s}{c_1}\right) \cos\left(\frac{s}{c_2}\right) + a_1 \cos\left(\frac{s}{c_1}\right) \sin\left(\frac{s}{c_2}\right)], \\ \Delta_{II}(is) &= -a_2 \sin\left(\frac{s}{c_1}\right) \sin\left(\frac{s}{c_2}\right) + a_1 \cos\left(\frac{s}{c_1}\right) \cos\left(\frac{s}{c_2}\right) + i a_1 a_2 k \cos\left(\frac{s}{c_1}\right) \sin\left(\frac{s}{c_2}\right), \\ \Delta_{III}(is) &= -a_1 a_2 k \sin\left(\frac{s}{c_1}\right) \sin\left(\frac{s}{c_2}\right) + i[a_2 \cos\left(\frac{s}{c_1}\right) \sin\left(\frac{s}{c_2}\right) + a_1 \sin\left(\frac{s}{c_1}\right) \cos\left(\frac{s}{c_2}\right)], \\ \Delta_{IV}(is) &= -k \sin\left(\frac{s}{c_1}\right) \sin\left(\frac{s}{c_2}\right) + i[a_1 \cos\left(\frac{s}{c_1}\right) \sin\left(\frac{s}{c_2}\right) + a_2 \sin\left(\frac{s}{c_1}\right) \cos\left(\frac{s}{c_2}\right)], \\ \Delta_V(is) &= a_1 \cos\left(\frac{s}{c_1}\right) \cos\left(\frac{s}{c_2}\right) - a_2 \sin\left(\frac{s}{c_1}\right) \sin\left(\frac{s}{c_2}\right) + i k \sin\left(\frac{s}{c_1}\right) \cos\left(\frac{s}{c_2}\right), \\ \Delta_{VI}(is) &= k \cos\left(\frac{s}{c_1}\right) \cos\left(\frac{s}{c_2}\right) + i[a_1 \sin\left(\frac{s}{c_1}\right) \cos\left(\frac{s}{c_2}\right) + a_2 \cos\left(\frac{s}{c_1}\right) \sin\left(\frac{s}{c_2}\right)]. \end{aligned} \quad (3.5)$$

Suppose that condition (3.3) does not hold. Then there exists a number $\bar{\eta} \in (0, 1/3)$ and a sequence (s_n) in \mathbb{R} with $|s_n| \geq 1$ and $|s_n| \rightarrow \infty$ such that

$$|\Delta_j(is_n)| \leq |s_n|^{-(2\mu_j-2)-3\bar{\eta}} \quad (3.6)$$

for all $n \in \mathbb{N}$ and some j .

We write $\Delta_j(is) = \Delta_j^{(1)}(is) + \Delta_j^{(2)}(is)$, where $\Delta_j^{(1)}(is)$ is the real part of $\Delta_j(is)$ if $j \in \{I, III, IV, VI\}$ and the imaginary part if $j \in \{II, V\}$. In each case, the formulas (3.5) yield the expression

$$\Delta_j^{(1)}(is_n) = \pm \kappa_1 \varphi_j(s_n/c_1) \psi_j(s_n/c_2), \quad (3.7)$$

for functions $\varphi_j, \psi_j \in \{\sin, \cos\}$ and a non-specified sign. Estimate (3.6) then implies that at least (a subsequence of) one of the sequences (s_n/c_1) and (s_n/c_2) approaches the set of zeros $N(1, j)$ of φ_j , respectively $N(2, j)$ of ψ_j . Here we have

$$\begin{array}{ll} \text{Case I:} & N(1, I) = \pi\mathbb{Z} + \frac{\pi}{2}, & N(2, I) = \pi\mathbb{Z} + \frac{\pi}{2}, \\ \text{Case II:} & N(1, II) = \pi\mathbb{Z} + \frac{\pi}{2}, & N(2, II) = \pi\mathbb{Z}, \\ \text{Case III:} & N(1, III) = \pi\mathbb{Z}, & N(2, III) = \pi\mathbb{Z}, \\ \text{Case IV:} & N(1, IV) = \pi\mathbb{Z}, & N(2, IV) = \pi\mathbb{Z}, \\ \text{Case V:} & N(1, V) = \pi\mathbb{Z}, & N(2, V) = \pi\mathbb{Z} + \frac{\pi}{2}, \\ \text{Case VI:} & N(1, VI) = \pi\mathbb{Z} + \frac{\pi}{2}, & N(2, VI) = \pi\mathbb{Z} + \frac{\pi}{2}. \end{array}$$

We now look at $\Delta_j^{(2)}(is_n)$ which is in all cases of the form

$$\Delta_j^{(2)}(is_n) = \tau_j^1 \kappa_2 \tilde{\varphi}_j(s_n/c_1) \psi_j(s_n/c_2) + \tau_j^2 \kappa_3 \varphi_j(s_n/c_1) \tilde{\psi}_j(s_n/c_2), \quad (3.8)$$

where $\tau_j^l \in \{1, -1\}$, as well as $\tilde{\varphi}_j = \sin$ if $\varphi_j = \cos$ and analogously in the other cases. If $(\varphi_j(s_n/c_1))$ tends to 0, the second summand in $\Delta_j^{(2)}(is_n)$ vanishes and the numbers $|\tilde{\varphi}_j(s_n/c_1)|$ converge to 1 as $n \rightarrow \infty$. Therefore also $\psi_j(s_n/c_2)$ has to approach $N(2, j)$. The roles of φ_j and ψ_j can be interchanged here. Hence, $\varphi_j(s_n/c_1)$ and $\psi_j(s_n/c_2)$ tend to 0 as $n \rightarrow \infty$.

We can thus find non-zero integers k_n and l_n satisfying $|k_n|, |l_n| \rightarrow \infty$ and null sequences (δ_n) and (ϵ_n) bounded by $\pi/2$ such that

$$\frac{s_n}{c_1} = k_n \pi + \xi_j \frac{\pi}{2} + \delta_n \quad \text{and} \quad \frac{s_n}{c_2} = l_n \pi + \zeta_j \frac{\pi}{2} + \epsilon_n \quad (3.9)$$

for all $n \in \mathbb{N}$. Here we have set

$$\begin{array}{ll} \text{Case I:} & \xi_I = 1, \quad \zeta_I = 1, & \text{Case II:} & \xi_{II} = 1, \quad \zeta_{II} = 0, \\ \text{Case III:} & \xi_{III} = 0, \quad \zeta_{III} = 0, & \text{Case IV:} & \xi_{IV} = 0, \quad \zeta_{IV} = 0, \\ \text{Case V:} & \xi_V = 0, \quad \zeta_V = 1, & \text{Case VI:} & \xi_{VI} = 1, \quad \zeta_{VI} = 1. \end{array} \quad (3.10)$$

It follows

$$\begin{aligned} c_1 k_n \pi + c_1 \xi_j \frac{\pi}{2} + c_1 \delta_n &= c_2 l_n \pi + c_2 \zeta_j \frac{\pi}{2} + c_2 \epsilon_n, \\ \frac{2k_n + \xi_j}{2l_n + \zeta_j} - d &= \frac{2}{\pi(2l_n + \zeta_j)} (d\epsilon_n - \delta_n). \end{aligned}$$

The assumption on d now implies that

$$\left| \frac{2k_n + \xi_j}{2l_n + \zeta_j} - d \right| \geq \frac{1}{|2l_n + \zeta_j|^{\mu_j + \bar{\eta}}}$$

for all $n \geq n_\eta$ and some index $n_\eta \in \mathbb{N}$, cf. (3.1), (3.2) and (3.10). After dropping finitely many members of the sequences, we then obtain the inequality

$$|d\epsilon_n - \delta_n| \geq \frac{\bar{c}}{|l_n|^{\mu_j-1+\eta}} \quad (3.11)$$

for all $n \in \mathbb{N}$, $j \in J$, and a constant $\bar{c} > 0$.

c) We claim that the sequences (δ_n) and (ϵ_n) satisfy the estimates

$$0 < \liminf_{n \rightarrow \infty} \frac{|\epsilon_n|}{|\delta_n|} \leq \limsup_{n \rightarrow \infty} \frac{|\epsilon_n|}{|\delta_n|} < \infty, \quad (3.12)$$

$$|\epsilon_n|, |\delta_n| \geq \frac{\bar{c}'}{|l_n|^{\mu_j-1+\eta}} \quad (3.13)$$

for a constant $\bar{c}' > 0$ and all $n \in \mathbb{N}$. We suppose that this claim was wrong.

First, there could be a subsequence such that $\lim_{n \rightarrow \infty} \frac{|\epsilon_n|}{|\delta_n|} = 0$. Because of (3.11), the numbers δ_n then have to satisfy (3.13) for all sufficiently large $n \in \mathbb{N}$ and some $\bar{c}' > 0$. Observe that $|\varphi_j(s_n/c_1)| \geq |\delta_n|/2$ and $|\psi_j(s_n/c_2)| \leq |\epsilon_n|$ for all sufficiently large n by (3.9), (3.10) and the Taylor series of sin and cos at their zeros. Similarly, the numbers $|\tilde{\varphi}_j(s_n/c_1)|$ and $|\tilde{\psi}_j(s_n/c_2)|$ tend to 1 as $n \rightarrow \infty$ and are thus contained in $[1/2, 1]$ starting from some index n_0 . We now use these facts and (3.9) to infer from (3.8) the lower bound

$$|\Delta_j(is_n)| \geq |\Delta_j^{(2)}(is_n)| \geq \frac{\kappa_3}{4} |\delta_n| - \kappa_2 |\epsilon_n| \geq \frac{\kappa_3}{8} |\delta_n| \geq \frac{\kappa_3 \bar{c}'}{8 |l_n|^{\mu_j-1+\eta}} \geq \frac{\kappa \bar{c}'}{|s_n|^{\mu_j-1+\eta}}$$

for all sufficiently large n . But this inequality contradicts (3.6).

Second, if $\lim_{n \rightarrow \infty} \frac{|\epsilon_n|}{|\delta_n|} = \infty$ was true for a subsequence, we infer a contradiction in a similar way. Hence, the relations (3.12) are fulfilled, and then (3.11) implies the inequality (3.13).

d) As we see in step 2), in $\Delta_j^{(2)}(is_n)$ cancellations may occur. So we look at the term $\Delta_j^{(1)}(is_n)$. Since the arguments approach the zero set of the functions φ_j and ψ_j , from (3.7) and (3.13) we derive the lower bound

$$|\Delta_j(is_n)| \geq |\Delta_j^{(1)}(is_n)| \geq \frac{\kappa_1}{4} |\delta_n| |\epsilon_n| \geq \frac{\kappa_1 (\bar{c}')^2}{4 |l_n|^{2\mu_j-2+2\eta}} \geq \frac{\kappa (\bar{c}')^2}{|s_n|^{2\mu_j-2+2\eta}}$$

as above for all $n \in \mathbb{N}$. But this inequality cannot be true because of (3.6). So assertion (3.3) is shown.

2a) By the assumption on d , for all $\eta > 0$ and $j \in J$ there are $k_n, l_n \in \mathbb{N}$ such that

$$|d_n| \leq \frac{1}{|l_n|^{\mu_j-\eta/2}}, \quad \text{where} \quad d_n := \frac{2k_n + \xi_j}{2l_n + \zeta_j} - d, \quad (3.14)$$

for all $n \in \mathbb{N}$. Due to [24] we can put here $\eta = 0$ if $\mu_j(d) = 2$.

To indicate the cases, we now denote the (positive) prefactors κ_2 and κ_3 of $\Delta_j^{(2)}(is)$ in (3.8) by α_j and β_j , respectively. (Only α_{II} and β_V differ from the corresponding factors in (3.5), which are equal to $-\alpha_{II}$ and $-\beta_V$, respectively.) Recall the numbers ξ_j and ζ_j from (3.10). We set

$$b_j := d + \frac{\alpha_j}{\beta_j}, \quad \delta_n := \left(\frac{d}{b_j} - 1 \right) \frac{\pi}{2} d_n (2l_n + \zeta_j), \quad \epsilon_n := \frac{\pi}{2b_j} d_n (2l_n + \zeta_j) \quad (3.15)$$

for $n \in \mathbb{N}$. By the estimate (3.14) there exists a constant $\tilde{c} > 0$ with

$$|\delta_n|, |\epsilon_n| \leq \frac{\tilde{c}}{|l_n|^{\mu_j - 1 - \eta/2}} \quad (3.16)$$

for every $n \in \mathbb{N}$, so that $|\delta_n|, |\epsilon_n| \leq \pi/2$ for all large n . Following (3.9), we next define the numbers

$$\tilde{s}_n := c_1 k_n \pi + c_1 \xi_j \frac{\pi}{2} + c_1 \delta_n. \quad (3.17)$$

Equations (3.15) then yield the formulas

$$\tilde{s}_n = c_2 l_n \pi + c_2 \zeta_j \frac{\pi}{2} + c_2 \epsilon_n \quad \text{and} \quad \alpha_j \epsilon_n + \beta_j \delta_n = 0 \quad (3.18)$$

for all $n \in \mathbb{N}$.

b) As in part 1c) we estimate

$$|\Delta_j^{(1)}(i\tilde{s}_n)| \leq \kappa_1 |\varphi_j(\tilde{s}_n/c_1)| |\psi_j(\tilde{s}_n/c_2)| \leq \kappa_1 |\delta_n| |\epsilon_n| \leq \frac{\kappa_1 \tilde{c}^2}{|l_n|^{2\mu_j - 2 - \eta}}, \quad (3.19)$$

for all $n \in \mathbb{N}$, using (3.7), (3.17), (3.18) and (3.16). In $\Delta_j^{(2)}(i\tilde{s}_n)$ from (3.8) our construction will lead to a cancellation. Set

$$\sigma_n^{(1)} := \begin{cases} 1, & k_n \text{ even,} \\ -1, & k_n \text{ odd,} \end{cases} \quad \sigma_n^{(2)} := \begin{cases} 1, & l_n \text{ even,} \\ -1, & l_n \text{ odd,} \end{cases} \quad \rho_j := \begin{cases} 1, & j \in \{III, IV\}, \\ -1, & \text{else,} \end{cases}$$

for $n \in \mathbb{N}$. We insert (3.17), (3.18) and (3.10) into the respective formulas in (3.5). Applying carefully the translation rules for sine and cosine, we arrive at the expression

$$\Delta_j^{(2)}(i\tilde{s}_n) = \rho_j \sigma_n^{(1)} \sigma_n^{(2)} [\alpha_j \cos(\delta_n) \sin(\epsilon_n) + \beta_j \sin(\delta_n) \cos(\epsilon_n)].$$

The power series for sine and cosine, (3.18), and (3.16) thus imply

$$|\Delta_j^{(2)}(i\tilde{s}_n)| \leq |\alpha_j \epsilon_n + \beta_j \delta_n| + O(|\delta_n|^3 + |\epsilon_n|^3) = O(|l_n|^{-3\mu_j + 3 + 3\eta/2}).$$

So the term (3.19) dominates in $\Delta(i\tilde{s}_n) = \Delta_j^{(1)}(i\tilde{s}_n) + \Delta_j^{(2)}(i\tilde{s}_n)$ and yields assertion (3.4). \square

4. MAIN RESULTS

Lemma 3.1 and Proposition 2.2 imply the crucial resolvent estimates. Throughout we use the standard norm of \mathcal{H} .

Proposition 4.1. *Let $j \in J$, $d = c_2/c_1$ be irrational, and μ_j in (3.2) be finite. Take any $\eta > 0$. Then the following assertions are true.*

a) *There exists a constant $C_\eta > 0$ such that*

$$\|R(is, A)\| \leq C_\eta |s|^{(2\mu_j - 2) + \eta} \quad (4.1)$$

for all $s \in \mathbb{R}$ with $|s| \geq 1$.

b) *There are numbers $\tilde{s}_n \in [1, \infty)$ with $\tilde{s}_n \rightarrow \infty$ and a constant $\tilde{C} > 0$ with*

$$\|R(i\tilde{s}_n, A)\| \geq \tilde{C} |s|^{(2\mu_j - 2) - \eta} \quad (4.2)$$

for all $n \in \mathbb{N}$. In (4.2) we can take $\eta = 0$ if $\mu_j = 2$.

Proof. The first assertion follows directly from (2.13) and (3.3). For the lower estimate, we take the numbers $\tilde{s}_n \rightarrow \infty$ from Lemma 3.1. Let $n \in \mathbb{N}$. To bound the norm of the resolvent, we use the functions

$$\tilde{\chi}_n : [-1, 0] \rightarrow \mathbb{R}; \quad \tilde{\chi}_n(x) = \begin{cases} \cos(\tilde{s}_n x / c_1), & j \in \{I, II, III\}, \\ \sin(\tilde{s}_n x / c_1), & j \in \{IV, V, VI\}, \end{cases} \quad \chi_n = \|\tilde{\chi}_n\|_{L^2}^{-1} \tilde{\chi}_n$$

and write $w^n = R(i\tilde{s}_n, A)(0, \chi_n, 0, 0)$. Since $\|\chi_n\|_{L^2} = 1$, it follows

$$\|R(i\tilde{s}_n, A)\| \geq \|w^n\|_{\mathcal{H}} \geq \|w_2^n\|_{L^2}. \quad (4.3)$$

In each case, formulas (2.9) and (2.8) yield

$$w_2^n(x) = i\tilde{s}_n w_1^n(x) = i\tilde{s}_n \alpha(i\tilde{s}_n) g_j\left(\frac{i\tilde{s}_n}{c_1} x\right) - \frac{i}{c_1} \int_{-1}^x \sin\left(\frac{\tilde{s}_n}{c_1}(x-r)\right) \chi_n(r) dr$$

for $x \in [-1, 0]$. By Lemma 3.3 of [7], the second summand is uniformly bounded by $c \|\chi_n\|_{L^2} = c$. The first one is given by

$$G_{n,j}(x) := i\tilde{s}_n \alpha(i\tilde{s}_n) g_j\left(\frac{i\tilde{s}_n}{c_1} x\right) = \begin{cases} i\tilde{s}_n \alpha(i\tilde{s}_n) \cos\left(\frac{\tilde{s}_n}{c_1}(x+1)\right), & j \in \{III, VI\}, \\ -\tilde{s}_n \alpha(i\tilde{s}_n) \sin\left(\frac{\tilde{s}_n}{c_1}(x+1)\right), & \text{else.} \end{cases}$$

Using the notation in (3.8), from formulas (2.10), (2.8), and (2.15) we derive the lower bound

$$\begin{aligned} |\tilde{s}_n \alpha(i\tilde{s}_n)| &\geq \frac{1}{|\Delta_j(i\tilde{s}_n)|} \left(\left| \tilde{\psi}_j\left(\frac{\tilde{s}_n}{c_2}\right) \frac{T_1}{c_1^2} \int_{-1}^0 \cos\left(\frac{\tilde{s}_n}{c_1} r\right) \chi_n(r) dr \right| \right. \\ &\quad \left. - \left| a_2 \tilde{\psi}_j\left(\frac{\tilde{s}_n}{c_2}\right) \left[\frac{-i}{c_1} \int_{-1}^0 \sin\left(\frac{\tilde{s}_n}{c_1} r\right) \chi_n(r) dr + \frac{kT_1}{c_1^2} \int_{-1}^0 \cos\left(\frac{\tilde{s}_n}{c_1} r\right) \chi_n(r) dr \right] \right| \right) \end{aligned}$$

for $j \in \{I, II, III\}$ and $x \in [-1, 0]$; whereas (2.11), (2.8), and (2.15) yield

$$\begin{aligned} |\tilde{s}_n \alpha(i\tilde{s}_n)| &\geq \frac{1}{|\Delta_j(i\tilde{s}_n)|} \left(\left| a_2 \tilde{\psi}_j\left(\frac{\tilde{s}_n}{c_2}\right) \frac{-i}{c_1} \int_{-1}^0 \sin\left(\frac{\tilde{s}_n}{c_1} r\right) \chi_n(r) dr \right| \right. \\ &\quad \left. - \left| \tilde{\psi}_j\left(\frac{\tilde{s}_n}{c_2}\right) \left[\frac{T_1}{c_1^2} \int_{-1}^0 \cos\left(\frac{\tilde{s}_n}{c_1} r\right) \chi_n(r) dr - \frac{ik}{c_1} \int_{-1}^0 \cos\left(\frac{\tilde{s}_n}{c_1} r\right) \chi_n(r) dr \right] \right| \right) \end{aligned}$$

for $j \in \{IV, V, VI\}$.

In the above expressions the integrals are bounded by $\|\chi_n\|_{L^2} = 1$. As in part 1c) of the proof of Lemma 3.1 the numbers $|\tilde{\psi}_j(\frac{\tilde{s}_n}{c_2})|$ tend to 1 and $\tilde{\psi}_j(\frac{\tilde{s}_n}{c_2})$ to 0 as $n \rightarrow \infty$. We thus obtain an index $n_0 \in \mathbb{N}$ such that

$$|\tilde{s}_n \alpha(i\tilde{s}_n)| \geq \frac{\kappa}{|\Delta_j(i\tilde{s}_n)| \|\tilde{\chi}_n\|_{L^2}} \int_{-1}^0 \tilde{\chi}_n(r)^2 dr = \frac{\kappa}{|\Delta_j(i\tilde{s}_n)|} \|\tilde{\chi}_n\|_{L^2}$$

for all $n \geq n_0$. Moreover, a substitution and equation (3.17) lead to the inequality

$$\begin{aligned} \int_{-1}^0 \cos^2\left(\frac{\tilde{s}_n}{c_1} r\right) dr &= \frac{c_1}{\tilde{s}_n} \int_{-\tilde{s}_n/c_1}^0 \cos^2(t) dt \geq \frac{1}{\pi(k_n + 1)} \int_0^{k_n \pi - \pi/2} \cos^2(t) dt \\ &= \frac{2k_n - 1}{\pi(k_n + 1)} \int_0^{\pi/2} \cos^2(t) dt \geq \gamma > 0, \end{aligned}$$

for some $\gamma > 0$ and all n , and analogously

$$\int_{-1}^0 \cos^2\left(\frac{\tilde{s}_n}{c_1}(r+1)\right) dr \geq \tilde{\gamma} > 0.$$

Similar inequalities are valid if we replace here \cos by \sin . Summing up, we have shown that

$$\|G_{n,j}\|_{L^2} \geq \frac{c}{|\Delta_j(i\tilde{s}_n)|}$$

for all $j \in J$, all sufficiently large n and a constant $c > 0$ independent of n . Lemma 3.1 and (4.3) then imply (4.2). \square

The above estimates and the results [6] and [8] imply our main theorem. Recall that $2\mu_j - 2 \geq 2$ and that here equality holds for a.e. d by Theorem E.3 in [10]. Moreover, $T_{III}(t)P_{III}w_0$ is constant in x and affine in t .

Theorem 4.2. *Let $j \in J$, $d = c_2/c_1$ be irrational, μ_j in (3.2) be finite, and $T_j(\cdot)$ the C_0 -semigroup on \mathcal{H}_j generated by A_j . Take any $\beta \in (0, (2\mu_j - 2)^{-1})$ and $\gamma > (2\mu_j - 2)^{-1}$. There are constants $c_\beta, c > 0$ and times $t_n = t_n(j) \geq 1$ tending to ∞ such that the following assertions hold.*

a) *Let $j \in \{I, IV, V\}$. The semigroups then satisfies*

$$\begin{aligned} \|T_j(t)A_j^{-1}\| &\leq c_\beta t^{-\beta} && \text{for all } t \geq 1, \\ \|T_j(t_n)A_j^{-1}\| &\geq ct_n^{-\gamma} && \text{for all } n \in \mathbb{N}. \end{aligned}$$

b) *Let $j \in \{II, III, VI\}$ and P_j be the spectral projection of A_j for $\{0\}$ from Proposition 2.2. For $j \in \{II, VI\}$ we obtain*

$$\begin{aligned} \|T_j(t)w_0 - P_j w_0\| &\leq c_\beta t^{-\beta} \|A_j w_0\| && \text{for all } t \geq 1, w_0 \in D(A_j), \\ \|T_j(t_n) - P_j\|_{\mathcal{B}(D(A_j), \mathcal{H}_j)} &\geq ct_n^{-\gamma} && \text{for all } n \in \mathbb{N}. \end{aligned}$$

Moreover, we have

$$\begin{aligned} \|T_{III}(t)w_0 - T_{III}(t)P_{III}w_0\| &\leq c_\beta t^{-\beta} \|A_{III}(I - P_{III})w_0\|, \\ \|T_{III}(t_n)(I - P_{III})\|_{\mathcal{B}(D(A_{III}), \mathcal{H})} &\geq ct_n^{-\gamma}, \end{aligned}$$

for alle $t \geq 1$, $w_0 \in D(A_{III})$, and $n \in \mathbb{N}$.

Proof. Let $j \in \{I, IV, V\}$. Part a) follows directly from Theorem 2.4 of [8] and Proposition 4.1 since $T_j(\cdot)$ is bounded on \mathcal{H}_j and $\sigma(A_j) \cap i\mathbb{R} = \emptyset$ by Proposition 2.2.

Let $j \in \{II, III, VI\}$. In these cases we use the restriction A_j^1 of A_j to $\ker P_j \subseteq \mathcal{H}_j$ which has the analogous properties as the generators in part a). The restriction $T_j^1(\cdot)$ of $T_j(\cdot)$ to $\ker P_j$ is generated by A_j^1 , see Proposition 2.2, and hence satisfies estimates as in a). For $j \in \{II, VI\}$ we further have $T_j(t)P_j = P_j$ and $A_j P_j = 0$. The result follows. \square

In the above theorem we do not quite get the decay rate $\beta = (2\mu_j - 2)^{-1}$ since μ_j was defined by a supremum. At least for a.e. d we can close this gap up to a logarithmic correction. Theorem 32 in [16] implies that for each $\eta > 0$ we have

$$\left|d - \frac{p}{q}\right| \geq \frac{1}{q^2 \ln^{1+\eta} q} \quad (4.4)$$

for a.e. irrational $d > 0$ and all rationals p/q with $q \geq q_\eta$ for some $q_\eta \in \mathbb{N}$. On the other hand, for a.e. irrational d one finds fractions p_n/q_n with

$$\left|d - \frac{p_n}{q_n}\right| \leq \frac{1}{q_n^2 \ln q_n}$$

for all $n \in \mathbb{N}$, so that (4.4) is essentially sharp.

Theorem 4.3. *For a.e. irrational $d = c_2/c_1$ and all $\epsilon > 0$ there is a constant c_ϵ such that*

$$\begin{aligned} \|T_j(t)A_j^{-1}\| &\leq c_\epsilon \frac{\ln(t)^{1+\epsilon}}{\sqrt{t}}, & j \in \{I, IV, V\}, \\ \|T_j(t)w_0 - P_j w_0\| &\leq c_\epsilon \frac{\ln(t)^{1+\epsilon}}{\sqrt{t}} \|A_j w_0\|, & j \in \{II, VI\}, \\ \|T_{III}(t)w_0 - T_{III}(t)P_{III}w_0\| &\leq c_\epsilon \frac{\ln(t)^{1+\epsilon}}{\sqrt{t}} \|A_{III}(I - P_{III})w_0\|, \end{aligned}$$

for all $t \geq 1$ and $w_0 \in D(A_j)$, where P_j is the spectral projection of A_j for $\{0\}$ from Proposition 2.2.

Proof. As in Lemma 3.1 one shows the lower bound

$$|\Delta_j(is)| \geq \frac{c_\eta}{s^2 \ln(|s|)^{2+2\eta}} \quad (4.5)$$

for all $|s| \geq 1$, $j \in J$ and $\eta > 0$. To this aim, we suppose that the opposite inequality is true for some $\bar{\eta} > 0$ and a sequence (s_n) with $|s_n| \rightarrow \infty$, cf. (3.6). We use (4.4) to replace (3.11) by

$$|d\epsilon_n - \delta_n| \geq \frac{\bar{c}}{|l_n| \ln(|l_n|)^{1+\bar{\eta}/2}}.$$

Arguing as in Lemma 3.1, we then arrive at a contradiction and infer (4.5).

Proposition 2.2 now yields the estimate $\|R(is, A_j)\| \leq c'_\eta s^2 \ln(|s|)^{2+2\eta}$ for $|s| \geq 1$. Based on Theorem 1.3 of [5], we finally derive the assertions as in the previous theorem. \square

REFERENCES

- [1] K. Ammari, M. Tucsnak and A. Henrot, *Optimal location of the actuator for the pointwise stabilization of a string*. C. R. Acad. Sci. Paris Sér. I Math. **330** (2000), 275–280.
- [2] N. Anantharaman and M. Léautaud, *Sharp polynomial decay rates for the damped wave equation on the torus*. Anal. PDE **7** (2014), 159–214.
- [3] A. Bátkai, K.-J. Engel, J. Prüss and R. Schnaubelt, *Polynomial stability of operator semigroups*. Math. Nachr. **279** (2006), 1425–1440.
- [4] C.J.K. Batty, A. Borichev and Y. Tomilov, *L^p -tauberian theorems and L^p -rates for energy decay*. J. Funct. Anal. **270** (2016), 1153–1201.
- [5] C.J.K. Batty, R. Chill and Y. Tomilov, *Fine scales of decay of operator semigroups*. J. Eur. Math. Soc. **18** (2016), 853–929.
- [6] C.J.K. Batty and T. Duyckaerts, *Non-uniform stability for bounded semi-groups on Banach spaces*. J. Evol. Equ. **8** (2008), 765–780.
- [7] C.J.K. Batty, L. Paunonen and D. Seifert, *Optimal energy decay in a one-dimensional coupled wave-heat system*. J. Evol. Equ. **16** (2016), 649–664.
- [8] A. Borichev and Y. Tomilov, *Optimal polynomial decay of functions and operator semigroups*. Math. Ann. **347** (2010), 455–478.
- [9] J.M. Borwein and P.B. Borwein, *Pi and the AGM. A Study in Analytic Number Theory and Computational Complexity*. John Wiley, 1987.
- [10] Y. Bugeaud, *Distribution Modulo One and Diophantine Approximation*. Cambridge University Press, 2012.
- [11] G. Chen, M. Coleman and H.H. West, *Pointwise stabilization in the middle of the span for second order systems nonuniform and uniform exponential decay of solutions*. SIAM J. Appl. Math. **47** (1987), 751–780.

- [12] B.Z. Guo and W.D. Zhu, *On the energy decay of two coupled strings through a joint damper*. Journal of Sound and Vibration 203 (1997), 447-455.
- [13] G.H. Hardy and E.M. Wright, *An Introduction to the Theory of Numbers*. 6th edition. Oxford University Press, 2008.
- [14] S. Jaffard, M. Tucsnak, E. Zuazua, *Singular internal stabilization of the wave equation*. J. Differential Equations **145** (1998), 184-215.
- [15] O. Kaviani and Q. Zhang, *Polynomial stabilization of solutions to a class of damped wave equations*. Preprint (2017), arXiv:1703.01735.
- [16] A.Ya. Khinchin, *Continued Fractions*. The University of Chicago Press, 1964.
- [17] L. Kuipers and B. Meulenbeld, *Some properties of continued fractions*. Acta Math. **87** (1952). 1-12.
- [18] G. Lebeau, *Équation des ondes amorties*. In: A. Boutet de Monvel and V. Marchenko (Eds.), "Algebraic and Geometric Methods in Mathematical Physics," Kluwer, 1996, pp. 73-109.
- [19] K.-S. Liu, *Energy decay problems in the design of a point stabilizer for coupled string vibrating systems*. SIAM J. Control Optim. **26** (1988), 1348-1356.
- [20] Z. Liu and B. Rao, *Characterization of polynomial decay rate for the solution of linear evolution equation*. Z. Angew. Math. Phys. **56** (2005), 630-644.
- [21] R. Marcovecchio, *The Rhin-Viola method for $\log 2$* . Acta Arith. **139** (2009), 147-184.
- [22] K.F. Roth, *Rational approximations to algebraic numbers*. Mathematika **2** (1955), 1-20.
- [23] V. Kh. Salikhov, *On the irrationality measure of π* . Russian Math. Surveys **63** (2008), 570-572.
- [24] W.T. Scott, *Approximation to real irrationals by certain classes of rational fractions*. Bull. Amer. Math. Soc. **46** (1940), 124-129.
- [25] J. Sondow, *Irrationality measures, irrationality basis, and a theorem of Jarník*. Preprint, arXiv:math/0406300.
- [26] M. Tucsnak, *On the pointwise stabilization of a string*. In: W. Desch, F. Kappel and K. Kunisch (Eds.), "Control and Estimation of Distributed Parameter Systems," Birkhäuser, 1998, pp. 287-295,

Ł. RZEPNICKI, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, NICOLAUS COPERNICUS UNIVERSITY, TORUŃ, POLAND.

E-mail address: keleb@mat.umk.pl

R. SCHNAUBELT, DEPARTMENT OF MATHEMATICS, KARLSRUHE INSTITUTE OF TECHNOLOGY, 76128 KARLSRUHE, GERMANY.

E-mail address: schnaubelt@kit.edu