

Multidimensional Thermoelasticity for Nonsimple Materials – Well-Posedness and Long-Time Behavior

Andrii Anikushyn, Michael Pokojov

CRC Preprint 2016/11, May 2016

KARLSRUHE INSTITUTE OF TECHNOLOGY

CRC 1173



Participating universities



Universität Stuttgart

EBERHARD KARLS
UNIVERSITÄT
TÜBINGEN



Funded by

DFG

ISSN 2365-662X

Multidimensional Thermoelasticity for Nonsimple Materials – Well-Posedness and Long-Time Behavior

Andrii Anikushyn*

Michael Pokojovy†

May 6, 2016

Abstract

An initial-boundary value problem for the multidimensional type III thermoelasticity for a nonsimple material with a center of symmetry is considered. In the linear case, the well-posedness with and without Kelvin-Voigt and/or frictional damping in the elastic part as well as the lack of exponential stability in the elastically undamped case is proved. Further, a frictional damping for the elastic component is shown to lead to the exponential stability. A Cattaneo-type hyperbolic relaxation for the thermal part is introduced and the well-posedness and uniform stability under a nonlinear frictional damping are obtained using a compactness-uniqueness-type argument. Additionally, a connection between the exponential stability and exact observability for unitary C_0 -groups is established.

Key words: thermoelasticity, nonsimple materials, semilinear systems, well-posedness, uniform stability, hyperbolic relaxation

AMS: 35B40; 35G46; 35G61; 35Q74; 74F05; 93D20

1 Introduction

In modern rational mechanics, when describing a material body, the general principles such as field equations or jump conditions are separated from the so-called constitutive equations or material laws (cf. [25, pp. 1–2]). Whereas the former are common among all representatives of a major class of materials such as solids or fluids, the latter are meant to uniquely characterize each particular material. Typically, a material law describes the response of a material to various stimuli applied and is modelled by an algebraic or an operator equation (see, e.g., [25, Chapter C3]). In macroscopic theories such as the classical theory of (thermo)elasticity or its modern generalizations and unifications such

*Faculty of Cybernetics, Taras Shevchenko National University of Kyiv, Kyiv, Ukraine
anik_andrii@univ.kiev.ua

†Department of Mathematics, Karlsruhe Institute of Technology, Karlsruhe, Germany
michael.pokojovy@kit.edu

as the one proposed by Green and Naghdi [5, 6, 7], etc., the microscopic structure of the material is ignored and the material law is obtained in the form of a stress-strain relation, which can be measured experimentally. Recently, Rajagopal [20] made a strong case for using implicit material laws, which are physically more sound as they let the stress, i.e., the force, induce the strain and not vice versa. In anelastic bodies, e.g., the thermoelastic ones, the entropy and the resulting irreversibility play a very important role. In the theory of thermoelasticity, a relation between the heat flux and the temperature gradient is postulated such that the second theorem of thermodynamics, usually in form of Clausius-Duhem inequality, is satisfied [25, Sections 96 and 96 bis]. Alternatively, an entropy balance equation can be used as proposed by Green and Naghdi [6, 7].

When being applied to various materials with microstructure, the macroscopic theory of (thermo)elasticity does not provide an adequate mechanical and thermodynamical description. They include but are not limited to porous elastic media, micropolar elastic solids, materials with microstructure and nonsimple elastic solids, which were first introduced in the works of Truesdell and Toupin [26], Green and Rivlin [8], Mindlin [15] and Toupin [24]. Their common feature is that they incorporate extra field variables such as microstresses or microrotations, hyperstresses or volume fractions, etc. For further details, we refer the reader to comprehensive monographs by Ciarletta and Ieşan [2] and Ieşan [10].

We now briefly summarize the linear thermoelasticity model for nonsimple materials without energy dissipation introduced by Quintanilla [19]. To this end, consider a rigid body occupying in a reference configuration a bounded domain Ω of \mathbb{R}^d with the Lipschitz boundary $\Gamma := \partial\Omega$. For a period of time $t \geq 0$ and a material point $\mathbf{x} = (x_i) \in \Omega$, let $\mathbf{u} = (u_i)$, (u_{iL}) and T , being functions \mathbf{x} and t , denote the displacement vector, the homogeneous deformation tensor of the ‘particle’ with its center of mass located at \mathbf{x} and the relative temperature measured with respect to a constant reference temperature T_0 , which is assumed to be attained at some time $t_0 \geq 0$. Further, we define the thermal displacement

$$\tau(\mathbf{x}, t) := \int_{t_0}^t T(\mathbf{x}, s) ds.$$

With $\mathbf{t} = (t_{Kj})$ denoting the first Piola & Kirchhoff stress tensor, the linear balance of momentum reads as

$$\rho \ddot{u}_i = t_{K^i, K} + \rho f_i,$$

where ρ stands for the material density and $\mathbf{f} = (f_i)$ is the volumetric force. Here and the sequel, we employ the Einstein’s summation convention as well as the standard notation for temporal and spatial derivatives of scalar and tensor fields (cf. [10, Chapter 1]). In contrast to the classical linear theory of thermoelasticity, which utilizes the (generalized) Hooke’s law to postulate a linear relation between the elastic part of \mathbf{t} and the infinitesimal Cauchy strain tensor

$$\boldsymbol{\varepsilon} = \frac{1}{2}(\nabla \mathbf{u}^T + \nabla \mathbf{u}),$$

the hyperstresses need to be accounted for. The latter can be shown to incorporate higher order derivatives of \mathbf{u} (cf. [10, Chapter 7.1]). Following Quintanilla [19, Sections 1 and 2], the linear equations of thermoelasticity without energy dissipation for a simple material

with a center of symmetry read as

$$\rho \ddot{u}_i = (A_{iJRs} u_{s,R} - \beta_{Ji} T - (C_{iJKSRl} u_{l,RS} + M_{iJKR\tau,R}),_K)_{,J} + \rho f_i, \quad (1)$$

$$a \ddot{\tau} = -\beta_{Ki} \dot{u}_{i,K} - M_{jLKI} u_{j,LKI} + K_{IJ} \tau_{,IJ} + \rho T_0^{-1} R \quad (2)$$

in $\Omega \times (0, \infty)$, where R stands for the volumetric heat sources. We refer the reader to [19, Section 2] for an explanation of material constants in Equations (1)–(2) as well as a discussion on their symmetry and positive definiteness properties. The homogeneous Dirichlet-Dirichlet boundary conditions for Equations (1)–(2) on Γ are given by

$$u_i = 0, \quad u_{i,A} = 0, \quad \tau = 0 \text{ in } \Gamma \times (0, \infty), \quad (3)$$

whereas the initial conditions are stated as

$$u_i(\cdot, 0) = u_i^0, \quad \dot{u}_i(\cdot, 0) = \dot{u}_i^1, \quad \tau(\cdot, 0) = \tau^0, \quad T(\cdot, 0) = T^0 \text{ in } \Omega. \quad (4)$$

In [19], Quintanilla proposed a logarithmically convex energy-like function to prove the uniqueness for Equations (1)–(4). He further employed the operator semigroup theory to obtain the existence of solutions.

Fernández Sare et al. [3] considered a one-dimensional counterpart of Equations (1)–(4) formulated in terms of Green & Naghdi's type I thermoelasticity (cf. [6]) under various sets of boundary conditions. Based on the operator semigroup theory and Gearhart & Prüss' theorem, the well-posedness and exponential stability of solution were shown. Further, the spectral analyticity criterion was employed to show the lack of analyticity for the underlying semigroup. Finally, the authors proved the impossibility of the solutions to localize in time. Similar results have later been obtained by Magaña and Quintanilla [14] also for the case of the more comprehensive type III thermoelasticity.

Pata and Quintanilla [16] considered a 3D version of Equation (1) with $T \equiv 0$ for the case vanishing anti-plane shear deformations (i.e., $u_1 = u_2 = 0$). The Equation for $u := u_3$ reduces then to

$$\rho \ddot{u} = \mu(0) \Delta u - l_2(0) \Delta^2 u + \int_0^\infty (\mu'(s) \Delta u(t-s) - l_2'(s) \Delta^2 u(t-s)) ds = 0 \quad (5)$$

together with the boundary condition

$$u = \Delta u = 0 \quad (6)$$

and the initial condition

$$u(\cdot, -s) = g(\cdot, s) \text{ for } s \in [0, \infty). \quad (7)$$

First, an abstract version of the initial-boundary value problem (5)–(7) was studied using the operator semigroup theory. Based on a modification of Gearhart & Prüss' theorem, a condition for the exponential stability was derived.

Gawinecki and Łazuka [4] considered a Cauchy problem for the genuinely nonlinear version of Equations (1)–(2) within Green & Naghdi's type I thermoelasticity for homogeneous isotropic media. Under appropriate conditions on the nonlinearity, a global classical solutions was obtained based on L^p - L^q -estimates.

Aouadi and Moulahi [1] studied an initial-boundary value problem for the anti-plane share component of a nonsimple thermoelastic body with a control distributed over an open subset ω of Ω

$$u_{tt} - c^2 \Delta u + \alpha \Delta^2 u + c^2 \gamma (-\Delta)^{1/2} \theta = \chi_\omega u_1, \quad (8)$$

$$\theta_t - \Delta \theta - \gamma (-\Delta)^{1/2} u_t = \chi_\omega u_2 \quad (9)$$

subject to the homogeneous Dirichlet-Dirichlet boundary conditions and the standard initial conditions. Here, χ_ω stands for the indicator function of ω . For the uncontrolled case, i.e., $u_1 = u_2 \equiv 0$, a well-posedness and exponential stability result for Equations (8)–(9) was proved. Further, the authors showed the system is approximately controllable over the standard state space by $L^2(0, T; L^2(\Omega))$ -controls at any time $T > 0$ if ω is nonempty.

In the present paper, we first consider Equations (1)–(4) restated within the type III thermoelasticity (cf. [14, Section 2] for the 1D system) with a (macroscopic) Kelvin-Voigt damping for \mathbf{u} . Hence, our equations read as

$$\rho \ddot{u}_i = (A_{iJKj} u_{j,K} - \beta_{Ji} \dot{\tau} - (C_{iJKLIj} u_{j,IL} + M_{iJKL} \tau_{,L} - B_{iJKj} \dot{u}_j)_{,K})_{,J} \quad (10)$$

$$a \dot{\tau} = -\beta_{Ki} \dot{u}_{i,K} + m_{IJ} \dot{\tau}_{,IJ} + M_{jLKI} u_{j,LKI} + K_{IJ} \tau_{,IJ}. \quad (11)$$

together with the boundary and initial conditions in Equations (3)–(4). Throughout the paper, the following (natural) positivity and positive definiteness conditions are postulated:

I. $a, \rho > 0$.

II. There exists $\alpha > 0$ such that

$$C_{iJKLj} u_{i,JI} u_{j,LK} + M_{iJKL} u_{i,KJ} \tau_{,L} + M_{jLKI} u_{j,LK} \tau_{,I} + K_{IJ} \tau_{,IJ} \tau_{,J} \geq \alpha (\tau_{,R} \tau_{,R} + u_{k,ST} u_{k,ST}), \quad (12)$$

$$A_{iKLj} u_{i,K} u_{j,L} \geq \alpha u_{k,R} u_{k,R}. \quad (13)$$

In Section 2, we show Equations (10)–(11), (3)–(4) are well-posed. Note that no positive definiteness conditions on (B_{iJKj}) or (m_{IJ}) are imposed. By a standard perturbation argument (e.g., [17, Chapter 3.1, Theorem 1.1]), the problem remains well-posed if the Kelvin-Voigt damping is replaced or complemented with a frictional damping.

In Section 3, we prove that, in contrast to the 1D situation (cf. [14, Section 2]), if $B_{iJKj} \equiv 0$, the positive definiteness of (m_{IJ}) is not sufficient to exponentially stabilize the system. Motivated by the necessity of an additional damping for \mathbf{u} , we let $B_{iJKj} \equiv 0$, but add a linear frictional damping to Equation (10) thus obtaining

$$\rho \ddot{u}_i = (A_{iJKj} u_{j,K} - \beta_{Ji} \dot{\tau} - (C_{iJKLIj} u_{j,IL} + M_{iJKL} \tau_{,L} - B_{iJKj} \dot{u}_j)_{,K})_{,J} - E_{ij} \dot{u}_j. \quad (14)$$

Further, in Section 4, under a positive definiteness assumption on (E_{ij}) , we use the Lyapunov's method to show the exponential stability of Equations (14), (11), (3)–(4).

In Section 5, a hyperbolization of (11) together a semilinear frictional damping in Equation (10) are considered. The resulting semilinear system reads then as

$$\rho \ddot{u}_i = (A_{iJKj} u_{j,K} - \beta_{Ji} \dot{\tau} - (C_{iJKLIj} u_{j,IL} + M_{iJKL} \tau_{,L})_{,K})_{,J} - E(|\dot{u}|) \dot{u}_i, \quad (15)$$

$$a\ddot{\tau} = -\beta_{K_i}\dot{u}_{i,K} + m_{IJ}q_{I,J} + M_{jLKI}u_{j,LKI} + K_{IJ}\tau_{,IJ}, \quad (16)$$

$$\kappa\dot{q}_i = \dot{\tau}_{,i} - q_i, \quad (17)$$

with a small parameter $\kappa > 0$. Here, $(m_{IJ}q_{I,J})$ plays the role of the heat flux. If the function E is constant, similar to the case $\kappa = 0$, Equations (15)–(17) are exponentially stable. In the nonlinear case, we exploit a technique due to Lasiecka and Tataru [12] to prove the uniform stability of the nonlinear system dependent on the behavior of E at 0 and infinity. Here, we use a generalization of a technique dating back to Haraux [9], which is given in the appendix A.

2 Linear sytem with a Kelvin-Voigt damping for the elastic part: Well-posedness

In this section, we consider the equations of type III thermoelasticity for a nonsimple material with a Kelvin-Voigt damping for \mathbf{u} . By a standard perturbation argument [17, Chapter 3.1, Theorem 1.1], the results remain valid also when a frictional damping, i.e., a term like $E_{ij}\dot{u}_j$, is considered instead of or in addition to $B_{iJKj}\dot{u}_{j,KJ}$. The equations read then as

$$\rho\ddot{u}_i = (A_{iJKj}u_{j,K} - \beta_{Ji}\dot{\tau} - (C_{iJKLIj}u_{j,IL} + M_{iJKL}\tau_{,L} - B_{iJKj}\dot{u}_j)_{,K})_{,J} \quad (18)$$

$$a\ddot{\tau} = -\beta_{K_i}\dot{u}_{i,K} + m_{IJ}\dot{\tau}_{,IJ} + M_{jLKI}u_{j,LKI} + K_{IJ}\tau_{,IJ} \quad (19)$$

in $\Omega \times (0, \infty)$ together with the boundary conditions

$$u_i = 0, \quad u_{i,J} = 0, \quad \tau = 0 \text{ in } \Gamma \times (0, \infty) \quad (20)$$

and the initial conditions

$$u_i(\cdot, 0) = u_i^0, \quad \dot{u}_i(\cdot, 0) = \dot{u}_i^0, \quad \tau(\cdot, 0) = \tau^0, \quad \dot{\tau}(\cdot, 0) = \dot{\tau}^0 \text{ in } \Omega, \quad (21)$$

where $\Gamma := \partial\Omega$ is assumed Lipschitzian. In addition to conditions I, II, we assume

$$B_{iJKj}\xi_{j,K}\xi_{i,J} \geq 0, \quad m_{IJ}\xi_I\xi_J \geq 0. \quad (22)$$

Letting $v = \dot{u}$ and $\theta = \dot{\tau}$, we rewrite the system (18)–(19) as follows:

$$\dot{u}_i = v_i, \quad (23)$$

$$\dot{v}_i = \frac{1}{\rho}(A_{iJKj}u_{j,K} - \beta_{Ji}\theta - (C_{iJKLIj}u_{j,IL} + M_{iJKL}\tau_{,L} - B_{iJKj}v_j)_{,K})_{,J}, \quad (24)$$

$$\dot{\tau} = \theta, \quad (25)$$

$$\dot{\theta} = \frac{1}{a}(-\beta_{K_i}v_{i,K} + (m_{IJ}\theta_{,J} + M_{jLKI}u_{j,LK} + K_{IJ}\tau_{,J})_{,I}). \quad (26)$$

We assume the evolution is taking place on the Hilbert space

$$\mathcal{H} = \left\{ U \mid U = (u, v, \tau, \theta)^T \in (H_0^2(\Omega))^d \times (L^2(\Omega))^d \times H_0^1(\Omega) \times L^2(\Omega) \right\},$$

where

$$H_0^s(\Omega) := \text{clos}(C_0^\infty(\Omega), \|\cdot\|_{H^s}) \text{ for } s \in \mathbb{N},$$

equipped with the scalar product

$$\begin{aligned} \langle U, U^* \rangle_{\mathcal{H}} &= \int_{\Omega} (\rho v_i v_i^* + a \theta \theta^* + A_{iKjL} u_{i,K} u_{j,L}^* + C_{iJKL} u_{i,JI} u_{j,LK}^* + K_{IJ} \tau_{,J} \tau_{,I}^*) dx \\ &\quad + \int_{\Omega} (M_{iJKL} u_{i,KJ} \tau_{,L}^* + M_{jLKI} u_{j,KL}^* \tau_{,I}) dx, \end{aligned} \quad (27)$$

for $U = (u, v, \tau, \theta)^T$, $U^* = (u^*, v^*, \tau^*, \theta^*)^T \in \mathcal{H}$. Here and in the following, for the sake of simplicity, we drop the physical convention to bold the vector and tensor fields. It is easy to verify that this scalar product is equivalent with the usual product induced by the product topology.

Consider the linear operator $\mathcal{A} : D(\mathcal{A}) \subset \mathcal{H} \rightarrow \mathcal{H}$ defined by

$$\mathcal{A}U = \begin{pmatrix} v_i, \\ \frac{1}{\rho}(A_{iJKj} u_{j,K} - \beta_{Ji} \theta - (C_{iJKLIj} u_{j,IL} + M_{iJKL} \tau_{,L} - B_{iJKj} v_j)_{,K})_{,J}, \\ \theta, \\ \frac{1}{a}(-\beta_{Ki} v_{i,K} + (m_{IJ} \theta_{,J} + M_{jLKI} u_{j,LK} + K_{IJ} \tau_{,J})_{,I}) \end{pmatrix}, \quad (28)$$

where

$$\begin{aligned} D(\mathcal{A}) = \{U \in \mathcal{H} \mid & v_i \in H_0^2(\Omega), \theta \in H_0^1(\Omega), \\ & (A_{iJKj} u_{j,K} - \beta_{Ji} \theta)_{,J} \\ & - (C_{iJKLIj} u_{j,IL} M_{iJKL} \tau_{,L} - B_{iJKj} v_j)_{,K} \}_{,J} \in L^2(\Omega), \\ & (m_{IJ} \theta_{,J} + M_{jLKI} u_{j,LK} + K_{IJ} \tau_{,J})_{,I} \in L^2(\Omega)\} \end{aligned} \quad (29)$$

Thus, the abstract form of Equations (23)–(26) is given by

$$\dot{U}(t) = \mathcal{A}U(t) \text{ for } t > 0, \quad U(0) = U_0. \quad (30)$$

with $U_0 := (u_i^0, u_i^1, \tau^0, \tau^1)^T$. We first prove the following auxiliary lemma.

Lemma 1. *The operator \mathcal{A} defined in Equations (28)–(29) is the infinitesimal generator of a C_0 -semigroup of contractions on \mathcal{H} .*

Proof. The proof of Lemma 1 resembles the one of [18, Theorem 2].

Denseness: Utilizing the fact that

$$(C_0^\infty(\Omega))^d \times (C_0^\infty(\Omega))^d \times C_0^\infty(\Omega) \times C_0^\infty(\Omega)$$

is a subset of the domain of \mathcal{A} , we conclude that $D(\mathcal{A})$ is dense in \mathcal{H} .

Disipativity: A straightforward calculation involving the Green's formula yields

$$\langle \mathcal{A}U, U \rangle_{\mathcal{H}} = - \int_{\Omega} (B_{iJKj} v_{i,K} v_{i,J} + m_{IJ} \theta_{,I} \theta_{,J}) dx. \quad (31)$$

Using Equation (22), we get

$$\langle \mathcal{A}U, U \rangle_{\mathcal{H}} \leq -\alpha \int_{\Omega} (v_{i,K}v_{i,K} + \theta_{,I}\theta_{,I}) dx \leq 0. \quad (32)$$

Maximality: Now we prove that $0 \in \rho(\mathcal{A})$ with $\rho(\mathcal{A})$ standing for the resolvent set of \mathcal{A} . For $F = (F^1, F^2, F^3, F^4)^T \in \mathcal{H}$, consider the operator equation $\mathcal{A}U = F$ or, explicitly,

$$\begin{aligned} v_i &= F_i^1, \\ \frac{1}{\rho} (A_{iJKj}u_{j,K} - \beta_{Ji}\theta - (C_{iJKLIj}u_{j,IL} + M_{iJKL}\tau_{,L} - B_{iJKj}v_{j,K})_{,J}) &= F_i^2, \\ \theta &= F^3, \\ \frac{1}{a} (-\beta_{Ki}v_{i,K} + (m_{IJ}\theta_{,J} + M_{jLKI}u_{j,LK} + K_{IJ}\tau_{,J})_{,I}) &= F^4. \end{aligned}$$

Eliminating v_i and θ , we obtain

$$\frac{1}{\rho} (A_{iJKj}u_{j,K} - \beta_{Ji}F^3 - (C_{iJKLIj}u_{j,IL} + M_{iJKL}\tau_{,L} - B_{iJKj}F_j^1)_{,K})_{,J} = F_i^2, \quad (33)$$

$$\frac{1}{a} (-\beta_{Ki}F_{i,K}^1 + (m_{IJ}F_{,J}^3 + M_{jLKI}u_{j,LK} + K_{IJ}\tau_{,J})_{,I}) = F^4. \quad (34)$$

To solve this system, we exploit the lemma of Lax & Milgram. We consider the Hilbert space

$$\mathcal{V} = (H_0^2(\Omega))^d \times H_0^1(\Omega)$$

equipped with the standard inner product associated with the product topology and introduce the bilinear form $\mathfrak{a} : \mathcal{V} \times \mathcal{V} \rightarrow \mathbb{R}$ via

$$\begin{aligned} \mathfrak{a}(V, V^*) &= \int_{\Omega} (A_{iJKj}u_{s,R}u_{i,J}^* + C_{iJKLj}u_{j,LK}u_{i,IJ}^*) dx \\ &+ \int_{\Omega} (M_{iJKL}\tau_{,L}u_{i,JK}^* + M_{jLKI}u_{j,KL}\tau_{,I}^* + K_{IJ}\tau_{,I}\tau_{,J}^*) dx, \end{aligned}$$

where $V = (u, \tau)^T$ and $V^* = (u^*, \tau^*)^T$. After multiplying Equations (33)–(34) in the inner product $(L^2(\Omega))^n$ and $L^2(\Omega)$ with ρW^1 and aW^2 , respectively, summing up the resulting equations and integrating by parts, we obtain a weak formulation of Equations (33)–(34) in the form:

$$\begin{aligned} \text{Determine } V \in \mathcal{V} \text{ such that} \\ \mathfrak{a}(V, W) &= -\langle G^1, W^1 \rangle_{(L^2(\Omega))^n} - \langle G^2, W^2 \rangle_{L^2(\Omega)} - m_{IJ} \langle F_{,J}^3, W_{,I}^2 \rangle_{L^2(\Omega)}. \end{aligned} \quad (35)$$

Here, $G_i^1 = \rho F_i^2 + \beta_{Ji}F_{,J}^3 - B_{iJKj}F_{j,KJ}^1$, $G^2 = aF^4 + \beta_{Ki}F_{i,K}^1$.

The bilinear form \mathfrak{a} is continuous and coercive on \mathcal{V} due to the conditions in (12)–(13). Clearly, the linear functional

$$(W^1, W^2) \mapsto \langle G^1, W^1 \rangle_{(L^2(\Omega))^n} + \langle G^2, W^2 \rangle_{L^2(\Omega)} + m_{IJ} \langle F_{,J}^3, W_{,I}^2 \rangle_{L^2(\Omega)}$$

is continuous on \mathcal{V} . Applying now the Lemma of Lax & Milgram, we deduce existence of a unique solution $V = (u, \tau)^T \in \mathcal{V}$ to Equations (33)–(34). Recalling the definition of

v_i and θ , we further get $v_i = F_i^1 \in H_0^2(\Omega)$ and $\theta = F^3 \in H_0^1(\Omega)$. Hence, the conditions in Equations (29) are satisfied. Therefore, we have $(u, v, \tau, \theta)^T \in D(\mathcal{A})$ implying that $(u, v, \tau, \theta)^T$ is a strong solution.

Finally we show the continuous dependence of the solution on F . It follows from the Lemma of Lax & Milgram that there exists $c_0 > 0$ such that

$$\|V\|_{\mathcal{V}} \leq c_0 \|\widehat{F}\|_{\mathcal{V}^*}, \quad (36)$$

where $\widehat{F} = (F^2, F^4)^T$. Since \mathcal{V} is continuously embedded into $(L^2(\Omega))^{d+1}$, we have

$$\|\widehat{F}\|_{\mathcal{V}^*} \leq c_1 \|\widehat{F}\|_{(L^2(\Omega))^{d+1}}. \quad (37)$$

Suppose $\|F_n\|_{\mathcal{H}} \rightarrow 0$ as $n \rightarrow \infty$. By definition, $\|F_n^1\|_{(H_0^2(\Omega))^d} \rightarrow 0$ and $\|F_n^3\|_{H_0^1(\Omega)} \rightarrow 0$ as $n \rightarrow \infty$ and, therefore, $\|(v_i)_n\|_{L^2(\Omega)} \rightarrow 0$ and $\|\theta_n\|_{L^2(\Omega)} \rightarrow 0$ as $n \rightarrow \infty$. Further, $\|F_n^2\|_{(L^2(\Omega))^d}, \|F_n^4\|_{L^2(\Omega)} \rightarrow 0$ as $n \rightarrow \infty$ and it follows from Equations (36)–(37) that $\|V\|_{\mathcal{V}} \rightarrow 0$. Therefore, the solution U_n of equation $\mathcal{A}U_n = F_n$ tends to 0 in \mathcal{H} as $n \rightarrow \infty$.

Now, after we have shown \mathcal{A} is m -dissipative, the claim of the present theorem follows from the Lumer & Phillips theorem (cf. [17, Theorem 4.3, p. 14]). \square

Now, by virtue of [17, Theorem 1.3, p. 102], Equations (18)–(21) as well as their abstract formulation (30) are well-posed.

Theorem 2. *Let $U_0 \in \mathcal{H}$. There exists then a unique mild solution $U \in C^0([0, \infty), \mathcal{H})$ to Equation (30). If $U_0 \in D(\mathcal{A})$, the mild solution is even a classical one satisfying $U \in C^1([0, \infty), \mathcal{H}) \cap C^0([0, \infty), D(\mathcal{A}))$.*

Applying [17, Chapter 3.1, Theorem 1.1], we further obtain

Corollary 3. *Replacing Equation (18) with*

$$\rho \ddot{u}_i = \left(A_{iJKj} u_{j,K} - \beta_{Ji} \dot{\tau} - (C_{iJKLIj} u_{j,IL} + M_{iJKL} \tau_{,L} - B_{iJKj} \dot{u}_j)_{,J} \right) + E_{ij} \dot{u}_j \quad (38)$$

for an arbitrary matrix $(E_{ij}) \in \mathbb{R}^{d \times d}$ in Equations (38), (19)–(21), the resulting abstract Cauchy problem is well-posed on \mathcal{H} .

3 Linear system with an undamped elastic part: Lack of exponential stability

In this section, we consider Equations (18)–(19) for the case $B_{iJKj} \equiv 0$, i.e.,

$$\rho \ddot{u}_i = \left(A_{iJKj} u_{j,K} - \beta_{Ji} \dot{\tau} - (C_{iJKLIj} u_{j,IL} + M_{iJKL} \tau_{,L}, \right. \quad (39)$$

$$\left. a \dot{\tau} = -\beta_{Ki} \dot{u}_{i,K} + m_{IJ} \dot{\tau}_{,IJ} + M_{jLKI} u_{j,LKI} + K_{IJ} \tau_{,IJ}. \right. \quad (40)$$

Equations (39)–(40) are precisely the type III thermoelasticity for nonsimple materials. Under a suitable choice of natural boundary conditions, i.e., the ones appearing in the

Green's formula, and a feasible selection of the coefficient tensors (A_{iJKj}) , (β_{Ji}) , (C_{iJKLIj}) , (m_{IJ}) , (K_{IJ}) as well as restricting Ω to the rectangular configuration

$$\Omega = (0, \pi) \times (0, \pi),$$

we show that the system (39)–(40) is lacking exponential stability. Selecting the relatively open in Γ disjunctive sets

$$\Gamma_1 := (0, \pi) \times \{0, \pi\} \text{ and } \Gamma_2 := \{0, \pi\} \times (0, \pi),$$

we have $\Gamma = \bar{\Gamma}_1 \cup \bar{\Gamma}_2$. The coefficient tensors (A_{iJKj}) , (β_{Ji}) , (C_{iJKLIj}) , (m_{IJ}) , (K_{IJ}) are chosen in [19, p. 5] to model an isotropic material with a center of symmetry. Further, we select

$$(m_{ij}) := \begin{pmatrix} \delta & 0 \\ 0 & \delta \end{pmatrix} \text{ for some } \delta > 0.$$

The balance equations and material laws (cf. [19]) read then as

$$\begin{aligned} \tau_{JI,J} &= \sigma_{RJI,RJ} + \rho \ddot{u}_I, \\ (\phi_I)_{,I} &= \beta \dot{e}_{LL} - D_{IJ} \dot{\tau}_{,IJ} + a \ddot{\tau}, \\ \tau_{IJ} &= \lambda \delta_{IJ} e_{LL} + 2\mu e_{IJ} - \beta \delta_{IJ} \dot{\tau}, \\ \sigma_{IJK} &= \frac{1}{2} a_1 (\delta_{JK} \kappa_{LLI} + 2\delta_{IJ} \kappa_{KLL} + \delta_{IK} \kappa_{LLJ}) + a_2 (\delta_{JI} \kappa_{ILL} + \delta_{IK} \kappa_{JLL}) \\ &\quad + 2a_3 \delta_{IJ} \kappa_{LLK} + 2a_4 \kappa_{IJK} + a_5 (\kappa_{KJI} + \kappa_{KIJ}) + m_1 \delta_{IJ} \tau_{,K} \\ &\quad + m_2 (\delta_{JK} \tau_{,I} + \delta_{IK} \tau_{,J}), \\ \phi_I &= m_1 \kappa_{LLI} + m_2 (\kappa_{ILL} + \kappa_{LIL}) + \delta_{IJ} \tau_{,J}, \\ e_{IJ} &= \frac{1}{2} (u_{I,J} + u_{J,I}), \quad \kappa_{IJK} = u_{K,IJ}. \end{aligned}$$

After plugging the constitutive equations into the balance equations, we arrive at the system

$$\begin{aligned} \rho \ddot{u}_1 &= (\lambda + 2\mu) u_{1,11} + \mu u_{1,22} + (\lambda + \mu) u_{2,12} - \beta \dot{\tau}_{,1} - b_1 u_{1,1111} - b_2 u_{1,1122} \\ &\quad - b_3 u_{1,2222} - b_4 u_{2,1112} - b_5 u_{2,1222} - (m_1 + 2m_2) (\tau_{,111} + \tau_{,122}), \end{aligned} \quad (41)$$

$$\begin{aligned} \rho \ddot{u}_2 &= (\lambda + 2\mu) u_{2,22} + \mu u_{2,11} + (\lambda + \mu) u_{1,12} - \beta \dot{\tau}_{,2} - b_1 u_{2,2222} - b_2 u_{2,1122} \\ &\quad - b_3 u_{2,1111} - b_4 u_{1,1112} - b_5 u_{1,1222} - (m_1 + 2m_2) (\tau_{,112} + \tau_{,222}), \end{aligned} \quad (42)$$

$$\begin{aligned} a \ddot{\tau} &= (m_1 + 2m_2) (u_{1,111} + u_{1,122} + u_{2,112} + u_{2,222}) + \tau_{,1} + \tau_{,2} \\ &\quad - \beta (\dot{u}_{1,1} + \dot{u}_{2,2}) + \delta (\dot{\tau}_{,11} + \dot{\tau}_{,22}), \end{aligned} \quad (43)$$

where

$$\begin{aligned} b_1 &= 2(a_1 + a_2 + a_3 + a_4 + a_5), & b_2 &= 2(a_1 + a_2 + 2a_3 + 2a_4 + a_5), \\ b_3 &= 2(a_3 + a_4), & b_4 &= b_5 = 2(a_1 + a_2 + a_5). \end{aligned}$$

Throughout this section, we employ the following set of boundary conditions, which naturally arise from partially integrating of the system (41)–(43):

$$\sigma_{111}|_{\Gamma} = a_1 (u_{1,11} + u_{1,22} + u_{1,12} + u_{2,12}) + 2a_2 (u_{1,11} + u_{2,21})$$

$$+2a_3(u_{1,11} + u_{1,22}) + 2(a_4 + a_5)u_{1,11} + (m_1 + 2m_2)\tau_{,1}|_\Gamma = 0, \quad (44)$$

$$\begin{aligned} \sigma_{221}|_\Gamma = a_1(u_{1,11} + u_{2,12}) + 2a_3(u_{1,11} + u_{1,22}) + 2a_4u_{1,22} \\ + 2a_5u_{2,12} + m_1\tau_{,1}|_\Gamma = 0, \end{aligned} \quad (45)$$

$$\begin{aligned} \sigma_{112}|_\Gamma = \frac{1}{2}a_1(u_{1,11} + u_{1,22}) + a_2(u_{1,11} + u_{2,21}) + 2a_4u_{2,12} \\ + a_5(u_{1,22} + u_{2,12}) + m_2\tau_{,1}|_\Gamma = 0, \end{aligned} \quad (46)$$

$$(\sigma_{112,1}, \sigma_{222,2}) \cdot (\nu_1, \nu_2)|_\Gamma = 0, \quad (47)$$

$$(\tau_{,1}, \tau_{,2})^T \cdot (\nu_1, \nu_2)^T|_{\Gamma_2} = 0, \tau|_{\Gamma_1} = 0, \quad (48)$$

where $\nu = (\nu_1, \nu_2)^T$ is the unit exterior normal vector to $\Gamma = \partial\Omega$. To eliminate the trivial kernel, the ground space is then selected as the Hilbert space

$$\mathcal{H}_* = (H^2(\Omega)/\{1\})^2 \times (L^2(\Omega)/\{1\})^2 \times (H^1(\Omega)/\{1\}) \times (L^2(\Omega)/\{1\})$$

i.e., each component is taken as the orthogonal complement of the one-dimensional subspace of constant functions. Without loss of generality, we equip this new space \mathcal{H}_* with the canonical unweighted inner product, which is equivalent with our original weighted definition in Equation (27).

The operator \mathcal{A} is defined similar to (28). The boundary conditions in Equations (44)–(48) are incorporated into the weak definition of the domain $D(\mathcal{A})$ similar to Equation (29).

Theorem 4. *Assume $a_3 + a_4 \geq 0$. Then the system (41)–(43) with boundary conditions (44)–(48) is not exponentially stable.*

Proof. We prove there exists a sequence $(\lambda_n)_n \subset \mathbb{R}$ with

$$\lim_{n \rightarrow \infty} |\lambda_n| = \infty$$

as well as the sequences $(U_n)_n \subset D(\mathcal{A})$ and $(F_n)_n \subset \mathcal{H}_0$ such that

$$(i\lambda_n - \mathcal{A})U_n = F_n \text{ is uniformly bounded w.r.t. } n \in \mathbb{N} \text{ and } \lim_{n \rightarrow \infty} \|U_n\|_{\mathcal{H}_*} = \infty.$$

The lack of exponential stability will then follow from the well-known Gearhart & Prüss' theorem (see e.g. [13, Theorem 1.3.2, p. 4]).

For $\lambda \in \mathbb{R}$, the solution $U = (u_1, u_2, v_1, v_2, \tau, \theta)^T$ of the resolvent equation $(\lambda i - \mathcal{A})U = F$ satisfies

$$\begin{aligned} \lambda_n u_{1i} - v_1 &= 0, \\ \lambda_n u_{2i} - v_2 &= 0, \\ \lambda_n v_{1i} - \frac{1}{\rho} &\left((\lambda + 2\mu)u_{1,11} + \mu u_{1,22} + (\lambda + \mu)u_{2,12} - \beta \dot{\tau}_{,1} - b_1 u_{1,1111} \right. \\ &\quad \left. - b_2 u_{1,1122} - b_3 u_{1,2222} - b_4 u_{2,1112} - b_5 u_{2,1222} \right. \\ &\quad \left. - (m_1 + 2m_2)(\tau_{,111} + \tau_{,122}) \right) = f_3, \\ \lambda_n v_{2i} - \frac{1}{\rho} &\left((\lambda + 2\mu)u_{2,22} + \mu u_{2,11} + (\lambda + \mu)u_{1,12} - \beta \dot{\tau}_{,2} - b_1 u_{2,2222} \right. \end{aligned}$$

$$\begin{aligned}
 & -b_2u_{2,1122} - b_3u_{2,1111} - b_4u_{1,1112} - b_5u_{1,1222} \\
 & \quad - (m_1 + 2m_2)(\tau_{,112} + \tau_{,222}) = f_4, \\
 & \quad \lambda_n \tau_i - \theta = 0, \\
 & \lambda_n \theta_i - \frac{1}{a} \left((m_1 + 2m_2)(u_{1,111} + u_{1,122} + u_{2,112} + u_{2,222}) \right. \\
 & \quad \left. + \tau_{,1} + \tau_{,2} - \beta(\dot{u}_{1,1} + \dot{u}_{2,2}) + \delta(\dot{\tau}_{,11} + \dot{\tau}_{,22}) \right) = f_6,
 \end{aligned}$$

where f_3, f_4, f_6 will be selected as

$$f_3 = \sin(nx) \sin(ny), \quad f_4 = \cos(nx) \cos(ny), \quad f_6 = \cos(nx) \sin(ny)$$

for $n \in \mathbb{N}$. Eliminating v_1, v_2, θ , we obtain for u_1, u_2, τ the following algebraic system

$$\begin{aligned}
 & -\lambda_n^2 u_1 - \frac{1}{\rho} \left((\lambda + 2\mu)u_{1,11} + \mu u_{1,22} + (\lambda + \mu)u_{2,12} \right. \\
 & \quad \left. - \beta \lambda_n i \tau_{,1} - b_1 u_{1,1111} - b_2 u_{1,1122} - b_3 u_{1,2222} \right. \\
 & \quad \left. - b_4 u_{2,1112} - b_5 u_{2,1222} - (m_1 + 2m_2)(\tau_{,111} + \tau_{,122}) \right) = f_3, \tag{49}
 \end{aligned}$$

$$\begin{aligned}
 & -\lambda_n^2 u_2 - \frac{1}{\rho} \left((\lambda + 2\mu)u_{2,22} + \mu u_{2,11} + (\lambda + \mu)u_{1,12} \right. \\
 & \quad \left. - \beta \lambda_n i \tau_{,2} - b_1 u_{2,2222} - b_2 u_{2,1122} - b_3 u_{2,1111} \right. \\
 & \quad \left. - b_4 u_{1,1112} - b_5 u_{1,1222} - (m_1 + 2m_2)(\tau_{,112} + \tau_{,222}) \right) = f_4, \tag{50}
 \end{aligned}$$

$$\begin{aligned}
 & -\lambda_n^2 \theta_i - \frac{1}{a} \left((m_1 + 2m_2)(u_{1,111} + u_{1,122} + u_{2,112} + u_{2,222}) \right. \\
 & \quad \left. + \tau_{,1} + \tau_{,2} - \beta \lambda_n i (u_{1,1} + u_{2,2}) + \delta \lambda_n i (\tau_{,11} + \tau_{,22}) \right) = f_6. \tag{51}
 \end{aligned}$$

To solve Equations (49)–(51), we employ the ansatz

$$u_1 = A \sin(nx) \sin(ny), \quad u_2 = B \cos(nx) \cos(ny), \quad \tau = C \cos(nx) \sin(ny),$$

where A, B, C will actually depend on n . It should be pointed out that this choice is compatible with the boundary conditions in Equations (44)–(48). Thus, system (49)–(51) is equivalent with finding A, B, C such that

$$\begin{aligned}
 & (-\lambda_n^2 \rho + (\lambda + 3\mu)n^2 + 2b_2 n^4)A - ((\lambda + \mu)n^2 + 2b_4 n^4)B \\
 & \quad + (-\beta \lambda_n n i + 2(m_1 + 2m + 2)n^3)C = \rho, \tag{52}
 \end{aligned}$$

$$\begin{aligned}
 & (-\lambda_n^2 \rho + (\lambda + 3\mu)n^2 + 2b_2 n^4)B - ((\lambda + \mu)n^2 + 2b_4 n^4)A \\
 & \quad + (\beta \lambda_n n i - 2(m_1 + 2m + 2)n^3)C = \rho, \tag{53}
 \end{aligned}$$

$$(2(m_1 + 2m_2)n^3 + \beta \lambda_n n i)(A - B) + (-\lambda_n^2 a + 2n^2 + 2\lambda_n \delta n^2 i)C = a. \tag{54}$$

Let

$$\lambda_n = \sqrt{(8(a_3 + a_4)n^4 + 2\mu n^2 - 1)/\rho}.$$

It is easy to verify that the linear algebraic system (52)–(54) has a unique solution. Summing up Equations (52) and (53), we get

$$(A + B)(-\lambda_n^2 \rho + 2\mu n^2 + 8(a_3 + a_4)n^4) = 2\rho,$$

where $a_3 + a_4 \geq 0$ follows from the positive definiteness of C_{iJKLj} . Equivalently,

$$A + B = 2\rho.$$

and, therefore, by virtue of Young's inequality,

$$\begin{aligned} \|U\|_{\mathcal{H}^*}^2 &\geq \|v_1\|_{L^2(\Omega)}^2 + \|v_2\|_{L^2(\Omega)}^2 = \lambda_n^2 \left(\|u_1\|_{L^2(\Omega)}^2 + \|u_2\|_{L^2(\Omega)}^2 \right) \\ &= \lambda_n^2 \frac{\pi^2}{2} (A^2 + B^2) \geq \frac{\lambda_n^2 \pi^2}{2} \left(\frac{A+B}{2} \right)^2 = \frac{\lambda_n^2 \pi^2 \rho^2}{2} \rightarrow \infty, \end{aligned}$$

whereas we estimate

$$\|F\|_{\mathcal{H}^*}^2 = \frac{3\pi^2}{4} \text{ for any } n \in \mathbb{N},$$

i.e., $(F_n)_n$ remains uniformly bounded w.r.t. $n \in \mathbb{N}$. Hence, the claim follows. \square

4 Linear system with a frictional damping for the elastic part: Exponential stability

In Section 3, if no additional damping for the elastic component is present, Equations (18)–(19) were shown, in general, not to exhibit an exponential decay rate. This justifies the necessity of introducing a damping mechanism for the elastic variable. In the following, we consider a frictional damping which leads to a system of partial differential equations reading as

$$\rho \ddot{u}_i = (A_{iJKj} u_{j,K} - \beta_{Ji} \dot{\tau} - (C_{iJKLIj} u_{j,IL} + M_{iJKL\tau,L})_{,K})_{,J} - E_{ij} \dot{u}_j, \quad (55)$$

$$a \ddot{\tau} = -\beta_{Ki} \dot{u}_{i,K} + m_{IJ} \dot{\tau}_{,IJ} + M_{jLKI} u_{j,LKI} + K_{IJ} \tau_{,IJ} \quad (56)$$

together with the boundary conditions (20) and initial conditions (21). For this system, we assume

$$E_{ij} \xi_i \xi_j \geq \alpha \xi_i \xi_i \text{ and } m_{IJ} \xi_I \xi_J \geq \alpha \xi_I \xi_I \text{ for some } \alpha > 0. \quad (57)$$

The natural first-order energy associated with the mild solution to Equations (55)–(56), (20)–(21) reads as

$$\begin{aligned} \mathcal{E}(t) &= \frac{1}{2} \int_{\Omega} (\rho \dot{u}_i^2 + a \dot{\tau}^2 + A_{iKLj} u_{i,K} u_{j,L} + C_{iJKLj} u_{i,JI} u_{j,LK}) dx \\ &\quad + \frac{1}{2} \int_{\Omega} (K_{IJ} \tau_{,I} \tau_{,J} + 2M_{iJKL} u_{i,KJ} \tau_{,L}) dx, \end{aligned} \quad (58)$$

where, as before, the Einstein's summation convention is applied to terms like $\dot{u}_i^2 = \dot{u}_i \dot{u}_i$, etc. Due to the assumptions in Equation (57), there exists a number $c_{\mathcal{E}} > 0$ such that

$$\mathcal{E}(t) \geq c_{\mathcal{E}} \int_{\Omega} (\dot{u}_i^2 + \dot{\tau}^2 + u_{i,J}^2 + u_{i,JK}^2 + \tau_{,I}^2)(t, x) dx. \quad (59)$$

Theorem 5. *Let $U_0 \in \mathcal{H}$. There exist then positive constants C and c_0 such that*

$$\mathcal{E}(t) \leq C\mathcal{E}(0)e^{-c_0 t} \text{ holds true for every } t \geq 0.$$

Proof. Without loss of generality, let $U_0 \in D(\mathcal{A})$. Indeed, if this is not the case, due to the dense embedding of $D(\mathcal{A}) \hookrightarrow \mathcal{H}$, an appropriate approximating sequence from $D(\mathcal{A})$ can be selected. Further, let $(u, \dot{u}, \tau, \dot{\tau})^T$ denote the classical solution to (55)–(56), (20)–(21) for the initial data U_0 . We construct a Lyapunov functional \mathcal{F} . Computing $\partial_t \mathcal{E}(t)$, we get

$$\begin{aligned} \partial_t \mathcal{E}(t) &= \int_{\Omega} \left(-\frac{1}{\rho} E_{ij} \dot{u}_i \dot{u}_j + \frac{1}{a} m_{IJ} \dot{\tau}_{,IJ} \dot{\tau} \right) dx \\ &= - \int_{\Omega} \left(\frac{1}{\rho} E_{ij} \dot{u}_i \dot{u}_j + \frac{1}{a} m_{IJ} \dot{\tau}_{,I} \dot{\tau}_{,J} \right) dx \leq -\alpha \int_{\Omega} (\dot{u}_i^2 + \dot{\tau}_{,I}^2) dx. \end{aligned}$$

Letting

$$\mathcal{F}_1(t) = \rho \int_{\Omega} \dot{u}_i u_i dx \text{ and } \mathcal{F}_2(t) = a \int_{\Omega} \dot{\tau} \tau dx, \quad (60)$$

after a partial integration, we arrive at

$$\begin{aligned} \partial_t \mathcal{F}_1(t) &= \int_{\Omega} (-A_{iKLj} u_{i,K} u_{j,L} - \beta_{Ji} \dot{\tau}_{,J} u_i - C_{iIJKLj} u_{i,JI} u_{j,LK}) dx \\ &\quad + \int_{\Omega} (-M_{iJKL} \tau_{,L} u_{i,KJ} - E_{ij} \dot{u}_i u_j + \rho \dot{u}^2) dx, \\ \partial_t \mathcal{F}_2(t) &= \int_{\Omega} (\beta_{Ki} \dot{u}_i \tau_{,K} - m_{IJ} \dot{\tau}_{,I} \tau_{,J} - M_{iJKL} \tau_{,L} u_{i,KJ} - K_{IJ} \tau_{,I} \tau_{,J} + a \dot{\tau}^2) dx. \end{aligned}$$

Let $R_0 > \max\{\beta_{Ji}, m_{IJ}, E_{ij}, a, \rho\}$. Utilizing the generalized Young's inequality and the first Poincaré's inequality, we obtain

$$\begin{aligned} \partial_t (\mathcal{F}_1 + \mathcal{F}_2)(t) &= \int_{\Omega} (-A_{iKLj} u_{i,K} u_{j,L} - C_{iIJKLj} u_{i,JI} u_{j,LK} - 2M_{iJKL} \tau_{,L} u_{i,KJ}) dx \\ &\quad + \int_{\Omega} (\beta_{Ki} \dot{u}_i \tau_{,K} - K_{IJ} \tau_{,I} \tau_{,J} - \beta_{Ji} \dot{\tau}_{,J} u_i - m_{IJ} \dot{\tau}_{,I} \tau_{,J} - E_{ij} \dot{u}_i u_j + \rho \dot{u}^2 + a \dot{\tau}^2) dx \\ &\leq -\alpha \int_{\Omega} (u_{i,J}^2 + u_{i,KL}^2 + \tau_{,I}^2) dx + \frac{R_0}{2} \int_{\Omega} \left(\frac{1}{\varepsilon} \dot{u}_i^2 + \varepsilon \tau_{,K}^2 \right) + \left(\frac{1}{\varepsilon} \dot{\tau}_{,J}^2 + \varepsilon u_i^2 \right) dx \\ &\quad + \frac{R_0}{2} \int_{\Omega} \left(\frac{1}{\varepsilon} \dot{\tau}_{,I}^2 + \varepsilon \tau_{,J}^2 \right) + \left(\frac{1}{\varepsilon} \dot{u}_i^2 + \varepsilon u_j^2 \right) + (\dot{u}^2 + \dot{\tau}^2) dx \\ &\leq -\alpha \int_{\Omega} (u_{i,J}^2 + u_{i,KL}^2 + \tau_{,I}^2) dx + \frac{R_0 n}{2} \int_{\Omega} \left(\frac{1}{\varepsilon} \dot{u}^2 + \varepsilon \tau_{,K}^2 + \frac{1}{\varepsilon} \dot{\tau}_{,I}^2 \right) dx \\ &\quad + \frac{R_0 n}{2} \int_{\Omega} \left(\varepsilon c_F u_{i,J}^2 + \frac{1}{\varepsilon} \dot{\tau}_{,I}^2 + \varepsilon \tau_{,J}^2 + \frac{1}{\varepsilon} \dot{u}^2 + \varepsilon c_F u_{j,I}^2 + \dot{u}^2 + c_F \dot{\tau}_{,I}^2 \right) dx. \end{aligned}$$

Now, selecting $\varepsilon > 0$ such that

$$\frac{\alpha}{2} > \frac{R_0 n}{2} \max\{2\varepsilon, 2\varepsilon c_F\},$$

we find

$$\begin{aligned} \partial_t(\mathcal{F}_1 + \mathcal{F}_2)(t) &\leq -\frac{\alpha}{2} \int_{\Omega} (u_{i,J}^2 + u_{i,KL}^2 + \tau_{,I}^2) \, dx \\ &\quad + \frac{R_0 n}{2} \int_{\Omega} \left(\frac{2}{\varepsilon} + 1 \right) \dot{u}^2 + \left(\frac{2}{\varepsilon} + c_F \right) \dot{\tau}_{,I}^2 \, dx. \end{aligned}$$

Next, we define

$$\mathcal{F}(t) = \mathcal{F}_1(t) + \mathcal{F}_2(t) + N\mathcal{E}(t)$$

for some $N > \frac{1}{2} + \frac{R_0 n}{2\alpha} \left(\frac{2}{\varepsilon} + \max\{1, c_F\} \right)$ to be fixed later. Then

$$\partial_t \mathcal{F}(t) \leq -\frac{\alpha}{2} \int_{\Omega} (u_{i,J}^2 + u_{i,KL}^2 + \tau_{,I}^2 + \dot{u}^2 + \dot{\tau}_{,I}^2) \, dx.$$

Using the first Poincaré's inequality and Equation (59), we obtain

$$\partial_t \mathcal{F}(t) \leq -\widehat{C}\mathcal{E}(t).$$

Taking into account

$$|(\mathcal{F}_1 + \mathcal{F}_2)(t)| \leq \widetilde{C}\mathcal{E}(t),$$

we conclude that

$$(N - \widetilde{C})\mathcal{E}(t) \leq \mathcal{F}(t) \leq (N + \widetilde{C})\mathcal{E}(t).$$

If necessary, N is increased to make $N - \widetilde{C}$ positive. Gronwall's inequality now yields

$$\mathcal{E}(t) \leq \frac{1}{N - \widetilde{C}} \mathcal{F}(t) \leq \frac{1}{N - \widetilde{C}} \mathcal{E}(0) e^{-\widehat{C}/(N + \widetilde{C})t} = C\mathcal{E}(0) e^{-c_0 t}.$$

This completes the proof. \square

5 Hyperbolized system with a frictional damping in the elastic part: Global existence and exponential stability

In this last section, we want to study the impact of a nonlinear frictional damping on 'the' equations of thermoelasticity for nonsimple materials. In contrast to Section 4, we replace Equation (19) with a Cattaneo-like hyperbolic relaxation (see, e.g., [22]). Thus, the resulting system becomes purely hyperbolic and we can use the well-known technique due to Lasiecka and Tataru [12] to utilize an observability inequality for the linear system to obtain a uniform decay for the nonlinear one.

Letting $\mathbf{q} = (q_i)$ such that $(m_{IJ}q_{I,J})$ represents the heat flux, for a (small) relaxation parameter $\kappa > 0$, we consider the following semilinear initial-boundary value problem

$$\rho \ddot{u}_i = (A_{iJK} u_{j,K} - \beta_{Ji} \dot{\tau} - (C_{iJKLI} u_{j,IL} + M_{iJKL} \tau_{,L})_{,K})_{,J} - E(|\dot{u}|) \dot{u}_i, \quad (61)$$

$$a \ddot{\tau} = -\beta_{Ki} \dot{u}_{i,K} + m_{IJ} q_{I,J} + M_{jLKI} u_{j,LKI} + K_{IJ} \tau_{,IJ}, \quad (62)$$

$$\kappa \dot{q}_i = \dot{\tau}_{,i} - q_i \quad (63)$$

in $\Omega \times (0, \infty)$ subject to the boundary conditions

$$u_i = 0, \quad u_{i,J} = 0, \quad \tau = 0 \text{ in } \Gamma \times (0, \infty) \quad (64)$$

and the initial conditions

$$\begin{aligned} u_i(\cdot, 0) &= u_i^0, & \dot{u}_i(\cdot, 0) &= \dot{u}_i^0, \\ \tau(\cdot, 0) &= \tau^0, & \dot{\tau}(\cdot, 0) &= \dot{\tau}^0, & q_i(\cdot, 0) &= q_i^0 \text{ in } \Omega. \end{aligned} \quad (65)$$

Note that if $\kappa = 0$ and $E(\cdot)$ is linear, Equations (61)–(63) reduce to (18)–(19).

5.1 Preliminaries

We assume the function $E: [0, \infty) \rightarrow \mathbb{R}$ satisfies the following conditions

- $E(s) > 0$ for $s > 0$,
- $s \mapsto E(s)s$ is continuous and monotonically increasing,
- $\lim_{s \searrow 0} E(s)s = 0$,
- There exist m, M such that

$$0 < m \leq E(s) \leq M \text{ for } s > 1, \quad (66)$$

- The function g_i given by $g_i(v) = E(|v|)v_i$ is uniformly Lipschitz-continuous, i.e.,

$$|g_i(\phi) - g_i(\psi)| \leq L|\phi - \psi| \quad (67)$$

for all $\phi, \psi \in \mathbb{R}^d$.

For Equations (61)–(63), we assume the evolution is taking place on the Hilbert space

$$\begin{aligned} \mathcal{H}_\kappa &= \left\{ U = (u, v, \tau, \theta, q)^T \in (H_0^2(\Omega))^d \times (L^2(\Omega))^d \times H_0^1(\Omega) \times L^2(\Omega) \times (L^2(\Omega))^d \right\} \\ &= \mathcal{H} \times (L^2(\Omega))^d \end{aligned}$$

equipped with the scalar product

$$\langle U, U^* \rangle_{\mathcal{H}_\kappa} = \langle (u, v, \tau, \theta)^T, (u^*, v^*, \tau^*, \theta^*)^T \rangle_{\mathcal{H}} + \int_{\Omega} \kappa m_{ij} q_i^* q_j \, dx$$

for $U = (u, v, \tau, \theta, q), U^* = (u^*, v^*, \tau^*, \theta^*, q^*) \in \mathcal{H}_\kappa$. (See Equation (27).) As before, one can easily prove the inner product is equivalent with the usual product on the Hilbert space \mathcal{H}_κ .

We rewrite the problem (61)–(65) in the abstract form:

$$\dot{U}(t) + \mathcal{K}(U(t)) = 0 \text{ for } t > 0, \quad U(0) = U_0. \quad (68)$$

Here, $\mathcal{K} = -\mathcal{L} + \mathcal{N}$ is a nonlinear operator with the domain $D(\mathcal{K}) := D(\mathcal{L})$, where

$$\mathcal{L}U = \begin{pmatrix} v_i \\ \frac{1}{\rho} (A_{iJKj}u_{j,K} - \beta_{Ji}\theta - (C_{iJKLIj}u_{j,IL} + M_{iJKL}\tau_{,L}),_{K}),_J \\ \theta \\ \frac{1}{a} (-\beta_{Ki}v_{i,K} + m_{IJ}q_{I,J} + M_{jLKI}u_{j,LKI} + K_{IJ}\tau_{,IJ}) \\ \frac{1}{\kappa}\theta_{,i} \end{pmatrix} \quad (69)$$

for $u \in D(\mathcal{L})$ and

$$\mathcal{N}(U) = \left(0, \frac{1}{\rho}E(|v|)v_i, 0, 0, \frac{1}{\kappa}q_i\right)^T \text{ for } u \in D(\mathcal{N}) \quad (70)$$

as well as

$$D(\mathcal{L}) = \left\{ U \in \mathcal{H} \mid v_i \in H_0^1(\Omega), \theta \in H_0^1(\Omega), \right. \quad (71)$$

$$\begin{aligned} & (A_{iJKj}u_{j,K} - \beta_{Ji}\theta \\ & - (C_{iJKLIj}u_{j,IL} + M_{iJKL}\tau_{,L}),_{K}),_J \in L^2(\Omega) \\ & \left. (m_{JI}q_J + M_{jLKI}u_{j,LK} + K_{IJ}\tau_{,J}),_I \in L^2(\Omega) \right\}, \end{aligned}$$

$$D(\mathcal{N}) = \mathcal{H}_\kappa. \quad (72)$$

Definition 6. Let $U_0 \in \mathcal{H}$ and assume \mathcal{L} generates a C_0 -semigroup $(S(t))_{t \geq 0}$ on \mathcal{H}_κ . A function $U \in C^0([0, \infty), \mathcal{H}_\kappa)$ satisfying the integral equation

$$U(t) = S(t)U_0 + \int_0^t S(t-s)\mathcal{N}(U(s))ds$$

is referred to as a mild solution to Equation (68). If U additionally satisfies

$$U \in H_{\text{loc}}^1(0, \infty; \mathcal{H}_\kappa) \cap L_{\text{loc}}^2(0, \infty; D(\mathcal{L})),$$

we call U a strong solution.

By $\mathcal{E}^{\text{nl}}(t)$ we denote the corresponding energy

$$\begin{aligned} \mathcal{E}^{\text{nl}}(t) &= \frac{1}{2} \int_{\Omega} (\rho|\dot{u}|^2 + a\dot{\tau}^2 + A_{iKLj}u_{i,K}u_{j,L} + C_{iJKLj}u_{i,JI}u_{j,LK} + K_{IJ}\tau_{,I}\tau_{,J}) \, dx \\ &+ \frac{1}{2} \int_{\Omega} (2M_{iJKL}u_{i,KJ}\tau_{,L} + \kappa|q|^2) \, dx \end{aligned}$$

associated with a mild solution U to nonlinear Equation (68). We observe

$$\langle \mathcal{L}V, V \rangle_{\mathcal{H}_\kappa} = 0 \text{ for any } V \in D(\mathcal{L}) \text{ and} \quad (73)$$

$$\partial_t \mathcal{E}^{\text{nl}}(t) = - \int_{\Omega} E(|v|)|v|^2 + m_{ij}q_iq_j \, dx \text{ a.e. in } (0, \infty) \quad (74)$$

for any strong solution U to Equation (68).

Remark 7. *By virtue of Stone's theorem (cf. [21], [27, Theorem 3.8.6, p. 105]), Equation (73) is equivalent with \mathcal{A} being skew-adjoint.*

A straightforward adaption of the proof of Lemma 1 yields:

Lemma 8. *The operator \mathcal{L} defined in Equation (69) is the infinitesimal generator of a C_0 -semigroup of contractions on \mathcal{H}_κ .*

For the nonlinear Cauchy problem (68), we have the following existence and uniqueness theorem.

Theorem 9. *For $U_0 \in \mathcal{H}_\kappa$, there exist unique mild solution U to Equation (68). If $U_0 \in D(\mathcal{K})$, the mild solution is strong.*

Proof. By virtue of Equation (67),

$$\begin{aligned} \|\mathcal{N}(U^1) - \mathcal{N}(U^2)\|_{\mathcal{H}_\kappa} &= \left(\rho \int_{\Omega} |E(|v^1|)v^1 - E(|v^2|)v^2|^2 dx \right)^{1/2} \\ &\leq L\sqrt{\rho}\|v^1 - v^2\|_{L^2} \leq L\|U^1 - U^2\|_{\mathcal{H}_\kappa} \end{aligned}$$

for $U^i = (u^i, v^i, \tau^i, \theta^i, q^i)^T$, $i = 1, 2$, i.e., the mapping \mathcal{N} is globally Lipschitzian on \mathcal{H}_κ . Hence, the first claim is a direct consequence of [17, Theorem 1.2, p. 184]. Since \mathcal{H}_κ , being a Hilbert space, is reflexive, the second claim readily follows from [17, Theorem 1.4, p. 189]. \square

By utilizing Lemma 8 and constructing a Lyapunov's functional similar to that one in Section 4 (see also [18]), we further obtain the following linear stability theorem.

Theorem 10. *For*

$$\mathcal{D}: \mathcal{H}_\kappa \rightarrow \mathcal{H}_\kappa, \quad \mathcal{D}U := \left(0, \frac{1}{\rho} E_{ij} v_j, 0, 0, \frac{1}{\kappa} q_i \right)^T \text{ for } U \in \mathcal{H}_\kappa,$$

the unique (mild or classical) solution to the Cauchy problem

$$\dot{U}(t) = \mathcal{L}U(t) - \mathcal{D}U(t) \text{ for } t > 0, \quad U(0) = U_0 \quad (75)$$

is exponentially stable.

5.2 Uniform stability

Recalling Equation (73) and using Theorem 10, we apply Theorem 15 in Appendix A to obtain the following observability result for the linear part of Equation (68).

Theorem 11. *There exist a time period $T_0 > 0$ and a positive constant C such that for any $T \geq T_0$ every mild solution $(u, \dot{u}, \tau, \dot{\tau}, q)^T$ to the Cauchy problem*

$$\dot{U}(t) = \mathcal{L}U(t) \text{ for } t > 0, \quad U(0) = U_0$$

satisfies

$$\mathcal{E}(0) \leq C \int_0^T \int_{\Omega} (|\dot{u}|^2 + |q|^2) dx dt.$$

In the following, we adapt the techniques introduced by Lasiecka and Tataru [12]. The technical difficulties in our case are due to the big system size and mixed-order structure of Equations (61)–(63).

Lemma 12. *For any $T > 0$ and $K > 0$, there exists a positive constant C such that*

$$\mathcal{E}^{\text{nlín}}(T) \leq C \int_0^T \int_{\Omega} (E^2(|\dot{u}|)|\dot{u}|^2 + |\dot{u}|^2 + |q|^2) \, dxdt$$

for any strong solution $\Psi = (u_i, \dot{u}_i, \tau, \dot{\tau}, q)^T$ to Equation (68) additionally satisfying $\mathcal{E}^{\text{nlín}}(0) \leq K$.

Proof. Suppose the converse is true which means there exist $T > 0$ and $K > 0$ together with a sequence of strong solutions $((u^{(m)}, \dot{u}^{(m)}, \tau^{(m)}, \dot{\tau}^{(m)}, q^{(m)}))_m^T$ such that $\mathcal{E}_m^{\text{nlín}}(0) \leq K$ to the Cauchy problem (68), whereas

$$\frac{\mathcal{E}_m^{\text{nlín}}(T)}{\int_0^T \int_{\Omega} E^2(|\dot{u}^{(m)}|)|\dot{u}^{(m)}|^2 + |\dot{u}^{(m)}|^2 + |q^{(m)}|^2 \, dxdt} \rightarrow \infty \text{ as } m \rightarrow \infty.$$

Denote

$$\Pi_m(\cdot) := \int_0^T \int_{\Omega} E^2(|\dot{u}^{(m)}|)|\dot{u}^{(m)}|^2 + |\dot{u}^{(m)}|^2 + |q^{(m)}|^2 \, dxdt.$$

Using Equation (74), we then get

$$\frac{K}{\Pi_m} \geq \frac{\mathcal{E}_m^{\text{nlín}}(0)}{\Pi_m} \geq \frac{\mathcal{E}_m^{\text{nlín}}(T)}{\Pi_m} \rightarrow \infty \text{ as } m \rightarrow \infty,$$

and, thus, $\Pi_m \rightarrow 0$, i.e.,

$$\int_0^T \|\dot{u}^{(m)}\|_{L^2}^2 dt \rightarrow 0 \text{ and } \int_0^T \|q^{(m)}\|_{L^2}^2 dt \rightarrow 0 \text{ as } m \rightarrow \infty.$$

Letting

$$\nu_m = \sqrt{\mathcal{E}_m^{\text{nlín}}(0)}, \quad \bar{u}^{(m)} = \frac{u^{(m)}}{\nu_m}, \quad \bar{\tau}^{(m)} = \frac{\tau^{(m)}}{\nu_m}, \quad \bar{q}^{(m)} = \frac{q^{(m)}}{\nu_m},$$

we observe $(\bar{u}^{(m)}, \bar{\tau}^{(m)}, \bar{q}^{(m)})^T$ is a strong solution to

$$\rho \ddot{u}_i = (A_{iJKj} u_{j,K} - \beta_{Ji} \dot{\tau} - (C_{iJKLIj} u_{j,IL} + M_{iJKL} \tau_{,L})_{,K})_{,J} - E(|\dot{u}| \nu_m) \dot{u}_i \quad (76)$$

$$a \ddot{\tau} = -\beta_{Ki} \dot{u}_{i,K} + m_{IJ} q_{I,J} + M_{jLKI} u_{j,LKI} + K_{IJ} \tau_{,IJ}, \quad (77)$$

$$\kappa \dot{q}_i = \dot{\tau}_{,i} - q_i \quad (78)$$

in $\Omega \times (0, \infty)$ subject to the boundary conditions

$$u_i = 0, \quad u_{i,J} = 0, \quad \tau = 0 \text{ on } \Gamma \times (0, \infty)$$

and the initial conditions

$$u_i = \frac{u_i^0}{\nu_m}, \quad \dot{u}_i = \frac{\dot{u}_i^0}{\nu_m}, \quad \tau = \frac{\tau^0}{\nu_m}, \quad \dot{\tau} = \frac{\dot{\tau}^0}{\nu_m}, \quad q_i = \frac{q_i^0}{\nu_m} \text{ in } \Omega \times \{0\}.$$

By $\bar{\mathcal{E}}_m^{\text{lin}}(t)$ denote the energy of the solution to system (76)–(78). Note that

$$\bar{\mathcal{E}}_m^{\text{lin}}(0) = \frac{\mathcal{E}_m^{\text{nl}}(0)}{\nu_m} = 1. \quad (79)$$

Now,

$$\int_0^T \int_{\Omega} (E^2(|\dot{u}^{(m)}|)|\dot{u}^{(m)}|^2 + |\dot{u}^{(m)}|^2 + |\bar{q}^{(m)}|^2) dx dt = \frac{\Pi_m}{\nu_m^2} = \frac{\Pi_m}{\mathcal{E}_m^{\text{nl}}(0)} \rightarrow 0$$

as $m \rightarrow \infty$. Therefore,

$$\int_0^T \|\dot{u}^{(m)}\|_{L^2}^2 dt \rightarrow 0 \text{ and } \int_0^T \|\bar{q}^{(m)}\|_{L^2}^2 dt \rightarrow 0 \text{ as } m \rightarrow \infty \quad (80)$$

and also

$$\int_0^T \|E(|\dot{u}^{(m)}|)|\dot{u}^{(m)}\|_{L^2}^2 dt \rightarrow 0 \text{ as } m \rightarrow \infty. \quad (81)$$

Now, consider the following linear system

$$\rho \ddot{v}_i = (A_{iJKj} v_{j,K} - \beta_{Ji} \dot{\omega} - (C_{iJKLIj} v_{j,IL} + M_{iJKL} \omega_{,L}),_{K}),_{J}, \quad (82)$$

$$a \ddot{\omega} = -\beta_{Ki} \dot{v}_{i,K} + m_{IJ} r_{I,J} + M_{jLKI} v_{j,LKI} + K_{IJ} \omega_{,IJ}, \quad (83)$$

$$\kappa \dot{r}_i = \dot{\omega}_{,i} - r_i \quad (84)$$

subject to the boundary conditions

$$v_i = 0, \quad v_{i,J} = 0, \quad \omega = 0 \text{ on } \Gamma \times (0, \infty) \quad (85)$$

and the initial conditions

$$v_i = \frac{u_i^0}{\nu_m}, \quad \dot{v}_i = \frac{\dot{u}_i^0}{\nu_m}, \quad \omega = \frac{\tau^0}{\nu_m}, \quad \dot{\omega} = \frac{\dot{\tau}^0}{\nu_m}, \quad r_i = \frac{q_i^0}{\nu_m} \text{ in } \Omega \times \{0\}. \quad (86)$$

Letting $(\bar{v}^{(m)}, \bar{\omega}^{(m)}, \bar{r}^{(m)})^T$ denote the strong solution to Equations (82)–(86) and $\mathcal{E}_m^{\text{lin}}(t)$ be the corresponding energy, the initial conditions (86) imply

$$\mathcal{E}_m^{\text{lin}}(0) = \bar{\mathcal{E}}_m^{\text{lin}}(0).$$

Exploring Equations (76)–(78) and (82)–(86), we deduce that $w_i^{(m)} = \bar{u}_i^{(m)} - \bar{v}_i^{(m)}$, $\chi^{(m)} = \bar{\tau}^{(m)} - \bar{\omega}^{(m)}$, $p_i^{(m)} = \bar{q}_i^{(m)} - \bar{r}_i^{(m)}$ strongly solves the ‘incremental’ system

$$\rho \ddot{w}_i = (A_{iJKj} w_{j,K} - \beta_{Ji} \dot{\chi} - (C_{iJKLIj} w_{j,IL} + M_{iJKL} \chi_{,L}),_{K}),_{J} \quad (87)$$

$$- E(|\dot{u}^{(m)}| \nu_m) \dot{w}_i^{(m)}$$

$$a \ddot{\chi} = -\beta_{Ki} \dot{w}_{i,K} + m_{IJ} p_{I,J} + M_{jLKI} w_{j,LKI} + K_{IJ} \chi_{,IJ}, \quad (88)$$

$$\kappa \dot{p}_i = \dot{\chi}_{,i} - p_i, \quad (89)$$

together with the boundary conditions

$$w_i = 0, \quad w_{i,J} = 0, \quad \chi = 0 \text{ on } \Gamma \times (0, \infty) \quad (90)$$

and the initial conditions

$$w_i(\cdot, 0) = \dot{w}_i(\cdot, 0) = 0, \quad \chi(\cdot, 0) = \dot{\chi}(\cdot, 0) = 0, \quad p_i(\cdot, 0) = 0 \text{ in } \Omega. \quad (91)$$

With $\mathcal{E}_m^{\text{diff}}(t)$ denoting the associated energy, Equation (91) implies $\mathcal{E}_m^{\text{diff}}(0) = 0$, for which we easily verify

$$\partial_t \mathcal{E}_m^{\text{diff}}(t) = \int_{\Omega} \left(-E(|\dot{u}^{(m)}|) \dot{u}_i^{(m)} \dot{w}_i^{(m)} - m_{ij} \bar{q}_j^{(m)} p_i \right) dx.$$

Therefore,

$$\begin{aligned} \mathcal{E}_m^{\text{diff}}(t) &= \mathcal{E}_m^{\text{diff}}(t) - \mathcal{E}_m^{\text{diff}}(0) \\ &= - \int_0^t \int_{\Omega} \left(E(|\dot{u}^{(m)}|) \dot{u}_i^{(m)} \dot{w}_i^{(m)} + m_{IJ} \bar{q}_j^{(m)} p_i \right) dx dt. \end{aligned} \quad (92)$$

Note that $\int_0^T \|\dot{w}^{(m)}\|_{L^2}^2 dt$ and $\int_0^T \|p^{(m)}\|_{L^2}^2 dt$ are bounded w.r.t. $m \in \mathbb{N}$. Indeed, a straightforward computation yields

$$\partial_t \mathcal{E}_m^{\text{lin}}(t) = 0.$$

Thus, $\mathcal{E}_m^{\text{lin}}(t) = \mathcal{E}_m^{\text{lin}}(0)$ and, therefore,

$$\frac{\rho}{2} \int_{\Omega} |\dot{v}^{(m)}|^2 dx + \frac{\kappa\alpha}{2} \int_{\Omega} |\bar{r}^{(m)}|^2 dx \leq \mathcal{E}_m^{\text{lin}}(t) = \mathcal{E}_m^{\text{lin}}(0) = \bar{\mathcal{E}}_m^{\text{lin}}(0) = 1,$$

whence

$$\int_0^T \int_{\Omega} \left(\frac{\rho}{2} |\dot{v}^{(m)}|^2 + \frac{\kappa\alpha}{2} |\bar{r}^{(m)}|^2 \right) dx dt \leq T.$$

Estimating

$$\begin{aligned} \left(\int_0^T \|\dot{w}_i^{(m)}\|_{L^2}^2 dt \right)^{1/2} &\leq \left(\int_0^T \|\dot{u}_i^{(m)}\|_{L^2}^2 dt \right)^{1/2} + \left(\int_0^T \|\dot{v}_i^{(m)}\|_{L^2}^2 dt \right)^{1/2} \text{ and} \\ \left(\int_0^T \|p_i^{(m)}\|_{L^2}^2 dt \right)^{1/2} &\leq \left(\int_0^T \|\bar{q}_i^{(m)}\|_{L^2}^2 dt \right)^{1/2} + \left(\int_0^T \|\bar{r}_i^{(m)}\|_{L^2}^2 dt \right)^{1/2} \end{aligned}$$

and recalling Equation (80), we arrive at the boundness of $\int_0^T \|\dot{w}^{(m)}\|_{L^2}^2 dt$ and $\int_0^T \|p^{(m)}\|_{L^2}^2 dt$ w.r.t. $m \in \mathbb{N}$. Using this fact and applying Cauchy & Schwarz' inequality to Equation (92), we further find a positive constant C such that

$$\begin{aligned} \mathcal{E}_m^{\text{diff}}(t) &\leq \left(\int_0^t \int_{\Omega} E^2(|\dot{u}^{(m)}|) |\dot{u}_i^{(m)}|^2 dx dt \right)^{1/2} \left(\int_0^t \int_{\Omega} |\dot{w}_i^{(m)}|^2 dx dt \right)^{1/2} \\ &\quad + \max_{i,j} |m_{ij}| \left(\int_0^t \int_{\Omega} |\bar{q}_j^{(m)}|^2 dx dt \right)^{1/2} \left(\int_0^t \int_{\Omega} |p_i|^2 dx dt \right)^{1/2} \\ &\leq C \left(\int_0^T \int_{\Omega} E^2(|\dot{u}^{(m)}|) |\dot{u}_i^{(m)}|^2 dx dt \right)^{1/2} + C \left(\int_0^T \int_{\Omega} |\bar{q}_j^{(m)}|^2 dx dt \right)^{1/2} \\ &\leq C \left(\left(\int_0^T \|E(|\dot{u}^{(m)}|) \dot{u}^{(m)}\|_{L^2}^2 dt \right)^{1/2} + \left(\int_0^T \|\bar{q}^{(m)}\|_{L^2}^2 dt \right)^{1/2} \right). \end{aligned}$$

Hence,

$$\max_{t \in [0, T]} \mathcal{E}_m^{\text{diff}}(t) \leq C \left(\left(\int_0^T \|E(|\dot{u}^{(m)}|)|\dot{u}^{(m)}\|_{L^2}^2 dt \right)^{1/2} + \left(\int_0^T \|\bar{q}^{(m)}\|_{L^2}^2 dt \right)^{1/2} \right). \quad (93)$$

Further, using the trivial inequality $a^2 = (a - b + b)^2 \leq 2((a - b)^2 + b^2)$, we get

$$\begin{aligned} \int_{\Omega} (|\dot{v}^{(m)}|^2 + |\bar{r}^{(m)}|^2) dx &\leq 2 \int_{\Omega} (|\dot{u}^{(m)} - \dot{v}^{(m)}|^2 + |\bar{q}^{(m)} - \bar{r}^{(m)}|^2 + |\dot{u}^{(m)}|^2 + |\bar{q}^{(m)}|^2) dx \\ &= 2 \int_{\Omega} (|\dot{w}^{(m)}|^2 + |p^{(m)}|^2 + |\dot{u}^{(m)}|^2 + |\bar{q}^{(m)}|^2) dx \\ &\leq C_1 \mathcal{E}_m^{\text{diff}}(t) + 2 \int_{\Omega} (|\dot{u}^{(m)}|^2 + |\bar{q}^{(m)}|^2) dx \\ &\leq C_1 \max_{t \in [0, T]} \mathcal{E}_m^{\text{diff}}(t) + 2 \int_{\Omega} (|\dot{u}^{(m)}|^2 + |\bar{q}^{(m)}|^2) dx. \end{aligned}$$

Integrating the latter inequality w.r.t. t and using Equations (80), (81) and (93), we obtain

$$\int_0^T \int_{\Omega} (|\dot{v}^{(m)}|^2 + |\bar{r}^{(m)}|^2) dx dt \rightarrow 0 \text{ as } m \rightarrow \infty.$$

On the other hand, recalling Equations (79), (87) and Theorem 11, we find

$$1 = \bar{\mathcal{E}}_m^{\text{lin}}(0) = \mathcal{E}_m^{\text{lin}}(0) \leq c \int_{\Omega} (|\dot{v}^{(m)}|^2 + |\bar{r}^{(m)}|^2) dx dt \rightarrow 0 \text{ as } m \rightarrow \infty,$$

which is a contradiction to our original assumption. \square

To proceed further, let us introduce some notations. Thanks to our assumptions on the function E , according to [12], there exists a real-valued function $h: [0, \infty) \rightarrow [0, \infty)$, being concave, strictly increasing and satisfying $h(0) = 0$ and

$$h(s^2 E(s)) \geq s^2 + E^2(s) s^2 \text{ for } s \in [0, 1]. \quad (94)$$

Further, we define a function r by means of

$$r(s) = h\left(\frac{s}{|Q|}\right) \text{ for } s \geq 0, \quad (95)$$

where $Q = \Omega \times (0, T)$ and $|Q|$ is the standard Borel measure of Q . Further, let

$$p(s) = (cI + r)^{-1}(Ms) \text{ for } s \geq 0, \quad (96)$$

where M and c are some positive constants to be defined later. Finally, let

$$q(s) = s - (I + p)^{-1}(s) \text{ for } s \geq 0. \quad (97)$$

We quote the following lemma due to Lasiecka and Tataru [12] we use below.

Lemma 13. *Let the functions p, q be defined as above. For any number sequence $(s_n)_n \subset (0, \infty)$ with $s_{m+1} + p(s_{m+1}) \leq s_m$, we have*

$$s_m \leq S(m) \text{ for every } m \in \mathbb{N},$$

where $S(t)$ is a solution of the scalar Cauchy problem

$$\dot{S}(t) + q(S(t)) = 0 \text{ for } t > 0, \quad S(0) = s_0. \quad (98)$$

Moreover, if p satisfies $p(s) > 0$ for $s > 0$, then

$$\lim_{t \rightarrow \infty} S(t) = 0.$$

Now, we are in position to prove the uniform stability result for the Cauchy problem (68).

Theorem 14. *For a sufficiently large number $T > 0$, we have*

$$\mathcal{E}^{\text{nl}}(t) \leq S\left(\frac{t}{T} - 1\right) \text{ for } t > T.$$

Moreover, $S(t) \rightarrow 0$ as $t \rightarrow \infty$, where $S(t)$ solves the ODE (98).

Proof. Consider the sets

$$\begin{aligned} Q_1 &= \{(x, t) \in \Omega \times (0, T) \mid |\dot{u}(x, t)| > 1\}, \\ Q_2 &= \{(x, t) \in \Omega \times (0, T) \mid |\dot{u}(x, t)| \leq 1\}. \end{aligned}$$

Due to the additivity property of Lebesgue integral, we have

$$\begin{aligned} \int_0^T \int_{\Omega} (E^2(|\dot{u}|)|\dot{u}|^2 + |\dot{u}|^2) dx dt &= \int_{Q_1} (E^2(|\dot{u}|)|\dot{u}|^2 + |\dot{u}|^2) dx dt \\ &\quad + \int_{Q_2} (E^2(|\dot{u}|)|\dot{u}|^2 + |\dot{u}|^2) dx dt. \end{aligned}$$

On the set with $|\dot{u}| > 1$, Equation (66) implies

$$E^2(|\dot{u}|)|\dot{u}|^2 \leq ME(|\dot{u}|)|\dot{u}|^2 \text{ and } |\dot{u}|^2 \leq \frac{1}{m}E(|\dot{u}|)|\dot{u}|^2$$

with m, M from Equation (66). Therefore,

$$\begin{aligned} \int_{Q_1} (E^2(|\dot{u}|)|\dot{u}|^2 + |\dot{u}|^2) dx dt &\leq \left(M + \frac{1}{m}\right) \int_{Q_1} E(|\dot{u}|)|\dot{u}|^2 dx dt \\ &\leq c_1 \int_Q E(|\dot{u}|)|\dot{u}|^2 dx dt, \end{aligned} \quad (99)$$

where $c_1 = M + m^{-1}$.

Next, consider the integral over Q_2 . Using Equation (94), we estimate

$$\int_{Q_2} E^2(|\dot{u}|)|\dot{u}|^2 + |\dot{u}|^2 dxdt \leq \int_{Q_2} h(E(|\dot{u}|)|\dot{u}|^2) dxdt.$$

Now, Jensen's inequality yields

$$\int_{Q_2} h(E(|\dot{u}|)|\dot{u}|^2) dxdt \leq |Q|h\left(\int_{Q_2} \frac{E(|\dot{u}|)|\dot{u}|^2}{|Q|} dxdt\right),$$

and, recalling the definition of function r in Equation (95), we get

$$\begin{aligned} \int_{Q_2} (E^2(|\dot{u}|)|\dot{u}|^2 + |\dot{u}|^2) dxdt &\leq |Q|r\left(\int_{Q_2} E(|\dot{u}|)|\dot{u}|^2 dxdt\right) \\ &\leq |Q|r\left(\int_Q E(|\dot{u}|)|\dot{u}|^2 dxdt\right). \end{aligned} \quad (100)$$

Combining Equations (99) and (100), we obtain

$$\int_0^T \int_{\Omega} (E^2(|\dot{u}|)|\dot{u}|^2 + |\dot{u}|^2) dxdt \leq (c_1\mathbf{I} + |Q|r)\left(\int_Q E(|\dot{u}|)|\dot{u}|^2 dxdt\right),$$

where \mathbf{I} is the identity function, i.e.,

$$(c_1\mathbf{I} + |Q|r)(s) = c_1s + |Q|r(s) \text{ for } s \geq 0.$$

Further, exploiting Lemma 12 and the monotonicity of r , we get

$$\begin{aligned} \mathcal{E}^{\text{nlín}}(T) &\leq C\left((c_1\mathbf{I} + |Q|r)\left(\int_Q E(|\dot{u}|)|\dot{u}|^2 dxdt\right) + \int_Q |q|^2 dxdt\right) \\ &\leq C\left(c_2 \int_Q (E(|\dot{u}|)|\dot{u}|^2 + |q|^2) dxdt + |Q|r\left(\int_Q E(|\dot{u}|)|\dot{u}|^2 dxdt\right)\right) \\ &\leq C\left(c_2 \int_Q (E(|\dot{u}|)|\dot{u}|^2 + |q|^2) dxdt + |Q|r\left(\int_Q (E(|\dot{u}|)|\dot{u}|^2 + |q|^2) dxdt\right)\right) \\ &= C\left((c_2\mathbf{I} + |Q|r)\left(\int_Q (E(|\dot{u}|)|\dot{u}|^2 + |q|^2) dxdt\right)\right), \end{aligned}$$

where $c_2 = \max\{c_1, 1\}$. Thus, we obtain

$$\frac{1}{C|Q|}\mathcal{E}^{\text{nlín}}(T) \leq \left(\frac{c_2}{|Q|}\mathbf{I} + r\right)\left(\int_Q (E(|\dot{u}|)|\dot{u}|^2 + |q|^2) dxdt\right),$$

or, equivalently,

$$M\mathcal{E}^{\text{nlín}}(T) \leq (c\mathbf{I} + r)(\mathcal{E}^{\text{nlín}}(0) - \mathcal{E}^{\text{nlín}}(T)).$$

Now, recalling Equation (96), we get

$$p(\mathcal{E}^{\text{nlín}}(T)) \leq \mathcal{E}^{\text{nlín}}(0) - \mathcal{E}^{\text{nlín}}(T),$$

or, equivalently,

$$p(\mathcal{E}^{\text{nl}}(T)) + \mathcal{E}^{\text{nl}}(T) \leq \mathcal{E}^{\text{nl}}(0).$$

It can easily be seen that, replacing $0, T$ with $mT, (m+1)T$ above, we obtain

$$p(\mathcal{E}^{\text{nl}}((m+1)T)) + \mathcal{E}^{\text{nl}}((m+1)T) \leq \mathcal{E}^{\text{nl}}(mT),$$

Note that the constants $C, c, c_1, c_2, |Q|$ and the functions h, r, p, q remain the same. As before, applying Lemma 13 with $s_m = \mathcal{E}^{\text{nl}}(mT)$, we conclude

$$E(mT) \leq S(m) \text{ for } m \in \mathbb{N} \text{ and } \lim_{t \rightarrow \infty} S(t) = 0.$$

It can further be easily shown that $q(s) \geq 0$ for $s \geq 0$ and, thus, $S(\cdot)$ is monotonically decreasing. Therefore, for $t = mT + \tau$ with $\tau \in [0, T]$, we have

$$E(t) \leq E(mT) \leq S(m) = S\left(\frac{t-\tau}{T}\right) \leq S\left(\frac{t}{T} - 1\right),$$

which finishes the proof. □

Funding

This work has been partially funded by the Young Scholar Fund at the University of Konstanz, Germany. MP has been supported by the ERC-CZ Project LL1202 ‘MOdelling REvisited + MOdel REduction’ at Charles University in Prague, Czech Republic and the Deutsche Forschungsgemeinschaft (DFG) through CRC 1173 at Karlsruhe Institute of Technology, Germany.

References

- [1] Aouadi M, Moulahi T. Approximate controllability of abstract nonsimple thermoelastic problem. *Evol. Equat. Contr. Theor.* 2015; 4(4):373–389.
- [2] Ciarletta M, Ieşan D. Non-Classical Elastic Solids. In: Pitman Research Notes in Mathematical Sciences, Vol. 293, Longman Scientific & Technical, Harlow, Essex, 1993.
- [3] Fernández Sare HD, Muñoz Rivera JE, Quintanilla R. Decay of solutions in non-simple thermoelastic bars. *Int. J. Eng. Sci.* 2010; 48:1233–1241.
- [4] Gawinecki JA, Łazuka J. Global solution on Cauchy problem in nonlinear non-simple thermoelastic materials. *Proc. Appl. Mat. Mech.* 2006; 6:371–372.
- [5] Green AE, Naghdi PM. A unified procedure for construction of theories of deformable media. I. Classical continuum physics, *Proc. Math. Phys. Eng. Sci.* 1995; 448.1934:335–356.

- [6] Green AE, Naghdi PM. A unified procedure for construction of theories of deformable media. II. Generalized continua, Proc. Math. Phys. Eng. Sci. 1995; 448.1934:357–377.
- [7] Green AE, Naghdi PM. A unified procedure for construction of theories of deformable media. III. Mixtures of interacting continua, Proc. Math. Phys. Eng. Sci. 1995; 448.1934:379–388.
- [8] Green AE, Rivlin RS. Multipolar continuum mechanics. Arch. Ration. Mech. An. 1964; 17:113–147.
- [9] Haraux A. Une remarque sur la stabilisation de certains systèmes du deuxième ordre en temps. Portugal. Math. 1989; 46:245–258.
- [10] Ieşan D. Thermoelastic Models of Continua, Solid Mechanics and Its Applications 118, Springer Netherlands; 2004.
- [11] Kadets V. Kurs funkcionalnogo analiza (A course on functional analysis). Kharkiv: University of Kharkiv; 2006. ISBN 966-623-199-9 (In Ukrainian)
- [12] Lasiecka I, Tataru D. Uniform boundary stabilization of semilinear wave equation with nonlinear boundary damping. Differ. Integral Equ.; 1993; 6:507–533.
- [13] Liu Z, Zheng S. Semigroups Associated with Dissipative Systems. Chapman and Hall/CRC: Boca Raton, London, New York, Washington, D.C.; 1999.
- [14] Magaña A, Quintanilla R. Exponential decay in nonsimple thermoelasticity of type III. Math. Method. Appl. Sci. 2014; 39:225–235.
- [15] Mindlin RD. Micro-structure in linear elasticity. Arch. Ration. Mech. An. 1964; 16:51–78.
- [16] Pata V, Quintanilla R. On the decay of solutions in nonsimple elastic solids with memory. J. Math. Anal. Appl. 2010; 363:19–28.
- [17] Pazy A. Semigroups of Linear Operators and Applications to Partial Differential Equations. Springer: New York; 1983.
- [18] Pokojovy M. On stability of hyperbolic thermoelastic Reissner-Mindlin-Timoshenko plates. Math. Meth. Appl. Sci. 2015; 38:1225–1246.
- [19] Quintanilla R. Thermoelasticity without dissipation of nonsimple materials. Z. Angew. Math. Mech. 2003; 83:172–180.
- [20] Rajagopal KR. On implicit constitutive theories. Appl. Math. 2003, 48:279–319.
- [21] Stone MH, On one-parameter unitary groups in Hilbert Space, Ann. Math. 1932; 33(3):643–648.
- [22] Tamma KK, Zhou X. Macroscale and microscale thermal transport and thermo-mechanical interactions: some noteworthy perspectives, J. Therm. Stresses 1998; 21(3–4):405–449.

- [23] Tebou L. Equivalence between observability and stabilization for a class of second order semilinear evolution equations. *Discret. Contin. Dyn. Syst.* 2009; 1:744–752.
- [24] Toupin RA. Theorie of elasticity with couple-stress. *Arch. Ration. Mech. An.* 1964; 17:85–112.
- [25] Truesdell C, Noll W. *The Non-Linear Field Theories of Mechanics*, 3rd ed., Antman SS (editor), Springer-Verlag Berlin Heidelberg New York; 2004.
- [26] Truesdell C, Toupin RA. The classical field theories. In: *Handbuch der Physik*, Vol. III/1, Flügge S (ed). Springer-Verlag: Berlin, 1960.
- [27] Tucsnak M, Weiss G. *Observation and Control for Operator Semigroups*. Springer Science & Business Media. Birkhäuser: Basel, Boston, Berlin; 2009.

A Relation between observability and stabilization

Consider a pair of evolution equations

$$\dot{U}(t) + \mathcal{A}U(t) + \mathcal{B}U(t) = 0 \text{ for } t \geq 0, \quad U(0) = U_0, \quad (101)$$

$$\dot{\phi}(t) + \mathcal{A}\phi(t) = 0 \text{ for } t \geq 0, \quad \phi(0) = \phi_0, \quad (102)$$

taking place on a Hilbert space \mathcal{H} endowed with inner product $\langle \cdot, \cdot \rangle$. By $E_U(t) := \frac{1}{2}\langle U, U \rangle_{\mathcal{H}}$ and $E_{\phi}(t) = \frac{1}{2}\langle \phi, \phi \rangle_{\mathcal{H}}$ we denote the corresponding natural energies. Further, let $\mathcal{B}: \mathcal{H} \rightarrow \mathcal{H}$ be a bounded linear operator. We are interested in establishing a relation between the following two properties:

- i)* There exist numbers $T > 0$ and $C > 0$ such that every mild solution of Equation (102) satisfies

$$E_{\phi}(0) \leq C \int_0^T \langle \mathcal{B}\phi, \phi \rangle dt. \quad (103)$$

- ii)* There exist numbers $M > 0$ and $\lambda > 0$ such that every mild solution of Equation (101) satisfies

$$E_U(t) \leq M e^{-\lambda t} E_U(0) \text{ for } t \geq 0. \quad (104)$$

Condition *i)* is referred to the (exact) observability inequality at time T for the operator pair $(\mathcal{A}, \mathcal{B})$, whereas *ii)* represents the exponential stability of the semigroup generated by $\mathcal{A} + \mathcal{B}$. Using the approach proposed by Haraux [9] and further developed Tebou [23] for a class of abstract wave-equation-like problems, we prove the following theorem.

Theorem 15. *Assume \mathcal{A} is a infinitesimal generator of a C_0 -semigroup of contractions on \mathcal{H} satisfying*

$$\langle \mathcal{A}U, U \rangle = 0 \text{ for any } U \in D(\mathcal{A}), \quad (105)$$

and \mathcal{B} is a self-adjoint, positive semidefinite operator. Then there exists a number $T_0 > 0$ such that the condition (104) implies (103) for any $T \geq T_0$.

Proof. Let ϕ be an arbitrary strong solution of (102). By U we denote a strong solution of the problem (101) with the initial condition

$$U(0) = \phi(0).$$

Multiplying Equation (101) by U in \mathcal{H} and using Equation (105), we get

$$\begin{aligned} \langle \dot{U}, U \rangle + \langle \mathcal{A}U, U \rangle &= -\langle \mathcal{B}U, U \rangle, \\ \partial_t E_U(t) &= -\langle \mathcal{B}U, U \rangle, \\ E_U(0) - E_U(T) &= \int_0^T \langle \mathcal{B}U, U \rangle dt. \end{aligned}$$

It follows from Equation (104) that there exists $T_0 > 0$ such that $E_U(T) \leq \frac{1}{2}E_U(0)$ for $T \geq T_0$. Hence,

$$\int_0^T \langle \mathcal{B}U, U \rangle dt \geq \frac{1}{2}E_U(0).$$

Letting $\psi = \phi - U$, we observe ψ is a strong solution of

$$\dot{\psi}(t) + \mathcal{A}\psi(t) + \mathcal{B}\psi(t) = \mathcal{B}\phi(t) \text{ for } t > 0, \quad \psi(0) = 0.$$

Denoting by $E_\psi(t)$ the associated energy, we get

$$\begin{aligned} \langle \dot{\psi}, \psi \rangle + \langle \mathcal{A}\psi, \psi \rangle + \langle \mathcal{B}\psi, \psi \rangle &= \langle \mathcal{B}\psi, \phi \rangle, \\ \langle \dot{\psi}, \psi \rangle + \langle \mathcal{B}\psi, \psi \rangle &= \langle \mathcal{B}\psi, \phi \rangle. \end{aligned}$$

Using Cauchy & Schwarz' inequality for bilinear forms (cf, e.g., [11, p. 358]), we get the following cascade of inequalities

$$\begin{aligned} \langle \dot{\psi}, \psi \rangle + \langle \mathcal{B}\psi, \psi \rangle &= \langle \mathcal{B}\psi, \phi \rangle \leq \frac{1}{2} (\langle \mathcal{B}\psi, \psi \rangle + \langle \mathcal{B}\phi, \phi \rangle), \\ \langle \dot{\psi}, \psi \rangle + \frac{1}{2} \langle \mathcal{B}\psi, \psi \rangle &\leq \frac{1}{2} \langle \mathcal{B}\phi, \phi \rangle, \\ \int_0^T \langle \dot{\psi}, \psi \rangle dt + \frac{1}{2} \int_0^T \langle \mathcal{B}\psi, \psi \rangle dt &\leq \frac{1}{2} \int_0^T \langle \mathcal{B}\phi, \phi \rangle dt, \\ E_\psi(T) - E_\psi(0) + \frac{1}{2} \int_0^T \langle \mathcal{B}\psi, \psi \rangle dt &\leq \frac{1}{2} \int_0^T \langle \mathcal{B}\phi, \phi \rangle dt. \end{aligned}$$

Now, using the fact $E_\psi(0) = 0$ and $E_\psi(t) \geq 0$, we get

$$\int_0^T \langle \mathcal{B}\psi, \psi \rangle dt \leq \int_0^T \langle \mathcal{B}\phi, \phi \rangle dt. \quad (106)$$

Tacking into account the fact $E_U(0) = E_\phi(0)$, Equation (106) and Cauchy & Schwarz' inequality for bilinear forms, we obtain

$$\frac{1}{2}E_\phi(0) = \frac{1}{2}E_U(0) \leq \int_0^T \langle \mathcal{B}U, U \rangle dt = \int_0^T \langle \mathcal{B}(\phi - \psi), (\phi - \psi) \rangle dt$$

$$\begin{aligned} &= \int_0^T \langle \mathcal{B}\phi, \phi \rangle dt + \int_0^T \langle \mathcal{B}\psi, \psi \rangle dt - 2 \int_0^T \langle \mathcal{B}\phi, \psi \rangle dt \\ &\leq 2 \left(\int_0^T \langle \mathcal{B}\phi, \phi \rangle dt + \int_0^T \langle \mathcal{B}\psi, \psi \rangle dt \right) \leq 8 \int_0^T \langle \mathcal{B}\phi, \phi \rangle dt, \end{aligned}$$

which finishes the proof. □