

Entropy decay for the Kac evolution

Federico Bonetto, Alissa Geisinger,
Michael Loss, Tobias Ried

CRC Preprint 2017/20, July 2017

KARLSRUHE INSTITUTE OF TECHNOLOGY

CRC 1173



Wave
phenomena

Participating universities



Universität Stuttgart

EBERHARD KARLS
UNIVERSITÄT
TÜBINGEN



Funded by

DFG

ISSN 2365-662X

ENTROPY DECAY FOR THE KAC EVOLUTION

FEDERICO BONETTO

School of Mathematics, Georgia Institute of Technology
Atlanta, GA 30332, United States of America

ALISSA GEISINGER

Universität Tübingen, Fachbereich Mathematik
Auf der Morgenstelle 10, 72076 Tübingen, Germany

MICHAEL LOSS

School of Mathematics, Georgia Institute of Technology
Atlanta, GA 30332, United States of America

TOBIAS RIED

Institute for Analysis, Karlsruhe Institute of Technology (KIT)
Englerstraße 2, 76131 Karlsruhe, Germany

ABSTRACT. We consider solutions to the Kac master equation for initial conditions where N particles are in a thermal equilibrium and $M \leq N$ particles are out of equilibrium. We show that such solutions have exponential decay in entropy relative to the thermal state. More precisely, the decay is exponential in time with an explicit rate that is essentially independent on the particle number. This is in marked contrast to previous results which show that the entropy production for arbitrary initial conditions is inversely proportional to the particle number. The proof relies on Nelson's hypercontractive estimate and the geometric form of the Brascamp-Lieb inequalities due to Franck Barthe. Similar results hold for the Kac-Boltzmann equation with uniform scattering cross sections.

1. Introduction

Among the models describing a gas of interacting particles, the Kac master equation [21], due to its simplicity, occupies a special place. It is useful in illuminating various issues in kinetic theory, e.g., providing a reasonably satisfactory derivation of the spatially homogeneous Boltzmann equation and giving a mathematical framework for investigating the approach to equilibrium. These issues were, in fact, the motivation for Kac's original work [21]. Although it does not have a foundation in Hamiltonian mechanics, the Kac master equation is based on simple probabilistic principles and yields a linear evolution equation for the velocity distribution for N particles undergoing collisions. It is in this context that Kac invented the notion of propagation of chaos and he used this notion to derive the spatially homogeneous, non-linear Kac-Boltzmann equation. The approach through master equations led Kac to formulate the notion of approach to equilibrium and suggested various avenues to investigate this problem as the number of particles, N , becomes large. He emphasized that this could be done in a quantitative way if one could show, e.g., that the gap of the generator is bounded below

2010 *Mathematics Subject Classification.* Primary: 82C22; Secondary: 60J25.

Key words and phrases. Kac model, Entropy decay, Thermostat, Correlation Inequalities, Maxwellian molecules.

© 2017 by the authors. Reproduction of this article by any means permitted for non-commercial purposes.

uniformly in N . This, known as Kac's conjecture [21], was proved by Élise Janvresse in [20] and, as a further sign of the simplicity of the model, the gap was computed explicitly in [9, 10], see also [24]. One of the problems in using the gap is that the approach to equilibrium is measured in terms of an L^2 distance. While this does seem to be a natural way to look at this problem, the size of the L^2 norm of approximately independent probability distributions increases exponentially with the size of the system. Thus, the half life of the L^2 norm is of order N .

A natural measure is, of course, given by the entropy, which is extensive, i.e., proportional to N . There has not been much success in proving exponential decay of the entropy with good rates. In [29] Cedric Villani showed that the entropy decays exponentially, albeit with a rate that is bounded below by a quantity that is inversely proportional to N . This estimate was complemented by Amit Einav [14], who gave an example of a state that has entropy production essentially of order $1/N$. His example is the initial state in which most of the energy is concentrated in a few particles while most of the others have very little energy. One might surmise, based on physical intuition, that this state is physically very improbable and still has low entropy production because most of the particles are in some sort of equilibrium. This intuition can be made rigorous, see [14], although by a quite difficult computation. One should add that low entropy production does not preclude exponential decay in entropy, i.e., large entropy production for the initial state might not be necessary for an exponential decay rate for the entropy.

A breakthrough was achieved by Mischler and Mouhot in [26, 25]. They undertook a general investigation of the Kac program for gases of hard spheres and true Maxwellian molecules in three dimensions. Among the results of Mischler and Mouhot is a proof that these systems relax towards equilibrium in relative entropy as well as in Wasserstein distance with a rate that is independent of the particle number. As expected, they achieve this not for any initial condition, but rather for a natural class of chaotic states. The rate of relaxation is, however, polynomial in time.

To summarize, there is so far no mathematical evidence that the entropy in the Kac model in general decays exponentially with a rate that is independent of N and physical intuition suggests that for highly "improbable" states, such as the one used by Einav, this cannot be expected. One can restrict the class of initial conditions by considering chaotic states as done by Mischler and Mouhot, which shifts the problem of finding suitable initial conditions for proving exponential decay to the level of the non-linear Boltzmann equation.

In this paper we take a different approach, one which is based on the idea of coupling a system of particles to a reservoir. Recall from [7] the master equation of M particles with velocities $\mathbf{v} = (v_1, v_2, \dots, v_M)$ interacting with a thermostat at temperature $1/\beta$,

$$\frac{\partial f}{\partial t} = \mathcal{L}_T f, \quad f(\mathbf{v}, 0) = f_0(\mathbf{v}). \quad (1)$$

The operator \mathcal{L}_T is given by

$$\mathcal{L}_T f = \mu \sum_{j=1}^M (B_j - I) f,$$

where

$$B_j[f](\mathbf{v}) := \int_{\mathbb{R}} dw \frac{1}{2\pi} \int_{-\pi}^{\pi} d\theta \sqrt{\frac{\beta}{2\pi}} e^{-\beta w_j^*(\theta)^2/2} f(\mathbf{v}_j(\theta, w)) ,$$

$$\mathbf{v}_j(\theta, w) = (v_1, \dots, v_j \cos(\theta) + w \sin(\theta), \dots, v_M) \quad \text{and} \quad w_j^*(\theta) = -v_j \sin(\theta) + w \cos(\theta) .$$

Thus, $B_j[f](\mathbf{v})$ describes the effect of a collision between particle j in the system and a particle in the reservoir. After the collision, the particle from the thermostat is discarded, which ensures that the thermostat stays in equilibrium. The interaction times with the thermostat are given by a Poisson process whose intensity μ is chosen so that the average time between two successive interactions of a given particle with the thermostat is independent of the number of particles in the system. For the case where $\rho(\theta) = (2\pi)^{-1}$, the entropy decays exponentially fast. In fact, abbreviating $\sqrt{\beta/(2\pi)}e^{-\beta/2\mathbf{v}^2} = \Gamma_\beta(\mathbf{v})$, we know from [7], that

$$S(f(\cdot, t)) := \int_{\mathbb{R}^M} f(\mathbf{v}, t) \log \left(\frac{f(\mathbf{v}, t)}{\Gamma_\beta(\mathbf{v})} \right) d\mathbf{v} \leq e^{-\mu t/2} S(f_0) .$$

Thus, one might guess that if a “small” *system* of M particles out of equilibrium interacts with a *reservoir*, that is a large system of $N \geq M$ particles in thermal equilibrium, then the entropy decays exponentially fast in time. This intuition is also supported by the results in [6]. There it was shown that if the thermostat is replaced by a large but finite reservoir *initially* in thermal equilibrium, this evolution is close to the evolution given by the thermostat. This results holds in various norms and, in particular, it is *uniform* in time. We would like to emphasize that the reservoir will not stay in thermal equilibrium as time progresses, nevertheless it will not veer far from it.

Since this is the model that we consider in this work, we will now describe it in detail. We consider probability distributions $F : \mathbb{R}^{M+N} \rightarrow \mathbb{R}_+$ and write $F(\mathbf{v}, \mathbf{w})$ where $\mathbf{v} = (v_1, \dots, v_M)$ describes the particles in the small system, whereas $\mathbf{w} = (w_{M+1}, \dots, w_{N+M})$ describes the particles in the large system. The Kac master equation is given by

$$\frac{\partial F}{\partial t} = \mathcal{L}F , \quad F(\mathbf{v}, \mathbf{w}, 0) = F_0(\mathbf{v}, \mathbf{w}) = f_0(\mathbf{v})e^{-\pi|\mathbf{w}|^2} , \quad (2)$$

where

$$\mathcal{L} = \frac{\lambda_S}{M-1} \sum_{1 \leq i < j \leq M} (R_{ij} - I) + \frac{\lambda_R}{N-1} \sum_{M < i < j \leq N+M} (R_{ij} - I) + \frac{\mu}{N} \sum_{i=1}^M \sum_{j=M+1}^{M+N} (R_{ij} - I) , \quad (3)$$

and R_{ij} is given as follows. For $1 \leq i < j \leq M$ we have

$$(R_{ij}F)(\mathbf{v}, \mathbf{w}) = \int_{-\pi}^{\pi} \rho(\theta) d\theta F(r_{ij}(\theta)^{-1}(\mathbf{v}, \mathbf{w})) ,$$

where

$$r_{ij}(\theta)^{-1}(\mathbf{v}, \mathbf{w}) = (v_1, \dots, v_i \cos \theta - v_j \sin \theta, \dots, v_i \sin \theta + v_j \cos \theta, \dots, v_M, \mathbf{w}) . \quad (4)$$

The other R_{ij} s are defined analogously. We assume that the probability measure ρ is smooth and satisfies

$$\int_{-\pi}^{\pi} \rho(\theta) d\theta \sin \theta \cos \theta = 0 . \quad (5)$$

In particular, we do *not* require \mathcal{L} to be self-adjoint on $L^2(\mathbb{R}^{N+M})$, a condition called *microscopic reversibility*. The initial state of the reservoir is assumed to be a thermal equilibrium state and we have chosen units in which the inverse temperature $\beta = 2\pi$. Note that λ_S is the rate at which one particle from the system will scatter with any other particle in the system and similarly for λ_R . Likewise, μ is the rate at which a single particle of the system will scatter with any particle in the reservoir. The rate at which a particular particle from the reservoir will scatter with a particle in the system is given by $\mu M/N$. Hence, when N is large compared to M this process is suppressed and one expects that the reservoir does not move far from its equilibrium. Indeed, it is shown in [6] that the solution of the master equation (3) stays close to the solution of a thermostated system in the Gabetta-Toscani-Wennberg metric,

$$d_{GTW}(F, G) := \sup_{k \neq 0} \frac{|\widehat{F}(k) - \widehat{G}(k)|}{|k|^2},$$

see [16]. Here, \widehat{F} denotes the Fourier transform of F . More precisely, with the initial conditions (1) and (2), it was shown that

$$d_{GTW}(f(\mathbf{v}, t)e^{-\pi|\mathbf{w}|^2}, F(\mathbf{v}, \mathbf{w}, t)) \leq C(f_0) \frac{M}{N},$$

where $C(f_0)$ is a constant that depends on the initial condition but is of order one. The distance varies inversely as N , the size of the reservoir and, moreover, this estimate holds *uniformly* in time. For a detailed description of the results we refer the reader to [6]. From this result and the fact that the entropy of the system interacting with a thermostat decays exponentially in time, one might surmise that the entropy of the system interacting with a finite reservoir also decays exponentially fast in time. In fact we shall show this to be true if we consider the entropy relative to the thermal state.

2. Results

For the solution of the master equation (2) we use interchangeably the notation

$$F(\mathbf{v}, \mathbf{w}, t) = (e^{\mathcal{L}t} F_0)(\mathbf{v}, \mathbf{w}). \quad (6)$$

This evolution preserves the energy and hence it suffices to consider it on $L^1(\mathbb{S}^{N+M}(\sqrt{N+M}))$ with the normalized surface measure. Likewise, it is easy to see that the evolution is ergodic on $\mathbb{S}^{N+M}(\sqrt{N+M})$ in the sense that $e^{\mathcal{L}t} F_0 \rightarrow 1$ as $t \rightarrow \infty$ and 1 is the only normalized equilibrium state.

For our purposes it is convenient to consider the evolution in $L^1(\mathbb{R}^{M+N})$ with Lebesgue measure. Then $e^{\mathcal{L}t} F_0$ converges to the spherical average of F_0 taken over spheres in \mathbb{R}^{M+N} . In this space we choose the initial condition

$$F_0(\mathbf{v}, \mathbf{w}) = f_0(\mathbf{v})e^{-\pi|\mathbf{w}|^2}. \quad (7)$$

Moreover, we introduce the function f ,

$$f(\mathbf{v}, t) := \int_{\mathbb{R}^N} [e^{\mathcal{L}t} F_0](\mathbf{v}, \mathbf{w}) d\mathbf{w} \quad (8)$$

and we call

$$S(f(\cdot, t)) := \int_{\mathbb{R}^M} f(\mathbf{v}, t) \log \left(\frac{f(\mathbf{v}, t)}{e^{-\pi|\mathbf{v}|^2}} \right) d\mathbf{v},$$

the entropy of f relative to the thermal state $e^{-\pi|\mathbf{v}|^2}$. Our main result is the following theorem.

Theorem 2.1. *Let $N \geq M$ and let ρ be a probability distribution with an absolutely convergent Fourier series such that (5) holds. The entropy of f relative of to the thermal state $e^{-\pi|v|^2}$ then satisfies*

$$S(f(\cdot, t)) \leq \left[\frac{M}{N+M} + \frac{N}{N+M} e^{-t\mu_\rho(N+M)/N} \right] S(f_0) ,$$

where

$$\mu_\rho = \mu \int_{-\pi}^{\pi} \rho(\theta) d\theta \sin^2(\theta) ,$$

and f_0 is as introduced in (7).

Remark 2.2.

1. Note that the theorem deals with the entropy relative to the thermal state and not with respect to the equilibrium state. The entropy relative to the equilibrium state tends to zero as $t \rightarrow \infty$. We do not know how to adapt our proof to this situation nor do we have any evidence that it does indeed tend to zero at an exponential rate. If this were the case, the rate would most likely depend on the initial condition.

2. The decay rate is universal in the sense that it only depends on μ and the distribution ρ . The intra-particle interactions in the system and in the reservoir do not seem to matter.

3. The statement of the theorem becomes particularly simple as $N \rightarrow \infty$. This corresponds to the thermostat problem treated in [7] with the exact same decay rate. It is known that for the thermostat the decay rate is optimal, see [28], and hence the decay rate here is optimal as well.

4. Although we assume that ρ is smooth, our result also holds for the case where ρ is a finite sum of Dirac measures. In particular Theorem 2.1 also holds if ρ is a delta measure that has its mass at the angles $\theta = \pm\pi/2$, that is, our result does not depend on ergodicity of the evolution.

As a consequence of Remark 2.2(2), one obtains a result for the standard Kac model. Recall that the generator of the standard Kac model is given by

$$\mathcal{L}_{\text{cl}} = \frac{2}{N+M-1} \sum_{1 \leq i < j \leq N+M} (R_{ij} - I) .$$

We may arbitrarily split the variables into two groups (v_1, \dots, v_M) and $(w_{M+1}, \dots, w_{M+N})$. Splitting the generator accordingly,

$$\begin{aligned} \mathcal{L}_{\text{cl}} &= \frac{2}{N+M-1} \sum_{1 \leq i < j \leq M} (R_{ij} - I) + \frac{2}{N+M-1} \sum_{M+1 \leq i < j \leq N+M} (R_{ij} - I) \\ &\quad + \frac{2}{N+M-1} \sum_{i=1}^M \sum_{j=M+1}^{N+M} (R_{ij} - I) , \end{aligned}$$

we see that the standard Kac model can be cast in the form (3) by setting

$$\lambda_S = \frac{2(M-1)}{N+M-1} , \quad \lambda_R = \frac{2(N-1)}{N+M-1} \quad \text{and} \quad \mu = \frac{2N}{N+M-1} .$$

Hence, we obtain the following Corollary.

Corollary 2.3. *Let $N \geq M$ and consider the time evolution defined by \mathcal{L}_{cl} with initial condition (7). Assume that the function f_0 in the initial condition has finite entropy. The entropy of the*

function

$$f(\mathbf{v}, t) := \int_{\mathbb{R}^N} [e^{\mathcal{L}t} F_0](\mathbf{v}, \mathbf{w}) d\mathbf{w}$$

relative to the thermal state $e^{-\pi|\mathbf{v}|^2}$, satisfies

$$S(f(\cdot, t)) \leq \left[\frac{M}{N+M} + \frac{N}{N+M} e^{-t\mu_\rho 2(N+M)/(N+M-1)} \right] S(f_0),$$

where

$$\mu_\rho = \int_{-\pi}^{\pi} \rho(\theta) d\theta \sin^2(\theta)$$

and ρ is a probability distribution such that (5) holds.

On a mathematical level, an efficient way of proving approach to equilibrium is through a logarithmic Sobolev inequality, which presupposes that the generator of the time evolution is given by a Dirichlet form. This kind of structure is notably absent in the Kac master equation. We shall see however, that the logarithmic Sobolev inequality in the form of Nelson's hypercontractive estimate is an important tool for the proof of Theorem 2.1. We will use an iterated version of it, which expresses the result in terms of marginals of the functions involved. This, coupled with an auxiliary computation and a sharp version of the Brascamp-Lieb inequalities [8] (see also [22]) will lead to the result.

In our opinion, the main result of this paper is the description of a simple mechanism for obtaining exponential relaxation towards equilibrium. One can extend the results to three dimensional momentum preserving collisions, however, so far only for a caricature of Maxwellian molecules. To carry this method over to the case of hard spheres and for true Maxwellian molecules is an open problem.

The plan of the paper is as follows: In Section 3 we derive a representation formula for the Kac evolution $e^{\mathcal{L}t}$ which is reminiscent of the Ornstein-Uhlenbeck process. This allows us to prove an entropy inequality based upon Nelson's hypercontractive estimate in Section 4. In Section 5 we show how the sharp version of the geometric Brascamp-Lieb inequality leads to a correlation inequality for the entropy involving marginals, which in turn proves our main entropy inequality. The fact that our Brascamp-Lieb datum is geometric relies on a sum rule which will be proved in Section 6. A short proof of the geometric form of the Brascamp-Lieb inequalities is deferred to Appendix A, as well as some technical details to ensure its applicability in Appendix B. In Section 7 we show how our method can be applied to three-dimensional Maxwellian collisions with a very simple angular dependence.

Acknowledgements: A.G. and T.R. would like to thank Georgia Tech for its hospitality. The work of M.L. and A.G. was supported in part by NSF grant DMS-1600560 and the Humboldt Foundation. A.G. and T.R. gratefully acknowledge financial support by the Deutsche Forschungsgemeinschaft (DFG) through GRK 1838 (A.G.) and CRC 1173 (T.R.). F.B. gratefully acknowledges financial support from the Simons Foundation award number 359963. T.R. thanks the Karlsruhe House of Young Scientists (KHYS) for a Research Travel Grant supporting the stay at Georgia Tech.

3. The representation formula

The aim of this section is to rewrite (6), that is $e^{\mathcal{L}t} F_0$, in a way which is reminiscent of the Ornstein-Uhlenbeck process. This representation will naturally lead to the next step in the proof of Theorem 2.1, namely the entropy inequality that will be presented in Theorem 4.1.

It is convenient to write

$$\mathcal{L} = \Lambda(Q - I) , \text{ where } \Lambda = \lambda_S \frac{M}{2} + \lambda_R \frac{N}{2} + \mu M ,$$

and the operator Q is a convex combination of R_{ij} s, given by

$$Q = \frac{\lambda_S}{\Lambda(M-1)} \sum_{1 \leq i < j \leq M} R_{ij} + \frac{\lambda_R}{\Lambda(N-1)} \sum_{M < i < j \leq M+N} R_{ij} + \frac{\mu}{\Lambda N} \sum_{i=1}^M \sum_{j=M+1}^{M+N} R_{ij} ,$$

i.e., Q is an average over rotation operators. The right hand side of (6) can be written as

$$(e^{\mathcal{L}t} F_0)(\mathbf{v}, \mathbf{w}) = e^{-\Lambda t} \sum_{k=0}^{\infty} \frac{t^k \Lambda^k}{k!} Q^k F_0(\mathbf{v}, \mathbf{w}) , \quad (9)$$

where

$$Q^k F_0(\mathbf{v}, \mathbf{w}) = \sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{[-\pi, \pi]^k} \rho(\theta_1) d\theta_1 \cdots \rho(\theta_k) d\theta_k F_0 \left(\left[\prod_{l=1}^k r_{\alpha_l}(\theta_l) \right]^{-1} (\mathbf{v}, \mathbf{w}) \right) . \quad (10)$$

Here, α labels pairs of particles, that is, $\alpha = (i, j)$, $1 \leq i < j \leq M + N$, $r_{\alpha}(\theta)$ is defined in (4) and λ_{α} is given by the rotation corresponding to the index α , that is,

$$\begin{aligned} \lambda_{(i,j)} &= \frac{\lambda_S}{\Lambda(M-1)} \quad \text{if } 1 \leq i < j \leq M , \\ \lambda_{(i,j)} &= \frac{\lambda_R}{\Lambda(N-1)} \quad \text{if } M+1 \leq i < j \leq M+N , \\ \lambda_{(i,j)} &= \frac{\mu}{\Lambda N} \quad \text{if } 1 \leq i \leq M , M+1 \leq j \leq M+N . \end{aligned}$$

Note that the sum over *all* pairs $\sum_{\alpha} \lambda_{\alpha} = 1$.

For our purpose, it is convenient to write the function f_0 , introduced in (7), as $f_0(\mathbf{v}) = h_0(\mathbf{v})e^{-\pi|\mathbf{v}|^2}$. Since the Gaussian function is invariant under rotations, (9) takes the form

$$(e^{\mathcal{L}t} F_0)(\mathbf{v}, \mathbf{w}) = e^{-\pi(|\mathbf{v}|^2 + |\mathbf{w}|^2)} e^{-\Lambda t} \sum_{k=0}^{\infty} \frac{t^k \Lambda^k}{k!} Q^k (h_0 \circ P)(\mathbf{v}, \mathbf{w}) .$$

We introduce the projection $P : \mathbb{R}^{N+M} \rightarrow \mathbb{R}^M$ by $P(\mathbf{v}, \mathbf{w}) = \mathbf{v}$, as a reminder that the semigroup $e^{\mathcal{L}t}$ acts on functions that depend on \mathbf{v} as well as \mathbf{w} . If we write

$$f(\mathbf{v}, t) = e^{-\pi|\mathbf{v}|^2} h(\mathbf{v}, t) ,$$

then (8) can be written as

$$h(\mathbf{v}, t) = e^{-\Lambda t} \sum_{k=0}^{\infty} \frac{t^k \Lambda^k}{k!} h_k(\mathbf{v}) ,$$

where the functions h_k are given by

$$h_k(\mathbf{v}) := \int_{\mathbb{R}^N} Q^k (h_0 \circ P)(\mathbf{v}, \mathbf{w}) e^{-\pi|\mathbf{w}|^2} d\mathbf{w} .$$

Likewise, the entropy of f is expressed as

$$S(f(\cdot, t)) = \int_{\mathbb{R}^M} h(\mathbf{v}, t) \log h(\mathbf{v}, t) e^{-\pi|\mathbf{v}|^2} d\mathbf{v} =: \mathcal{S}(h(\cdot, t)) .$$

Expanding the function $Q^k(h_0 \circ P)(\mathbf{v}, \mathbf{w})$, we find that

$$h_k(\mathbf{v}) = \sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{[-\pi, \pi]^k} \rho(\theta_1) d\theta_1 \cdots \rho(\theta_k) d\theta_k \times \\ \times \int_{\mathbb{R}^N} (h_0 \circ P) \left(\left[\prod_{l=1}^k r_{\alpha_l}(\theta_l) \right]^{-1} (\mathbf{v}, \mathbf{w}) \right) e^{-\pi|\mathbf{w}|^2} d\mathbf{w}, \quad (11)$$

where, as before, see (10), $r_\alpha(\theta)$ rotates the plane given by the index pair α by an angle θ while keeping the other directions fixed. Since $P(\mathbf{v}, \mathbf{w}) = \mathbf{v}$, it is natural to write

$$\left[\prod_{j=1}^k r_{\alpha_j}(\theta_j) \right]^{-1} = \begin{pmatrix} A_k(\underline{\alpha}, \underline{\theta}) & B_k(\underline{\alpha}, \underline{\theta}) \\ C_k(\underline{\alpha}, \underline{\theta}) & D_k(\underline{\alpha}, \underline{\theta}) \end{pmatrix},$$

where $A_k \in \mathbb{R}^{M \times M}$ is an $M \times M$ matrix, $B_k \in \mathbb{R}^{M \times N}$, $C_k \in \mathbb{R}^{N \times M}$ and $D_k \in \mathbb{R}^{N \times N}$. Further, $\underline{\alpha} = (\alpha_1, \dots, \alpha_k)$ and $\underline{\theta} = (\theta_1, \dots, \theta_k)$. This notation allows us to rewrite (11) as

$$h_k(\mathbf{v}) = \sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{[-\pi, \pi]^k} \rho(\theta_1) d\theta_1 \cdots \rho(\theta_k) d\theta_k \times \\ \times \int_{\mathbb{R}^N} h_0(A_k(\underline{\alpha}, \underline{\theta})\mathbf{v} + B_k(\underline{\alpha}, \underline{\theta})\mathbf{w}) e^{-\pi|\mathbf{w}|^2} d\mathbf{w}.$$

Note that, by the definition of rotations,

$$A_k(\underline{\alpha}, \underline{\theta})A_k^T(\underline{\alpha}, \underline{\theta}) + B_k(\underline{\alpha}, \underline{\theta})B_k^T(\underline{\alpha}, \underline{\theta}) = I_M. \quad (12)$$

Lemma 3.1. *Let $A \in \mathbb{R}^{M \times M}$ and $B \in \mathbb{R}^{M \times N}$ be matrices that satisfy $AA^T + BB^T = I_M$. Then*

$$\int_{\mathbb{R}^N} h(A\mathbf{v} + B\mathbf{w})e^{-\pi|\mathbf{w}|^2} d\mathbf{w} = \int_{\mathbb{R}^M} h(A\mathbf{v} + (I_M - AA^T)^{1/2}\mathbf{u})e^{-\pi|\mathbf{u}|^2} d\mathbf{u}$$

for any integrable function h .

Proof. Denote the range of B by $H \subset \mathbb{R}^M$ and its kernel by $K \subset \mathbb{R}^N$. We may write

$$\int_{\mathbb{R}^N} h(A\mathbf{v} + B\mathbf{w})e^{-\pi|\mathbf{w}|^2} d\mathbf{w} = \int_K \int_{K^\perp} h(A\mathbf{v} + B\mathbf{u})e^{-\pi|\mathbf{u}|^2} e^{-\pi|\mathbf{u}'|^2} d\mathbf{u}d\mathbf{u}' \\ = \int_{K^\perp} h(A\mathbf{v} + B\mathbf{u})e^{-\pi|\mathbf{u}|^2} d\mathbf{u}.$$

The symmetric map $BB^T : \mathbb{R}^M \rightarrow \mathbb{R}^M$ has H as its range and H^\perp , that is the orthogonal complement of H in \mathbb{R}^M , as its kernel. Indeed, suppose that there exists $x \in \mathbb{R}^M$ with $BB^T x = 0$, then $B^T x = 0$, i.e., $x \in \text{Ker} B^T$ or x is perpendicular to H . Hence, the map $BB^T : H \rightarrow H$ is invertible. Define the linear map $R : \mathbb{R}^N \rightarrow H$ by

$$R = (BB^T)^{-1/2} B$$

and note that $RR^T = I_H$ while $R^T R$ projects the space K^\perp orthogonally onto H . Since K^\perp and H have the same dimension, it follows that R^T restricted to H defines an isometry between

H and K^\perp . Hence,

$$\begin{aligned} \int_{K^\perp} h(\mathbf{A}\mathbf{v} + B\mathbf{u}) e^{-\pi|\mathbf{u}|^2} d\mathbf{u} &= \int_{K^\perp} h\left(\mathbf{A}\mathbf{v} + (BB^T)^{1/2} R\mathbf{u}\right) e^{-\pi|\mathbf{u}|^2} d\mathbf{u} \\ &= \int_H h\left(\mathbf{A}\mathbf{v} + (BB^T)^{1/2} RR^T\mathbf{u}\right) e^{-\pi|R^T\mathbf{u}|^2} d\mathbf{u} \\ &= \int_H h\left(\mathbf{A}\mathbf{v} + (BB^T)^{1/2} \mathbf{u}\right) e^{-\pi|\mathbf{u}|^2} d\mathbf{u} . \end{aligned}$$

The assumption $AA^T + BB^T = I_M$, together with the fact that

$$\int_H h\left(\mathbf{A}\mathbf{v} + (BB^T)^{1/2} \mathbf{u}\right) e^{-\pi|\mathbf{u}|^2} d\mathbf{u} = \int_{H^\perp} \int_H h\left(\mathbf{A}\mathbf{v} + (BB^T)^{1/2} \mathbf{u}\right) e^{-\pi|\mathbf{u}|^2} d\mathbf{u} e^{-\pi|\mathbf{u}'|^2} d\mathbf{u}'$$

now implies the lemma. \square

The matrix $A_k(\underline{\alpha}, \underline{\theta})$ has an orthogonal singular value decomposition,

$$A_k(\underline{\alpha}, \underline{\theta}) = U_k(\underline{\alpha}, \underline{\theta}) \Gamma_k(\underline{\alpha}, \underline{\theta}) V_k^T(\underline{\alpha}, \underline{\theta}) , \quad (13)$$

where $\Gamma_k(\underline{\alpha}, \underline{\theta}) = \text{diag}[\gamma_{k,1}(\underline{\alpha}, \underline{\theta}), \dots, \gamma_{k,M}(\underline{\alpha}, \underline{\theta})]$ is the diagonal matrix whose entries $\gamma_{k,j}(\underline{\alpha}, \underline{\theta})$, $j = 1, \dots, M$, are the singular values of $A_k(\underline{\alpha}, \underline{\theta})$, and $U_k(\underline{\alpha}, \underline{\theta})$ and $V_k(\underline{\alpha}, \underline{\theta})$ are rotations in \mathbb{R}^M . Note that (12) implies $\gamma_{k,j}(\underline{\alpha}, \underline{\theta}) \in [0, 1]$ for $j = 1, \dots, M$. We shall use the abbreviation

$$h_0(U_k(\underline{\alpha}, \underline{\theta})\mathbf{v}) = h_{0,U_k(\underline{\alpha}, \underline{\theta})}(\mathbf{v}) .$$

These considerations can be summarized by the representation formula presented in the following theorem.

Theorem 3.2 (Representation formula). *The function h_k can be written as*

$$\begin{aligned} h_k(\mathbf{v}) &= \sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{[-\pi, \pi]^k} \rho(\theta_1) d\theta_1 \cdots \rho(\theta_k) d\theta_k \times \\ &\quad \times \int_{\mathbb{R}^M} h_{0,U_k(\underline{\alpha}, \underline{\theta})} \left(\Gamma_k(\underline{\alpha}, \underline{\theta}) V_k^T(\underline{\alpha}, \underline{\theta}) \mathbf{v} + (I_M - \Gamma_k^2(\underline{\alpha}, \underline{\theta}))^{1/2} \mathbf{w} \right) e^{-\pi|\mathbf{w}|^2} d\mathbf{w} , \quad (14) \end{aligned}$$

where $h_{0,U_k(\underline{\alpha}, \underline{\theta})}$, $\Gamma_k(\underline{\alpha}, \underline{\theta})$ and V_k are as defined above.

4. The hypercontractive estimate

Starting from (14) and using convexity of the entropy and Jensen's inequality together with

$$\sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{[-\pi, \pi]^k} \rho(\theta_1) d\theta_1 \cdots \rho(\theta_k) d\theta_k = 1 ,$$

we get

$$\mathcal{S}(h_k) \leq \sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{[-\pi, \pi]^k} \rho(\theta_1) d\theta_1 \cdots \rho(\theta_k) d\theta_k \mathcal{S}(g_k(\cdot, \underline{\alpha}, \underline{\theta})),$$

where we set

$$g_k(\mathbf{v}, \underline{\alpha}, \underline{\theta}) = \int_{\mathbb{R}^M} h_{0,U_k(\underline{\alpha}, \underline{\theta})} \left(\gamma_k(\underline{\alpha}, \underline{\theta}) \mathbf{v} + (I_M - \gamma_k^2(\underline{\alpha}, \underline{\theta}))^{1/2} \mathbf{w} \right) e^{-\pi|\mathbf{w}|^2} d\mathbf{w} , \quad (15)$$

and we removed the rotation $V_k^T(\underline{\alpha}, \underline{\theta})$ by a change of variables.

To explain the main observation in this section we look at (15) when $M = 1$. Since $0 \leq \gamma_k(\underline{\alpha}, \underline{\theta}) \leq 1$, we can write $\gamma_k(\underline{\alpha}, \underline{\theta}) = e^{-t}$ and we get $g_k(v, \underline{\alpha}, \underline{\theta}) = N_t(h_{0, U_k(\underline{\alpha}, \underline{\theta})})$ where N_t is the Ornstein-Uhlenbeck semigroup, that is

$$N_t h(x) = \int_{\mathbb{R}} h\left(e^{-t}x + \sqrt{1 - e^{-2t}}y\right) e^{-\pi y^2} dy .$$

Thus Theorem 3.2 renders the function h_k as a convex combination of terms reminiscent of the Ornstein-Uhlenbeck process, albeit in matrix form. We make use of this observation to find a bound for $\mathcal{S}(g_k(\cdot, \underline{\alpha}, \underline{\theta}))$. This bound together with a suitable correlation inequality proved in the next section will lead to a bound for $\mathcal{S}(h_k)$.

In addition to the notation developed in the previous section, we need various marginals of the function $h_{0, U_k(\underline{\alpha}, \underline{\theta})}$. Quite generally, if h is a function of M variables and $\sigma \subset \{1, \dots, M\}$, we shall denote by h^σ the marginals of h with respect to the variables $v_j, j \in \sigma$, for instance,

$$h^{\{1,2\}}(v_3, \dots, v_M) = \int_{\mathbb{R}^2} h(v_1, v_2, v_3, \dots, v_M) e^{-\pi(v_1^2 + v_2^2)} dv_1 dv_2 .$$

It will be convenient to use the matrix $P_\sigma : \mathbb{R}^M \rightarrow \mathbb{R}^{|\sigma|}$ that projects \mathbb{R}^M orthogonally onto $\mathbb{R}^{|\sigma|}$ which we will identify with subspace of \mathbb{R}^M . To give an example, let $\mathbf{v} = (v_1, \dots, v_M)$. Then $P_{\{1,2\}}\mathbf{v} = (v_1, v_2)$. The following theorem is the main result of this section.

Theorem 4.1 (Partial entropy bound). *Let $h_0 \in L^1(\mathbb{R}^M, e^{-\pi|\mathbf{v}|^2} d\mathbf{v})$ be nonnegative and assume that $\mathcal{S}(h_0) < \infty$. Then*

$$\begin{aligned} & \mathcal{S}(g_k(\cdot, \underline{\alpha}, \underline{\theta})) \\ & \leq \sum_{\sigma \subset \{1, \dots, M\}} \prod_{i \in \sigma^c} \gamma_{k,i}^2 \prod_{j \in \sigma} (1 - \gamma_{k,j}^2) \int_{\mathbb{R}^M} h_0(\mathbf{v}) \log h_{0, U_k(\underline{\alpha}, \underline{\theta})}^\sigma (P_{\sigma^c} U_k(\underline{\alpha}, \underline{\theta})^T \mathbf{v}) e^{-\pi|\mathbf{v}|^2} d\mathbf{v} , \end{aligned} \quad (16)$$

where σ^c is the complement of the set σ in $\{1, \dots, M\}$.

A key role in the proof of Theorem 4.1 is played by Nelson's hypercontractive estimate.

Theorem 4.2 (Nelson's hypercontractive estimate). *The Ornstein-Uhlenbeck semigroup,*

$$N_t h(x) = \int_{\mathbb{R}} h\left(e^{-t}x + \sqrt{1 - e^{-2t}}y\right) e^{-\pi y^2} dy ,$$

for $t \geq 0$, is bounded from $L^p(\mathbb{R}, e^{-\pi x^2} dx)$ to $L^q(\mathbb{R}, e^{-\pi x^2} dx)$ if and only if

$$(p - 1) \geq e^{-2t}(q - 1) .$$

For such values of p and q ,

$$\|N_t h\|_q \leq \|h\|_p$$

with equality if and only if h is constant.

Proof. For a proof we refer the reader to [27]. For other proofs see [17, 18, 15, 12]. \square

Nelson's hypercontractive estimate, that is Theorem 4.2, implies the following Corollary, which will be useful in the proof of Theorem 4.1.

Corollary 4.3 (Entropic version of Nelson's hypercontractive estimate). *Let $h : \mathbb{R} \rightarrow \mathbb{R}_+$ be a function in $L^1(\mathbb{R}, e^{-\pi x^2} dx)$ with finite entropy, i.e.,*

$$\mathcal{S}(h) = \int_{\mathbb{R}} h(x) \log h(x) e^{-\pi x^2} dx < \infty .$$

Then

$$\mathcal{S}(N_t h) \leq e^{-2t} \mathcal{S}(h) + (1 - e^{-2t}) \|h\|_1 \log \|h\|_1$$

for all $t \geq 0$.

Proof. Let $h \in L^p(\mathbb{R}, e^{-\pi x^2} dx)$, for $p \geq 1$ small, be a nonnegative function. As $\|N_t h\|_1 = \|h\|_1$, we can apply Nelson's hypercontractive estimate, which implies that for p, q that satisfy $(p - 1) = e^{-2t}(q - 1)$,

$$\frac{\|N_t h\|_q - \|N_t h\|_1}{q - 1} \leq \frac{\|h\|_p - \|h\|_1}{p - 1} = e^{-2t} \frac{\|h\|_p - \|h\|_1}{p - 1} .$$

Sending $p \rightarrow 1$ and hence $q \rightarrow 1$, we get the claimed estimate for such functions h . If h just has finite entropy one cuts off h at large values, uses the above estimate and removes the cutoff using the monotone convergence theorem. \square

We are now ready to prove Theorem 4.1.

Proof of Theorem 4.1. Remember that $0 \leq \gamma_{k,j}(\underline{\alpha}, \underline{\theta}) \leq 1$ for $j = 1, \dots, M$. Thus, by inductively applying Corollary 4.3 to

$$\int_{\mathbb{R}^M} h_{0,U_k(\underline{\alpha}, \underline{\theta})} \left(\gamma_{k,1} v_1 + \sqrt{1 - \gamma_{k,1}^2} u_1, \dots, \gamma_{k,M} v_M + \sqrt{1 - \gamma_{k,M}^2} u_M \right) e^{-\pi \sum_{j=1}^M u_j^2} du_1 \cdots du_M ,$$

we obtain

$$\mathcal{S}(g_k(\cdot, \underline{\alpha}, \underline{\theta})) \leq \sum_{\sigma \subset \{1, \dots, M\}} \prod_{i \in \sigma^c} \gamma_{k,i}^2 \prod_{j \in \sigma} (1 - \gamma_{k,j}^2) \int_{\mathbb{R}^{|\sigma^c|}} h_{0,U_k(\underline{\alpha}, \underline{\theta})}^\sigma(\mathbf{u}) \log h_{0,U_k(\underline{\alpha}, \underline{\theta})}^\sigma(\mathbf{u}) e^{-\pi |\mathbf{u}|^2} d\mathbf{u} .$$

Inserting the definition of the marginal $h_{0,U_k(\underline{\alpha}, \underline{\theta})}^\sigma$, we see that

$$\begin{aligned} \int_{\mathbb{R}^{|\sigma^c|}} h_{0,U_k(\underline{\alpha}, \underline{\theta})}^\sigma(\mathbf{u}) \log h_{0,U_k(\underline{\alpha}, \underline{\theta})}^\sigma(\mathbf{u}) e^{-\pi |\mathbf{u}|^2} d\mathbf{u} &= \int_{\mathbb{R}^M} h_{0,U_k(\underline{\alpha}, \underline{\theta})}^\sigma(P_{\sigma^c} \mathbf{v}) \log h_{0,U_k(\underline{\alpha}, \underline{\theta})}^\sigma(P_{\sigma^c} \mathbf{v}) e^{-\pi |\mathbf{v}|^2} d\mathbf{v} \\ &= \int_{\mathbb{R}^M} h_{0,U_k(\underline{\alpha}, \underline{\theta})}(\mathbf{v}) \log h_{0,U_k(\underline{\alpha}, \underline{\theta})}^\sigma(P_{\sigma^c} \mathbf{v}) e^{-\pi |\mathbf{v}|^2} d\mathbf{v} \\ &= \int_{\mathbb{R}^M} h_0(\mathbf{v}) \log h_{0,U_k(\underline{\alpha}, \underline{\theta})}^\sigma(P_{\sigma^c} U_k(\underline{\alpha}, \underline{\theta})^T \mathbf{v}) e^{-\pi |\mathbf{v}|^2} d\mathbf{v}, \end{aligned}$$

which finishes the proof of Theorem 4.1. \square

5. The key entropy bound

Collecting the results of the previous sections we get the following bound

$$\begin{aligned} \mathcal{S}(h_k) &\leq \sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{[-\pi, \pi]^k} \rho(\theta_1) d\theta_1 \cdots \rho(\theta_k) d\theta_k \times \\ &\quad \times \sum_{\sigma \subset \{1, \dots, M\}} \prod_{i \in \sigma^c} \gamma_{k,i}^2 \prod_{j \in \sigma} (1 - \gamma_{k,j}^2) \int_{\mathbb{R}^M} h_0(\mathbf{v}) \log h_{0, U_k(\underline{\alpha}, \underline{\theta})}^\sigma (P_{\sigma^c} U_k(\underline{\alpha}, \underline{\theta})^T \mathbf{v}) e^{-\pi|\mathbf{v}|^2} d\mathbf{v}. \end{aligned} \quad (17)$$

The right-hand side of (17) contains a large sum over the entropy of marginals of h_0 . In order to bound such a sum in terms of the entropy of h_0 one may try to apply some version of the Loomis-Whitney inequality [23] or, more precisely, of an inequality by Han [19]. This is essentially correct, but will require a substantial generalization of this inequality. Let us first formulate the main theorem of this section.

Theorem 5.1 (Entropy bound). *The estimate*

$$\mathcal{S}(h_k) \leq \left[\frac{M}{N+M} + \frac{N}{N+M} \left(1 - \mu_\rho \frac{N+M}{N\Lambda} \right)^k \right] \mathcal{S}(h_0) \quad (18)$$

holds.

As mentioned before, to prove Theorem 5.1, we need a generalized version of an inequality by Han. This generalization was proven by Carlen-Cordero-Erausquin in [11]. It is based on the geometric Brascamp-Lieb inequality due to Ball [1], see also [2], in the rank one case, and due to Barthe [3] in the general case.

Theorem 5.2 (Correlation inequality). *For $i = 1, \dots, K$, let $H_i \subset \mathbb{R}^M$ be subspaces of dimension d_i and $B_i : \mathbb{R}^M \rightarrow H_i$ be linear maps with the property that $B_i B_i^T = I_{H_i}$, the identity map on H_i . Assume further that there are non-negative constants $c_i, i = 1, \dots, K$ such that*

$$\sum_{i=1}^K c_i B_i^T B_i = I_M. \quad (19)$$

Then, for nonnegative functions $f_i : H_i \rightarrow \mathbb{R}$,

$$\int_{\mathbb{R}^M} \prod_{i=1}^K f_i^{c_i}(B_i \mathbf{v}) e^{-\pi|\mathbf{v}|^2} d\mathbf{v} \leq \prod_{i=1}^K \left(\int_{H_i} f_i(u) e^{-\pi|u|^2} du \right)^{c_i}. \quad (20)$$

Moreover,

$$\int_{\mathbb{R}^M} h(\mathbf{v}) \log h(\mathbf{v}) e^{-\pi|\mathbf{v}|^2} d\mathbf{v} \geq \sum_{i=1}^K c_i \left[\int_{\mathbb{R}^M} h(\mathbf{v}) \log f_i(B_i \mathbf{v}) e^{-\pi|\mathbf{v}|^2} d\mathbf{v} - \log \int_{H_i} f_i(u) e^{-\pi|u|^2} du \right], \quad (21)$$

for any nonnegative function $h \in L^1(\mathbb{R}^M, e^{-\pi|\mathbf{v}|^2} d\mathbf{v})$.

Since Theorem 5.2 is very useful in a number of applications, and for the readers convenience, we will give an elementary proof in Appendix A.

Remark 5.3. *By taking the trace in (19) one sees that*

$$\sum_{i=1}^K c_i d_i = M .$$

We would like to apply (21) to the right hand side of (17). An immediate problem is that (17) is in terms of integrals and not sums. While there are some results available for continuous indices (see, e.g., [4]), they do not apply to our situation and hence we will take a more direct approach and approximate the measure $\rho(\theta)d\theta$ by a discrete measure. It is important that the approximation also satisfies the constraint (5). The following lemma establishes such an approximation. Its proof is given in Appendix B.

Lemma 5.4. *Let ρ be a probability density on $[-\pi, \pi]$ whose Fourier series converges absolutely and assume that (5) is satisfied. There exists a sequence of discrete probability measures ν_K , $K = 1, 2, \dots$, such that for every continuous function f on $[-\pi, \pi]$*

$$\lim_{K \rightarrow \infty} \int_{-\pi}^{\pi} f(\theta) \nu_K(d\theta) = \int_{-\pi}^{\pi} f(\theta) \rho(\theta) d\theta .$$

Moreover,

$$\int_{-\pi}^{\pi} \cos \theta \sin \theta \nu_K(d\theta) = 0 ,$$

for all $K \in \mathbb{N}$. More precisely,

$$\nu_K(d\theta) = \frac{2\pi}{4K+1} \sum_{\ell=-2K}^{2K} \rho_K \left(\frac{2\pi\ell}{4K+1} \right) \delta \left(\theta - \frac{2\pi\ell}{4K+1} \right) d\theta ,$$

where

$$\rho_K(\theta) = \int_{-\pi}^{\pi} \rho(\theta - \phi) p_K(\theta) d\phi \quad \text{and} \quad p_K(\theta) := \frac{1}{2K+1} \left(\sum_{k=-K}^K e^{ik\theta} \right)^2 .$$

At this point we can prepare the ground for the application of Theorem 5.2 to inequality (17). We first replace $\rho(\theta)d\theta$ in (17) with $\nu_K(d\theta)$. Setting

$$\omega_{\ell_j} = \rho_K(\theta_j) , \quad \theta_{\ell_j} = \frac{2\pi\ell_j}{4K+1} , \quad \text{and} \quad \underline{\theta} = (\theta_{\ell_1}, \dots, \theta_{\ell_k}) ,$$

we obtain

$$\begin{aligned} & \sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{[-\pi, \pi]^k} \nu_K(d\theta_1) \cdots \nu_K(d\theta_k) \sum_{\sigma \subset \{1, \dots, M\}} \prod_{i \in \sigma^c} \gamma_{k,i}(\underline{\alpha}, \underline{\theta})^2 \prod_{j \in \sigma} (1 - \gamma_{k,j}(\underline{\alpha}, \underline{\theta})^2) \times \\ & \quad \times \int_{\mathbb{R}^M} h_0(\mathbf{v}) \log h_{0, U_k(\underline{\alpha}, \underline{\theta})}^{\sigma} (P_{\sigma^c} U_k(\underline{\alpha}, \underline{\theta})^T \mathbf{v}) e^{-\pi|\mathbf{v}|^2} d\mathbf{v} \\ & = \sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \sum_{-K \leq \ell_1, \dots, \ell_k \leq K} \prod_{j=1}^k \omega_{\ell_j} \sum_{\sigma \subset \{1, \dots, M\}} \prod_{i \in \sigma^c} \gamma_{k,i}(\underline{\alpha}, \underline{\theta})^2 \prod_{j \in \sigma} (1 - \gamma_{k,j}(\underline{\alpha}, \underline{\theta})^2) \times \\ & \quad \times \int_{\mathbb{R}^M} h_0(\mathbf{v}) \log h_{0, U_k(\underline{\alpha}, \underline{\theta})}^{\sigma} (P_{\sigma^c} U_k(\underline{\alpha}, \underline{\theta})^T \mathbf{v}) e^{-\pi|\mathbf{v}|^2} d\mathbf{v} . \end{aligned} \tag{22}$$

In order to apply Theorem 5.2 to (22) we have to replace the sum over the index i with a sum over the indices $\alpha_1, \dots, \alpha_k, \ell_1, \dots, \ell_k$ and all subsets $\sigma \subset \{1, \dots, M\}$. Moreover, we substitute

$$\begin{aligned} \text{the constants } c_i & \quad \text{by } \frac{1}{C_{k,M}} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \prod_{j=1}^k \omega_{\ell_j} \prod_{i \in \sigma^c} \gamma_{k,i}(\underline{\alpha}, \underline{\theta})^2 \prod_{j \in \sigma} (1 - \gamma_{k,j}(\underline{\alpha}, \underline{\theta})^2) , \\ \text{the functions } f_i(\mathbf{w}) & \quad \text{by } h_{0,U_k(\underline{\alpha}, \underline{\theta})}^\sigma(\mathbf{w}) , \\ \text{the linear maps } B_i & \quad \text{by } P_{\sigma^c} U_k(\underline{\alpha}, \underline{\theta})^T , \\ \text{the functions } f_i(B_i \mathbf{v}) & \quad \text{by } h_{0,U_k(\underline{\alpha}, \underline{\theta})}^\sigma(P_{\sigma^c} U_k(\underline{\alpha}, \underline{\theta})^T \mathbf{v}) , \\ \text{and the subspaces } H_i & \quad \text{by } \mathbb{R}^{|\sigma^c|} . \end{aligned}$$

For any given index i the condition $B_i B_i^T = I_{H_i}$ corresponds to $P_{\sigma^c} U_k(\underline{\alpha}, \underline{\theta})^T U_k(\underline{\alpha}, \underline{\theta}) P_{\sigma^c} = P_{\sigma^c}$ which is the identity on $\mathbb{R}^{|\sigma^c|}$.

The next theorem establishes the sum rule (19) in our setting and hence ensures the applicability of Theorem 5.2 to (22).

Theorem 5.5 (The sum rule). *If $\nu(d\theta)$ is a probability measure satisfying (5), then*

$$\begin{aligned} & \sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{[-\pi, \pi]^k} \nu(d\theta_1) \cdots \nu(d\theta_k) \times \\ & \times \sum_{\sigma \subset \{1, \dots, M\}} \prod_{i \in \sigma^c} \gamma_{k,i}(\underline{\alpha}, \underline{\theta})^2 \prod_{j \in \sigma} (1 - \gamma_{k,j}(\underline{\alpha}, \underline{\theta})^2) U_k(\underline{\alpha}, \underline{\theta}) P_{\sigma^c}^T P_{\sigma^c} U_k(\underline{\alpha}, \underline{\theta})^T = C_{k,M} I_M , \end{aligned} \quad (23)$$

where

$$C_{k,M} = \left[\frac{M}{N+M} + \frac{N}{N+M} \left(1 - \mu_\nu \frac{N+M}{N\Lambda} \right)^k \right]$$

with

$$\mu_\nu = \mu \int \nu(d\theta) \sin^2 \theta .$$

The proof will be given in Section 6. We observe here that it follows from Theorem 5.4 that $\mu_\rho = \lim_{K \rightarrow \infty} \mu_{\nu_K}$.

Proof of Theorem 5.1. First we consider the case where ρ is replaced by ν_K and use Theorem 5.2 together with Theorem 5.5 and the identification rules described above. The entropy inequality (21) now says that

$$\begin{aligned} & \int_{\mathbb{R}^M} h_0(\mathbf{v}) \log h_0(\mathbf{v}) e^{-\pi|\mathbf{v}|^2} d\mathbf{v} \\ & \geq \frac{1}{C_{k,M}} \sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \sum_{-K \leq \ell_1, \dots, \ell_k \leq K} \prod_{j=1}^k \omega_{\ell_j} \sum_{\sigma \subset \{1, \dots, M\}} \prod_{i \in \sigma^c} \gamma_{k,i}(\underline{\alpha}, \underline{\theta})^2 \prod_{j \in \sigma} (1 - \gamma_{k,j}(\underline{\alpha}, \underline{\theta})^2) \times \\ & \times \left[\int_{\mathbb{R}^M} h_0(\mathbf{v}) \log h_{0,U_k(\underline{\alpha}, \underline{\theta})}^\sigma(P_{\sigma^c} U_k(\underline{\alpha}, \underline{\theta})^T \mathbf{v}) e^{-\pi|\mathbf{v}|^2} d\mathbf{v} - \log \int_{\mathbb{R}^{|\sigma^c|}} h_{0,U_k(\underline{\alpha}, \underline{\theta})}^\sigma(u) e^{-\pi|u|^2} du \right] . \end{aligned}$$

However, since h_0 is normalized and $U_k(\underline{\alpha}, \underline{\theta})$ is orthogonal, we find that

$$\begin{aligned} \int_{\mathbb{R}^{|\sigma^c|}} h_{0,U_k(\underline{\alpha}, \underline{\theta})}^\sigma(u) e^{-\pi|u|^2} d u &= \int_{\mathbb{R}^{|\sigma^c|}} \int_{\mathbb{R}^{|\sigma|}} h_{0,U_k(\underline{\alpha}, \underline{\theta})}(v, u) e^{-\pi|v|^2} d v e^{-\pi|u|^2} d u \\ &= \int_{\mathbb{R}^M} h_0(U_k(\underline{\alpha}, \underline{\theta})\mathbf{v}) e^{-\pi|\mathbf{v}|^2} d \mathbf{v} \\ &= 1 . \end{aligned}$$

Thus we find

$$\begin{aligned} \sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \sum_{-K \leq \ell_1, \dots, \ell_k \leq K} \prod_{j=1}^k \omega_{\ell_j} \sum_{\sigma \subset \{1, \dots, M\}} \prod_{i \in \sigma^c} \gamma_{k,i}(\underline{\alpha}, \underline{\theta})^2 \prod_{j \in \sigma} (1 - \gamma_{k,j}(\underline{\alpha}, \underline{\theta})^2) \times \\ \times \int_{\mathbb{R}^M} h_0(\mathbf{v}) \log h_{0,U_k(\underline{\alpha}, \underline{\theta})}^\sigma(P_{\sigma^c} U_k(\underline{\alpha}, \underline{\theta})^T \mathbf{v}) e^{-\pi|\mathbf{v}|^2} d \mathbf{v} \leq C_{k,M} \mathcal{S}(h_0) . \end{aligned} \quad (24)$$

As $K \rightarrow \infty$, the left-hand side of (24) converges to the right-hand side of (17). \square

We now have all ingredients to give the proof of Theorem 2.1.

Proof of Theorem 2.1. Recall from Section 3, that

$$f(\mathbf{v}, t) = e^{-\pi|\mathbf{v}|^2} e^{-\Lambda t} \sum_{k=0}^{\infty} \frac{t^k \Lambda^k}{k!} h_k(\mathbf{v}) ,$$

and that $S(f(\cdot, t)) = \mathcal{S}(h(\cdot, t))$. Combining Theorem 4.1 and Theorem 5.1, we obtain

$$\mathcal{S}(h_k) \leq C_{k,M} \mathcal{S}(h_0) ,$$

and computing

$$e^{-\Lambda t} \sum_{k=0}^{\infty} \frac{\Lambda^k t^k}{k!} C_{k,M}$$

yields Theorem 2.1. \square

6. The sum rule. Proof of Theorem 5.5

We have to compute the matrix

$$\begin{aligned} Z &:= \sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{[-\pi, \pi]^k} \nu(d\theta_1) \cdots \nu(d\theta_k) \times \\ &\quad \times \sum_{\sigma \subset \{1, \dots, M\}} \prod_{i \in \sigma^c} \gamma_{k,i}(\underline{\alpha}, \underline{\theta})^2 \prod_{j \in \sigma} (1 - \gamma_{k,j}(\underline{\alpha}, \underline{\theta})^2) U_k(\underline{\alpha}, \underline{\theta}) P_{\sigma^c}^T P_{\sigma^c} U_k(\underline{\alpha}, \underline{\theta})^T . \end{aligned}$$

Obviously $P_{\sigma^c}^T P_{\sigma^c} = P_{\sigma^c}$ and hence

$$\begin{aligned} Z &= \sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{[-\pi, \pi]^k} \nu(d\theta_1) \cdots \nu(d\theta_k) \times \\ &\quad \times U_k(\underline{\alpha}, \underline{\theta}) \left[\sum_{\sigma \subset \{1, \dots, M\}} \prod_{i \in \sigma^c} \gamma_{k,i}(\underline{\alpha}, \underline{\theta})^2 \prod_{j \in \sigma} (1 - \gamma_{k,j}(\underline{\alpha}, \underline{\theta})^2) P_{\sigma^c} \right] U_k(\underline{\alpha}, \underline{\theta})^T . \end{aligned}$$

The sum on σ is easily evaluated and yields the matrix $\Gamma_k^2(\underline{\alpha}, \underline{\theta})$. Hence, recalling the orthogonal singular value decomposition (13) of $A_k(\underline{\alpha}, \underline{\theta})$, that is, $A_k(\underline{\alpha}, \underline{\theta}) = U_k(\underline{\alpha}, \underline{\theta})\Gamma_k(\underline{\alpha}, \underline{\theta})V_k^T(\underline{\alpha}, \underline{\theta})$, we find that

$$Z = \sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{[-\pi, \pi]^k} \nu(d\theta_1) \cdots \nu(d\theta_k) A_k(\underline{\alpha}, \underline{\theta}) A_k^T(\underline{\alpha}, \underline{\theta}) . \quad (25)$$

One can think about this expression in the following fashion. Recall that

$$\left[\prod_{l=1}^k r_{\alpha_l}(\theta_l) \right]^{-1} = \begin{pmatrix} A_k(\underline{\alpha}, \underline{\theta}) & B_k(\underline{\alpha}, \underline{\theta}) \\ C_k(\underline{\alpha}, \underline{\theta}) & D_k(\underline{\alpha}, \underline{\theta}) \end{pmatrix} .$$

With this notation, the matrix Z equals the top left entry of the matrix

$$\sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{[-\pi, \pi]^k} \nu(d\theta_1) \cdots \nu(d\theta_k) \left[\prod_{l=1}^k r_{\alpha_l}(\theta_l) \right]^{-1} \begin{pmatrix} I_M & 0 \\ 0 & 0 \end{pmatrix} \left[\prod_{l=1}^k r_{\alpha_l}(\theta_l) \right] .$$

The computation hinges on a repeated application of the elementary identity

$$\begin{aligned} & \int_{-\pi}^{\pi} \nu(d\theta) \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} m_1 & 0 \\ 0 & m_2 \end{pmatrix} \begin{pmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{pmatrix} \\ &= \begin{pmatrix} (1 - \tilde{\nu})m_1 + \tilde{\nu}m_2 & 0 \\ 0 & (1 - \tilde{\nu})m_2 + \tilde{\nu}m_1 \end{pmatrix} , \end{aligned}$$

where $\tilde{\nu} = \int \nu(d\theta) \sin^2(\theta)$. For this to be true we just need (5). We easily check that for the rotations $r_{\alpha}(\theta)$

$$\begin{aligned} & \sum_{\alpha} \lambda_{\alpha} \int_{-\pi}^{\pi} \nu(d\theta) r_{\alpha}(\theta)^{-1} \begin{pmatrix} m_1 I_M & 0 \\ 0 & m_2 I_N \end{pmatrix} r_{\alpha}(\theta) \\ &= \frac{1}{\Lambda} \left(\frac{M\lambda_S}{2} + \frac{N\lambda_R}{2} \right) \begin{pmatrix} m_1 I_M & 0 \\ 0 & m_2 I_N \end{pmatrix} \\ & \quad + \frac{\mu}{\Lambda N} \begin{pmatrix} N(M-1) + N((1-\tilde{\nu})m_1 + \tilde{\nu}m_2) I_M & 0 \\ 0 & (N-1)M + M(\tilde{\nu}m_1 + (1-\tilde{\nu})m_2) I_N \end{pmatrix} \\ &= \begin{pmatrix} m_1 I_M & 0 \\ 0 & m_2 I_N \end{pmatrix} + \frac{\mu_{\nu}}{\Lambda N} \begin{pmatrix} N(m_2 - m_1) I_M & 0 \\ 0 & M(m_1 - m_2) I_N \end{pmatrix} . \end{aligned} \quad (26)$$

where $\mu_{\nu} = \tilde{\nu}\mu$. Denote by $L(\nu_1, \nu_2)$ the $(N+M) \times (N+M)$ matrix

$$L(m_1, m_2) = \begin{pmatrix} m_1 I_M & 0 \\ 0 & m_2 I_N \end{pmatrix} ,$$

and set

$$\mathcal{P} = I_2 - \frac{\mu_{\nu}}{\Lambda N} \begin{pmatrix} N & -N \\ -M & M \end{pmatrix} .$$

Then (26) is recast as

$$\sum_{\alpha} \lambda_{\alpha} \int_{-\pi}^{\pi} \nu(d\theta) r_{\alpha}(\theta)^{-1} L(m_1, m_2) r_{\alpha}(\theta) = L(m'_1, m'_2) , \quad (27)$$

where

$$\begin{pmatrix} m'_1 \\ m'_2 \end{pmatrix} = \mathcal{P} \begin{pmatrix} m_1 \\ m_2 \end{pmatrix} .$$

By a repeated application of (27) we obtain

$$\sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{[-\pi, \pi]^k} \nu(d\theta_1) \cdots \nu(d\theta_k) \left[\prod_{j=1}^k r_{\alpha_j}(\theta_j) \right]^T L(\underline{m}) \left[\prod_{j=1}^k r_{\alpha_j}(\theta_j) \right] = L(\mathcal{P}^k \underline{m}) .$$

Thus,

$$Z = \left(\mathcal{P}^k \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right)_1 I_M .$$

It is easy to see that \mathcal{P} has eigenvalues $\ell_1 = 1$ and $\ell_2 = 1 - \mu_\nu(M + N)/(\Lambda N)$ with eigenvectors $\underline{m}_1 = (1, 1)$ and $\underline{m}_2 = (N, -M)^T/(M + N)$. Consequently,

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} = \frac{M}{N + M} \underline{m}_1 + \underline{m}_2 ,$$

which yields

$$\left(\mathcal{P}^k \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right)_1 = \frac{M}{N + M} + \frac{N}{M + N} \left(1 - \mu_\nu \frac{M + N}{\Lambda N} \right)^k .$$

This proves Theorem 5.5. □

7. Boltzmann-Kac collisions

In this section we show that the above results can also be extended, at least in a particular case, to three-dimensional Boltzmann-Kac collisions.

Again we consider a system of M particles coupled to a reservoir consisting of N particles, but now with velocities $v_1, \dots, v_M, w_1, \dots, w_N \in \mathbb{R}^3$. The collisions between a pair of particles have to conserve energy and momentum,

$$\begin{aligned} z_i^2 + z_j^2 &= (z_i^*)^2 + (z_j^*)^2 \\ z_i + z_j &= z_i^* + z_j^* , \end{aligned}$$

where z can be either the velocity of a system particle v or of a reservoir particle w . A convenient parametrization of the post-collisional velocities in terms of the velocities before the collision is given by

$$\begin{aligned} z_i^*(\omega) &= z_i - \omega \cdot (z_i - z_j) \omega \\ z_j^*(\omega) &= z_j + \omega \cdot (z_i - z_j) \omega, \quad \text{where } \omega \in \mathbb{S}^2 . \end{aligned}$$

This is the so-called ω -representation. This representation is particularly useful, because the velocities are related to each other by a *linear* transformation, and the strategy used to proof the results for the one-dimensional Kac system carries over rather directly. The direction ω will be chosen according to the uniform probability distribution on the unit sphere \mathbb{S}^2 .

Introduce the operators

$$(R_{ij}f)(\mathbf{z}) = \int_{\mathbb{S}^2} f(r_{ij}(\omega)^{-1}\mathbf{z}) d\omega ,$$

where $d\omega$ denotes the uniform probability measure on the sphere and the matrices $r_{ij}(\omega)$ are symmetric involutions acting as

$$\begin{pmatrix} z_i^* \\ z_j^* \end{pmatrix} = \begin{pmatrix} I - \omega\omega^T & \omega\omega^T \\ \omega\omega^T & I - \omega\omega^T \end{pmatrix} \begin{pmatrix} z_i \\ z_j \end{pmatrix}$$

on the velocities of the particles i and j , and as identities otherwise. They will replace the one-dimensional Kac collision operators in (3) in the otherwise unchanged generator of the time evolution. Notice that the matrices $r_{ij}(\omega)$ are orthogonal, so that the expansion formula (10) still holds with the obvious changes in the dimension of the single-particle spaces.

We prove an analog of Theorem 2.1 for the case of three-dimensional Boltzmann-Kac collisions and pseudo-Maxwellian molecules.

Theorem 7.1. *Let $N \geq M$ and $F_0(\mathbf{v}, \mathbf{w}) = f_0(\mathbf{v}) e^{-\pi|\mathbf{w}|^2}$ for some probability distribution f_0 on \mathbb{R}^{3M} . Then the entropy of the marginal*

$$f(\mathbf{v}, t) := \int_{\mathbb{R}^{3N}} (e^{\mathcal{L}t} F_0)(\mathbf{v}, \mathbf{w}) d\mathbf{w}$$

with respect to the thermal state $e^{-\pi|\mathbf{v}|^2}$ is bounded by

$$S(f(\cdot, t)) \leq \left[\frac{N}{N+M} + \frac{N}{N+M} e^{-\frac{\mu}{3} \frac{N+M}{N} t} \right] S(f_0) .$$

Remark 7.2. *The result in three dimensions is very similar to the case of one-dimensional Kac collisions, with the difference that the rate of exponential decay is $\mu/3$ instead of μ_ρ . The factor $1/3$ comes from the fact that $\int_{\mathbb{S}^2} d\omega \omega\omega^T = I_3/3$. It would be interesting to cover the true Maxwellian molecules interaction*

$$(R_{ij}f)(z) = \int_{\mathbb{S}^2} b \left(\frac{v_i - v_j}{|v_i - v_j|} \cdot \omega \right) f(r_{ij}(\omega)^{-1}z) d\omega .$$

However, the dependence of the scattering rate b on the velocities doesn't seem to be treatable with the above methods.

The proof of Theorem 7.1 essentially deviates from the one-dimensional case in only two places: the sum rule and the discrete approximation of the integrals. We begin by proving an analogue of Theorem 5.5. Most of the steps for the computation of the matrix Z in (25) are the same. What remains is to compute

$$Z := \sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{\mathbb{S}^2 \times \dots \times \mathbb{S}^2} d\omega_1 \cdots d\omega_k A_k(\underline{\alpha}, \underline{\omega}) A_k(\underline{\alpha}, \underline{\omega})^T ,$$

which is somewhat different for the case of Boltzmann-Kac collisions. Recall that $A_k(\underline{\alpha}, \underline{\omega})$ is the upper left $3M \times 3M$ block of $[\prod_{j=1}^k r_{\alpha_j}(\omega_j)]^{-1}$, i.e.,

$$A_k(\underline{\alpha}, \underline{\omega}) = P_{3M} [\prod_{j=1}^k r_{\alpha_j}(\omega_j)]^{-1} P_{3M}^T$$

with the projection $P_{3M} = (I_{3M} \ 0)$ from $\mathbb{R}^{3M+3N} \rightarrow \mathbb{R}^{3M}$. In particular, by linearity,

$$Z = P_{3M} \left(\sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{(\mathbb{S}^2)^k} d\underline{\omega} \left[\prod_{j=1}^k r_{\alpha_j}(\omega_j) \right]^{-1} \begin{pmatrix} I_{3M} & 0 \\ 0 & 0 \end{pmatrix} \left[\prod_{j=1}^k r_{\alpha_j}(\omega_j) \right] \right) P_{3M}^T .$$

As in the proof of Theorem 5.5 we have

Lemma 7.3. *Let $\alpha, \beta \geq 0$. Then*

$$\sum_{1 \leq i < j \leq M+N} \lambda_{ij} \int_{\mathbb{S}^2} d\omega r_{ij}(\omega)^{-1} \begin{pmatrix} \alpha I_{3M} & 0 \\ 0 & \beta I_{3N} \end{pmatrix} r_{ij}(\omega) = \begin{pmatrix} \alpha' I_{3M} & 0 \\ 0 & \beta' I_{3N} \end{pmatrix},$$

where α', β' are related to α, β by

$$\begin{pmatrix} \alpha' \\ \beta' \end{pmatrix} = \mathcal{P} \begin{pmatrix} \alpha \\ \beta \end{pmatrix}, \quad \mathcal{P} = I_2 - \frac{\mu}{3\Lambda} \begin{pmatrix} 1 & -1 \\ -\frac{M}{N} & \frac{M}{N} \end{pmatrix}.$$

Notice that the matrix \mathcal{P} of Lemma 7.3 has eigenvalues 1 and $1 - \mu/(3\Lambda)(1 + M/N)$ with corresponding eigenvectors $(1 \ 1)^T$ and $(-N/M \ 1)^T$. Repeated application of Lemma 7.3 then implies, see also the argument in the one-dimensional case,

$$\sum_{\alpha_1, \dots, \alpha_k} \lambda_{\alpha_1} \cdots \lambda_{\alpha_k} \int_{(\mathbb{S}^2)^k} d\omega \left[\prod_{j=1}^k r_{\alpha_j}(\omega_j) \right]^{-1} \begin{pmatrix} \alpha I_{3M} & 0 \\ 0 & \beta I_{3N} \end{pmatrix} \left[\prod_{j=1}^k r_{\alpha_j}(\omega_j) \right] = \begin{pmatrix} \alpha^{(k)} I_{3M} & 0 \\ 0 & \beta^{(k)} I_{3N} \end{pmatrix},$$

where

$$\begin{pmatrix} \alpha^{(k)} \\ \beta^{(k)} \end{pmatrix} = \mathcal{P}^k \begin{pmatrix} \alpha \\ \beta \end{pmatrix}.$$

Before we prove Lemma 7.3, let us make an easy observation.

Corollary 7.4. *In the particular case $\alpha = 1, \beta = 0$, we get*

$$Z = \left[\frac{M}{M+N} + \frac{N}{M+N} \left(1 - \frac{\mu}{3\Lambda} \left(1 + \frac{M}{N} \right) \right)^k \right] I_{3M}.$$

Proof of Lemma 7.3. For $1 \leq i < j \leq M$ (respectively for $M+1 \leq i < j \leq M+N$) the operators $r_{ij}(\omega)$ only act non-trivially in the first $3M$ (last $3N$) variables. Taking into account that $r_{ij}(\omega)^{-1} I r_{ij}(\omega) = I$, we obtain

$$\frac{\lambda_S}{M-1} \sum_{1 \leq i < j \leq M} \int_{\mathbb{S}^2} d\omega r_{ij}(\omega)^{-1} \begin{pmatrix} \alpha I_{3M} & 0 \\ 0 & \beta I_{3N} \end{pmatrix} r_{ij}(\omega) = \frac{M\lambda_S}{2} \begin{pmatrix} \alpha I_{3M} & 0 \\ 0 & \beta I_{3N} \end{pmatrix},$$

and

$$\frac{\lambda_R}{N-1} \sum_{M+1 \leq i < j \leq M+N} \int_{\mathbb{S}^2} d\omega r_{ij}(\omega)^{-1} \begin{pmatrix} \alpha I_{3M} & 0 \\ 0 & \beta I_{3N} \end{pmatrix} r_{ij}(\omega) = \frac{N\lambda_R}{2} \begin{pmatrix} \alpha I_{3M} & 0 \\ 0 & \beta I_{3N} \end{pmatrix}.$$

It remains to look at the interaction terms $i = 1, \dots, M$ and $j = M+1, \dots, M+N$. Notice that

$$\begin{aligned} & r_{ij}(\omega)^{-1} \begin{pmatrix} \alpha I_{3M} & 0 \\ 0 & \beta I_{3N} \end{pmatrix} r_{ij}(\omega) \\ &= \begin{pmatrix} \alpha I_{3M} & 0 \\ 0 & \beta I_{3N} \end{pmatrix} + \left(\begin{array}{c|c} 0 & 0 \\ \hline (\beta - \alpha)\omega\omega^T & 0 \\ \hline 0 & 0 \\ \hline 0 & (\beta - \alpha)\omega\omega^T \\ & 0 \end{array} \right), \end{aligned}$$

where the non-zero entries in the second summand on the right-hand side correspond to the i^{th} , respectively j^{th} , 3×3 block on the diagonal. Since $\int_{\mathbb{S}^2} d\omega \omega \omega^T = 1/3 I_3$, we obtain

$$\begin{aligned} & \frac{\mu}{N} \sum_{i=1}^M \sum_{j=M+1}^{M+N} \int_{\mathbb{S}^2} d\omega r_{ij}(\omega)^{-1} \begin{pmatrix} \alpha I_{3M} & 0 \\ 0 & \beta I_{3N} \end{pmatrix} r_{ij}(\omega) \\ &= \mu M \begin{pmatrix} \alpha I_{3M} & 0 \\ 0 & \beta I_{3N} \end{pmatrix} + \frac{\mu}{3} (\alpha - \beta) \begin{pmatrix} -I_{3M} & 0 \\ 0 & \frac{M}{N} I_{3N} \end{pmatrix}. \end{aligned}$$

Recall the definition of $\Lambda = M\lambda_S/2 + N\lambda_R/2 + \mu M$. Hence summation of all the three contributions yields the statement of the Lemma. \square

As in the one-dimensional case, in order to apply the geometric Brascamp-Lieb inequality Theorem 5.2, we need to approximate the uniform probability measure $d\omega$ on the sphere by a suitable sequence of discrete measures as in the one-dimensional case (see Lemma 5.4). Additionally, in each step of the discretization, the constraint $\int_{\mathbb{S}^2} d\omega \omega \omega^T = 1/3 I$, has to hold. This is important, because it guarantees that the geometric Brascamp-Lieb condition, i.e., the sum rule (19), holds in each step.

In order to find such an approximation, we parametrize the sphere in the usual way by spherical coordinates

$$\omega = \omega(\theta, \varphi) = \begin{pmatrix} \sin \theta \cos \varphi \\ \sin \theta \sin \varphi \\ \cos \theta \end{pmatrix}$$

for $\theta \in [0, \pi]$ and $\varphi \in [0, 2\pi]$. For $K, L \in \mathbb{N}$ we introduce the measures

$$\begin{aligned} \Phi_K &:= \frac{\pi}{K} \sum_{j=0}^{2K-1} \delta_{\frac{\pi}{K}j} \quad \text{on } [0, 2\pi], \quad \text{and} \\ \Theta_L &:= \sum_{i=1}^L \frac{2}{(1-u_i^2)^{3/2} (P'_L(u_i))^2} \delta_{\arccos u_i} \quad \text{on } [0, \pi], \end{aligned}$$

where P_L is the Legendre polynomial of order L on $[-1, 1]$, and u_i , $i = 1, \dots, L$, are its zeros. Then, if $f \in \mathcal{C}[0, 2\pi]$ and $g \in \mathcal{C}[-1, 1]$,

$$\int_0^{2\pi} f(\varphi) \Phi_k(d\varphi) = \frac{\pi}{K} \sum_{j=0}^{2K-1} f\left(\frac{\pi}{K}j\right) \rightarrow \int_0^{2\pi} f(\varphi) d\varphi$$

as $K \rightarrow \infty$ as Riemann sum. Furthermore,

$$\int_0^\pi g(\cos \theta) \sin \theta \Theta_L(d\theta) = \sum_{i=1}^L \frac{2}{(1-u_i^2)(P'_L(u_i))^2} g(u_i) \rightarrow \int_{-1}^1 g(u) du = \int_0^\pi g(\cos \theta) \sin \theta d\theta$$

as $L \rightarrow \infty$ by Gauss-Legendre quadrature. The latter approximation is exact for polynomials of order less or equal to $2L - 1$. In particular, we have

$$\int_0^\pi \cos^2 \theta \sin \theta \Theta_L(d\theta) = \int_0^\pi \cos^2 \theta \sin \theta d\theta = \frac{2}{3}, \quad \text{and} \quad \int_0^\pi \sin^3 \theta \Theta_L(d\theta) = \int_0^\pi \sin^3 \theta d\theta = \frac{4}{3},$$

for all $L \geq 2$. It is easy to check that

$$\begin{aligned} \int_0^{2\pi} \sin \varphi \cos \varphi \Phi_k(d\varphi) &= 0, \\ \int_0^{2\pi} \sin \varphi \Phi_k(d\varphi) &= \int_0^{2\pi} \cos \varphi \Phi_k(d\varphi) = 0, \\ \int_0^{2\pi} \sin^2 \varphi \Phi_k(d\varphi) &= \int_0^{2\pi} \cos^2 \varphi \Phi_k(d\varphi) = \pi, \end{aligned}$$

for all $K \geq 2$. Consequently,

$$\begin{aligned} & \frac{1}{4\pi} \int_0^{2\pi} \omega(\theta, \varphi) \omega(\theta, \varphi)^T \Theta_L(d\theta) \Phi_k(d\varphi) \\ &= \frac{1}{2K} \sum_{j=0}^{2K-1} \sum_{i=0}^L \frac{\omega(\arccos u_i, \pi j/K) \omega(\arccos u_i, \pi j/K)^T}{(1-u_i^2)(P'_L(u_i))^2} = \frac{1}{3} I_3 \end{aligned}$$

for all $K, L \geq 2$. It follows that Z is not changed by replacing the uniform measure on \mathbb{S}^2 by the above discrete approximation, in particular, Z is still proportional to the identity matrix, which guarantees the applicability of the geometric Brascamp-Lieb inequality.

This concludes the proof of Theorem 7.1. \square

APPENDIX A. The Geometric Brascamp-Lieb inequality and the entropy inequality

In this section we prove Theorem 5.2. We use the same strategy as in [13] and [5] which consists of transporting the functions f_i with the heat kernel in such a way that the right-hand side of (20) remains fixed while the left-hand side of that inequality increases. The results in [5] are quite general but for the special case in which the sum rule (19) holds, the proof is quite simple and this is one of the reasons why we include it here.

Proof of Theorem 5.2. The inequality (20) is equivalent to

$$\int_{\mathbb{R}^M} \prod_{i=1}^K f_i^{c_i}(B_i \mathbf{v}) d\mathbf{v} \leq \prod_{i=1}^K \left(\int_{H_i} f_i(u) du \right)^{c_i}. \quad (28)$$

This follows from the identity

$$\prod_{i=1}^K \left(e^{-\pi |B_i \mathbf{v}|^2} \right)^{c_i} = e^{-\pi \sum_{i=1}^K (\mathbf{v} \cdot c_i B_i^T B_i \mathbf{v})} = e^{-\pi |\mathbf{v}|^2}.$$

We transport the functions f_i by the heat flow, that is we define

$$f_i(B_i \mathbf{v}, t) := \frac{1}{(4\pi t)^{M/2}} \int_{\mathbb{R}^M} e^{-|\mathbf{v}-\mathbf{w}|^2/(4t)} f_i(B_i \mathbf{w}) d\mathbf{w}. \quad (29)$$

For the above definition to make sense, we have to show that the right-hand side is a function of $B_i \mathbf{v}$ alone. The condition $B_i B_i^T = I_{H_i}$ means that the matrix $P_i = B_i^T B_i$ is an orthogonal projection onto a d_i dimensional subspace of \mathbb{R}^M . Moreover, $B_i P_i = I_{H_i} B_i = B_i$. We rewrite the integral (29) by splitting it in an integral over $\mathbf{w}' \in \text{Ran } P_i$ and one over integration over

$\mathbf{w}'' \in \text{Ran } P_i^\perp$. Carrying out the integration over \mathbf{w}'' we obtain

$$\begin{aligned}
f_i(B_i \mathbf{v}, t) &= \frac{1}{(4\pi t)^{M/2}} \int_{\text{Ran } P_i} \int_{\text{Ran } P_i^\perp} e^{-|(P_i \mathbf{v} - P_i \mathbf{w}')|^2/(4t)} e^{-|(P_i^\perp \mathbf{v} - \mathbf{w}'')|^2/(4t)} f_i(B_i P_i \mathbf{w}') d\mathbf{w}' d\mathbf{w}'' \\
&= \frac{1}{(4\pi t)^{d_i/2}} \int_{\text{Ran } P_i} e^{-|(P_i \mathbf{v} - P_i \mathbf{w}')|^2/(4t)} f_i(B_i P_i \mathbf{w}') d\mathbf{w}' \\
&= \frac{1}{(4\pi t)^{d_i/2}} \int_{\text{Ran } P_i} e^{-|(B_i \mathbf{v} - B_i \mathbf{w}')|^2/(4t)} f_i(B_i \mathbf{w}') d\mathbf{w}' \\
&= \frac{1}{(4\pi t)^{d_i/2}} \int_{H_i} e^{-|(B_i \mathbf{v} - u)|^2/(4t)} f_i(u) du .
\end{aligned}$$

where, in the last equality, we have used that B_i maps the range of P_i isometrically onto H_i . This justifies (29). Moreover, the above computation also shows that

$$\int_{H_i} f_i(u, t) du = \int_{H_i} f_i(u) du$$

so that the right-hand side of the inequality (28) does not change under the heat flow.

We now show that the left-hand side of (28) is an increasing function of t . It is convenient to set $\phi_i(u, t) = \log f_i(u, t)$. Differentiating the function $\phi_i(B_i \mathbf{v}, t)$ with respect to t yields

$$\frac{d}{dt} \phi_i(B_i \mathbf{v}, t) = \Delta_{\mathbf{v}} \phi_i(B_i \mathbf{v}, t) + |\nabla_{\mathbf{v}} \phi_i(B_i \mathbf{v}, t)|^2 .$$

Moreover,

$$\frac{d}{dt} \int_{\mathbb{R}^M} \prod_{i=1}^K f_i^{c_i}(B_i \mathbf{v}, t) d\mathbf{v} = \sum_{m=1}^K c_m \int_{\mathbb{R}^M} [\Delta_{\mathbf{v}} \phi_m(B_m \mathbf{v}, t) + |\nabla_{\mathbf{v}} \phi_m(B_m \mathbf{v}, t)|^2] \prod_{i=1}^K f_i^{c_i}(B_i \mathbf{v}, t) d\mathbf{v} .$$

Integrating by parts the term containing the Laplacian yields

$$\begin{aligned}
\frac{d}{dt} \int_{\mathbb{R}^M} \prod_{i=1}^K f_i^{c_i}(B_i \mathbf{v}, t) d\mathbf{v} &= \\
&\sum_{m=1}^K c_m \int_{\mathbb{R}^M} |\nabla_{\mathbf{v}} \phi_m(B_m \mathbf{v}, t)|^2 \prod_{i=1}^K f_i^{c_i}(B_i \mathbf{v}, t) d\mathbf{v} \\
&\quad - \sum_{m, \ell=1}^K c_m c_\ell \int_{\mathbb{R}^M} \nabla_{\mathbf{v}} \phi_m(B_m \mathbf{v}, t) \cdot \nabla_{\mathbf{v}} \phi_\ell(B_\ell \mathbf{v}, t) \prod_{i=1}^K f_i^{c_i}(B_i \mathbf{v}, t) d\mathbf{v} .
\end{aligned}$$

Finally, using that

$$\nabla_{\mathbf{v}} \phi_m(B_m \mathbf{v}, t) = B_m^T (\nabla \phi_m)(B_m \mathbf{v})$$

we get

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^M} \prod_{i=1}^K f_i^{c_i}(B_i \mathbf{v}, t) d\mathbf{v} = \\ \sum_{m=1}^K c_m \int_{\mathbb{R}^M} |B_m^T(\nabla \phi_m)(B_m \mathbf{v}, t)|^2 \prod_{i=1}^K f_i^{c_i}(B_i \mathbf{v}, t) d\mathbf{v} \\ - \sum_{m,\ell=1}^K c_m c_\ell \int_{\mathbb{R}^M} B_m^T(\nabla \phi_m)(B_m \mathbf{v}, t) \cdot B_\ell^T(\nabla \phi_\ell)(B_\ell \mathbf{v}, t) \prod_{i=1}^K f_i^{c_i}(B_i \mathbf{v}, t) d\mathbf{v} . \end{aligned}$$

We claim that this expression is non-negative. The vectors $\nabla \phi_m \in H_m$ are arbitrary and hence the problem is reduced to proving that for any set of vectors $V_m \in H_m$, $m = 1, \dots, K$, it holds

$$\sum_{m=1}^K c_m |B_m^T V_m|^2 - \sum_{m,\ell=1}^K c_m c_\ell B_m^T V_m \cdot B_\ell^T V_\ell \geq 0 .$$

Recalling that $B_m B_m^T = I_{H_m}$ and setting $Y = \sum_{\ell} c_\ell B_\ell^T V_\ell$ we conclude that it is enough to show that

$$|Y|^2 \leq \sum_{m=1}^K c_m |V_m|^2 .$$

This follows easily, since, by applying Schwarz's inequality, we find that

$$|Y|^2 = \sum_{\ell=1}^K c_\ell Y \cdot B_\ell^T V_\ell = \sum_{\ell=1}^K c_\ell B_\ell Y \cdot V_\ell \leq \left(\sum_{\ell=1}^K c_\ell |B_\ell Y|^2 \right)^{1/2} \left(\sum_{\ell=1}^K c_\ell |V_\ell|^2 \right)^{1/2} .$$

Combining this with (19), we learn that

$$|Y|^2 \leq \left(Y \cdot \sum_{\ell=1}^K c_\ell B_\ell^T B_\ell Y \right)^{1/2} \left(\sum_{\ell=1}^K c_\ell |V_\ell|^2 \right)^{1/2} = |Y| \left(\sum_{\ell=1}^K c_\ell |V_\ell|^2 \right)^{1/2} .$$

Thus we have that, when applying (28) to the functions $f_i(u, t)$, the left hand side is an increasing function of t while the right hand side does not depends on t . It is thus enough to show that the inequality holds for large t . Using once more the sum-rule (19), we see that

$$\begin{aligned} \int_{\mathbb{R}^M} \prod_{i=1}^K \frac{1}{(4\pi t)^{c_i d_i/2}} \left[\int_{H_i} e^{-\frac{|B_i \mathbf{v} - u|^2}{4t}} f_i(u) du \right]^{c_i} d\mathbf{v} = \\ \frac{1}{(4\pi)^{M/2}} \int_{\mathbb{R}^M} \prod_{i=1}^K \left[\int_{H_i} e^{-\frac{|B_i \mathbf{v} - t^{-1/2} u|^2}{4}} f_i(u) du \right]^{c_i} d\mathbf{v} \xrightarrow{t \rightarrow \infty} \frac{1}{(4\pi)^{M/2}} \int_{\mathbb{R}^M} e^{-\frac{|\mathbf{v}|^2}{4}} \prod_{i=1}^K \left[\int_{H_i} f_i(u) du \right]^{c_i} d\mathbf{v} \\ = \prod_{i=1}^K \left[\int_{H_i} f_i(u) du \right]^{c_i} \end{aligned}$$

which proves the first part of Theorem 5.2.

To prove the entropy inequality (21) we follow [11]. Let h be a non-negative function whose L^1 norm is one and whose entropy is finite. An elementary computation then shows that

$$\int_{\mathbb{R}^M} h(\mathbf{v}) \log h(\mathbf{v}) e^{-\pi|\mathbf{v}|^2} d\mathbf{v} = \sup_{\Phi} \left\{ \int_{\mathbb{R}^M} h(\mathbf{v}) \Phi(\mathbf{v}) e^{-\pi|\mathbf{v}|^2} d\mathbf{v} - \log \int_{\mathbb{R}^M} e^{\Phi(\mathbf{v})} e^{-\pi|\mathbf{v}|^2} d\mathbf{v} \right\} .$$

Now, we set

$$\Phi(\mathbf{v}) = \sum_{i=1}^K c_i \log f_i(B_i \mathbf{v}) .$$

This leads to the lower bound

$$\begin{aligned} & \int_{\mathbb{R}^M} h(\mathbf{v}) \log h(\mathbf{v}) e^{-\pi|\mathbf{v}|^2} d\mathbf{v} \\ & \geq \sum_{i=1}^K c_i \int_{\mathbb{R}^M} h(\mathbf{v}) \log f_i(B_i \mathbf{v}) e^{-\pi|\mathbf{v}|^2} d\mathbf{v} - \log \int_{\mathbb{R}^M} \prod_{i=1}^K f_i(B_i \mathbf{v})^{c_i} e^{-\pi|\mathbf{v}|^2} d\mathbf{v} \\ & \geq \sum_{i=1}^K c_i \int_{\mathbb{R}^M} h(\mathbf{v}) \log f_i(B_i \mathbf{v}) e^{-\pi|\mathbf{v}|^2} d\mathbf{v} - \log \left[\prod_{i=1}^K \left(\int_{H_i} f_i(u) e^{-\pi|u|^2} du \right)^{c_i} \right] , \end{aligned}$$

where the second step is a consequence of the Brascamp-Lieb inequality (20). \square

APPENDIX B. Proof of Lemma 5.4

Proof. For K any positive integer we convolve $\rho(\theta)$ with the non-negative trigonometric polynomial

$$p_K(\theta) := \frac{1}{2K+1} \left(\sum_{k=-K}^K e^{ik\theta} \right)^2 = \sum_{m=-2K}^{2K} \left(1 - \frac{|m|}{2K+1} \right) e^{im\theta} ,$$

and obtain a probability density $\rho_K(\theta)$. The Fourier coefficients of $\rho_K(\theta)$ are given by

$$\widehat{\rho}_K(m) = \widehat{\rho}(m) \left(1 - \frac{|m|}{2K+1} \right)$$

for $|m| \leq 2K$ and are zero otherwise. In particular,

$$\widehat{\rho}_K(2) - \widehat{\rho}_K(-2) = 4i \int_{-\pi}^{\pi} \rho_K(\theta) \sin \theta \cos \theta d\theta = 0 .$$

With ρ_K we construct the measure

$$\nu_K(d\theta) = \frac{2\pi}{4K+1} \sum_{\ell=-2K}^{2K} \rho_K \left(\frac{2\pi\ell}{4K+1} \right) \delta \left(\theta - \frac{2\pi\ell}{4K+1} \right) d\theta .$$

The measure ν_K is positive since $\rho_K((2\pi\ell)/(4K+1)) \geq 0$. Moreover, for all $m \in \mathbb{Z}$ with $|m| \leq 2K$ the Fourier coefficients $\widehat{\nu}_K(m)$ and $\widehat{\rho}_K(m)$ coincide. In particular, we have

$$\int_{-\pi}^{\pi} \nu_K(d\theta) \sin \theta \cos \theta = 0 .$$

To see this, we compute

$$\widehat{\nu}_K(m) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \nu_K(\theta) e^{-im\theta} d\theta = \frac{1}{4K+1} \sum_{\ell=-2K}^{2K} \rho_K \left(\frac{2\pi\ell}{4K+1} \right) e^{-2\pi i m \ell / (4K+1)}$$

for $|m| \leq 2K$. Observe that

$$\rho_K \left(\frac{2\pi\ell}{4K+1} \right) = \sum_{k=-2K}^{2K} \widehat{\rho}_K(k) e^{2\pi i k \ell / (4K+1)},$$

and, as a consequence,

$$\widehat{\nu}_K(m) = \frac{1}{4K+1} \sum_{\ell=-2K}^{2K} \sum_{k=-2K}^{2K} \widehat{\rho}_K(k) e^{2\pi i \ell (k-m) / (4K+1)}.$$

But

$$\sum_{\ell=-2K}^{2K} e^{2\pi i \ell (k-m) / (4K+1)} = \begin{cases} 4K+1 & \text{if } k = m \\ 0 & \text{if } k \neq m, \end{cases}$$

and hence we conclude that

$$\widehat{\nu}_K(m) = \widehat{\rho}_K(m) \tag{30}$$

for $|m| \leq 2K$. It is easy to see that for any continuous function f on $[-\pi, \pi]$,

$$\lim_{K \rightarrow \infty} \int_{-\pi}^{\pi} f(\theta) \nu_K(d\theta) = \int_{-\pi}^{\pi} f(\theta) \rho(\theta) d\theta.$$

□

REFERENCES

- [1] K. BALL, [Volumes of sections of cubes and related problems](#). In *Geometric Aspects of Functional Analysis: Israel Seminar (GAFA) 1987–88*, Volume **1376** of the series *Lecture Notes in Mathematics*, pp 251–260. Springer Berlin Heidelberg, 1989. [MR 1008726](#). [Zbl 0674.46008](#).
- [2] K. BALL, Volume ratios and a reverse isoperimetric inequality. *Journal of the London Mathematical Society (Second Series)* **44** (1991), 351–359. [MR 1136445](#). [Zbl 0694.46010](#).
- [3] F. BARTHE, On a reverse form of the Brascamp-Lieb inequality. *Inventiones mathematicae* **134** (1998), 335–361. [MR 1650312](#). [Zbl 0901.26010](#).
- [4] F. BARTHE, [A continuous version of the Brascamp-Lieb inequalities](#). In *Geometric Aspects of Functional Analysis: Israel Seminar 2002–2003*, Volume **1850** of the series *Lecture Notes in Mathematics*, pp 53–63. Springer Berlin Heidelberg, 2004. [MR 2087150](#). [Zbl 1084.26011](#).
- [5] J. BENNETT, A. CARBERY, M. CHRIST, and T. TAO, The Brascamp-Lieb inequalities: finiteness, structure and extremals. *Geometric and Functional Analysis* **17** (2008), 1343–1415. [MR 2377493](#). [Zbl 1132.26006](#).
- [6] F. BONETTO, M. LOSS, H. TOSSOUNIAN, and R. VAIDYANATHAN, Uniform approximation of a Maxwellian thermostat by finite reservoirs. *Communications in Mathematical Physics* **351** (2017), 311–339. [MR 3613506](#). [Zbl 06702028](#).
- [7] F. BONETTO, M. LOSS, and R. VAIDYANATHAN, The Kac model coupled to a thermostat. *Journal of Statistical Physics* **156** (2014), 647–667. [MR 3226836](#). [Zbl 1302.35290](#).
- [8] H.J. BRASCAMP, and E.H. LIEB, Best constants in Young’s inequality, its converse, and its generalization to more than three functions. *Advances in Mathematics* **20**(1976), 151–173. [MR 0412366](#). [Zbl 0339.26020](#).
- [9] E.A. CARLEN, M.C. CARVALHO, and M. LOSS, [Many-body aspects of approach to equilibrium](#). In *Journées “Équations aux Dérivées Partielles” (La Chapelle sur Erdre, 2000)*, Exp. No. XI, 12 pp., Université de Nantes, Nantes, 2000. [MR 1775687](#). [Zbl 1213.81230](#).

- [10] E.A. CARLEN, M.C. CARVALHO, and M. LOSS, Determination of the spectral gap for Kac's master equation and related stochastic evolution. *Acta Mathematica* **191** (2003), 1–54. MR 2020418. Zbl 1080.60091.
- [11] E.A. CARLEN, and D. CORDERO-ERAUSQUIN, Subadditivity of the entropy and its relation to Brascamp-Lieb type inequalities. *Geometric and Functional Analysis* **19** (2009), 373–405. MR 2545242. Zbl 1231.26015.
- [12] E.A. CARLEN, and M. LOSS, Extremals of functionals with competing symmetries. *Journal of Functional Analysis* **88** (1990), 437–456. MR 1038450. Zbl 0705.46016.
- [13] E.A. CARLEN, E.H. LIEB, and M. LOSS, A sharp analog of Young's inequality on S^N and related entropy inequalities. *The Journal of Geometric Analysis* **14** (2004), 487–520. MR 2077162. Zbl 1056.43002.
- [14] A. EINAV, On Villani's conjecture concerning entropy production for the Kac master equation. *Kinetic and Related Models* **4** (2011), 479–497. MR 2786394. Zbl 1335.82023.
- [15] P. FEDERBUSH, Partially alternate derivation of a result of Nelson. *Journal of Mathematical Physics* **10** (1969), 50–52. Zbl 0165.58301.
- [16] E. GABETTA, G. TOSCANI, and B. WENNERBERG, Metrics for probability distributions and the trend to equilibrium for solutions of the Boltzmann equation. *Journal of Statistical Physics* **81** (1995), 901–934. MR 1361302. Zbl 1081.82616.
- [17] L. GROSS, Logarithmic Sobolev inequalities. *American Journal of Mathematics* **97** (1975), 1061–1083. MR 0420249. Zbl 0318.46049.
- [18] L. GROSS, Logarithmic Sobolev inequalities and contractivity properties of semigroups. In *Dirichlet Forms (Varenna, 1992)*, Volume **1563** of the series *Lecture Notes in Mathematics*, pp 54–88. Springer Berlin Heidelberg, 1993. MR 1292277. Zbl 0812.47037.
- [19] T.S. HAN, Nonnegative entropy measures of multivariate symmetric correlations. *Information and Control* **36** (1978), 133–156. MR 0464499. Zbl 0367.94041.
- [20] E. JANVRESSE, Spectral gap for Kac's model of Boltzmann equation. *The Annals of Probability* **29** (2001), 288–304. MR 1825150. Zbl 1034.82049.
- [21] M. KAC, Foundations of kinetic theory. In *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, 1954–1955, vol. III*, pp 171–197. University of California Press, Berkeley and Los Angeles, 1956. MR 0084985. Zbl 0072.42802.
- [22] E.H. LIEB, Gaussian kernels have only Gaussian maximizers. *Inventiones mathematicae* **102** (1990), 179–208. MR 1069246. Zbl 0726.42005.
- [23] L.H. LOOMIS, and H. WHITNEY, An inequality related to the isoperimetric inequality. *Bulletin of the American Mathematical Society* **55** (1949), 961–962. MR 0031538. Zbl 0035.38302.
- [24] D.K. MASLEN, The eigenvalues of Kac's master equation. *Mathematische Zeitschrift* **243** (2003), 291–331. MR 1961868. Zbl 1016.82016.
- [25] S. MISCHLER, and C. MOUHOT, About Kac's program in kinetic theory. *Comptes Rendus Mathématique (Académie des Sciences Paris)* **349** (2011), 1245–1250. MR 2861993. Zbl 1233.35153.
- [26] S. MISCHLER, and C. MOUHOT, Kac's program in kinetic theory. *Inventiones mathematicae* **193** (2013), 1–147. MR 3069113. Zbl 1274.82048.
- [27] E. NELSON, The free Markoff field. *Journal of Functional Analysis* **12** (1973), 211–227. MR 0343816. Zbl 0273.60079.
- [28] H. TOSSOUNIAN, and R. VAIDYANATHAN, Partially thermostated Kac model. *Journal of Mathematical Physics* **56** (2015), 083301 [16 pages]. MR 3455336. Zbl 1330.82036.
- [29] C. VILLANI, Cercignani's conjecture is sometimes true and always almost true. *Communications in Mathematical Physics* **234** (2003), 455–490. MR 1964379. Zbl 1041.82018.

E-mail address: bonetto@math.gatech.edu

E-mail address: alissa.geisinger@uni-tuebingen.de

E-mail address: loss@math.gatech.edu

E-mail address: tobias.ried@kit.edu