

Unified error analysis for non-conforming space discretizations of wave-type equa- tions

David Hipp, Marlis Hochbruck, Christian Stohrer

CRC Preprint 2017/29 (revised), April 2018

KARLSRUHE INSTITUTE OF TECHNOLOGY

CRC 1173



Wave
phenomena

Participating universities



Universität Stuttgart

EBERHARD KARLS
UNIVERSITÄT
TÜBINGEN



Funded by

DFG

ISSN 2365-662X

Unified error analysis for non-conforming space discretizations of wave-type equations

DAVID HIPPT[†], MARLIS HOCHBRUCK[‡], CHRISTIAN STOHRER[§]

*Karlsruhe Institute of Technology
Institute for Applied and Numerical Mathematics
Englerstr. 2, 76131 Karlsruhe, Germany*

[Received on 6 June 2018]

This paper provides a unified error analysis for non-conforming space discretizations of linear wave equations in time-domain. We propose a framework which studies wave equations as first-order evolution equations in Hilbert spaces and their space discretizations as differential equations in finite dimensional Hilbert spaces. A lift operator maps the semi-discrete solution from the approximation space to the continuous space. Our main results are a priori error bounds in terms of interpolation, data and conformity errors of the method. Such error bounds are the key to the systematic derivation of convergence rates for a large class of problems. To show that this approach significantly eases the proof of new convergence rates, we apply it to an isoparametric finite element discretization of the wave equation with acoustic boundary conditions in a smooth domain. Moreover, our results reproduce known convergence rates for already investigated conforming and non-conforming space discretizations in a concise and unified way. The examples discussed in this paper comprise discontinuous Galerkin discretizations of Maxwell's equations and finite elements with mass lumping for the acoustic wave equation.

This article has been accepted for publication in IMA Journal of Numerical Analysis published by Oxford University Press.

Keywords: wave equation, non-conforming space discretization, abstract error analysis, a priori error bounds, linear evolution equations, operator semigroups, linear monotone operators in Hilbert spaces, dynamic boundary conditions, isoparametric finite elements

1. Introduction

In this paper we investigate the convergence behavior of possibly non-conforming space discretizations of wave equations written as first-order or as second-order partial differential equations in time and space. Both types of equations arise as mathematical models for wave phenomena in elastodynamics, electromagnetics and acoustics, cf. Joly (2003). In the last few decades, there has been a remarkable progress in understanding and analyzing such numerical approximations. Despite sharing the main ideas of proof, most contributions focus on a particular wave-type equation and a particular space discretization method, cf. e.g. Baker (1976), Zhao (2004), Cohen & Pernet (2017).


Only a few papers proceed in a unified way and harness the analogies shared between the individual studies. Among them are several works which develop a unified error analysis for a particular class of space discretization methods for wave equations, cf. Fujita *et al.* (2001), Joly (2003), Burman *et al.*

[†]Corresponding author. Email:david.hipp@kit.edu

[‡]Email: marlis.hochbruck@kit.edu

[§]Email: christian.stohrer@kit.edu

(2010). On the other hand, abstract approximation theory for evolution equations mostly shows convergence but does not provide error bounds or convergence rates, e.g. Ito & Kappel (2002), Guidetti *et al.* (2004), Batkai *et al.* (2012). The few abstract error estimates available in the literature are not ready-to-apply such that information about basic approximation properties of the numerical method lead to a convergence rate, cf. for example Brenner *et al.* (1982) and Hussein & Kappel (2002). Finally, there is the framework of gradient discretization methods which was designed for a unified error analysis, but only covers elliptic and parabolic problems so far, cf. Droniou *et al.* (2017).



In order to systemize the derivation of convergence rates for other wave-type equations or new space discretizations thereof, we proceed in three steps. First, we propose a unified and abstract framework for wave-type equations and their space discretizations. Second, within this framework, we show that wave-type equations are well-posed and that the errors of their abstract space discretizations are bounded by a sum of interpolation errors, data errors and discretization errors. Third, we use more specific properties of the numerical method to prove the final a priori estimates. These a priori estimates allow to infer convergence rates by inserting known information about the numerical method in a modular way. We demonstrate the easy handling of our results by deriving new convergence results for an isoparametric bulk-surface finite element discretization of the wave equation with acoustic boundary conditions in a smooth domain Ω . Such discretizations are non-conforming since the computational domain does not coincide with Ω . In Hochbruck *et al.* (2017), the authors apply our a priori bounds to prove convergence rates of a heterogeneous multiscale discretization of Maxwell's equations using edge elements. Moreover, our results generalize former error estimates as they successfully reproduce convergence rates for several examples as the discontinuous Galerkin method for linear Maxwell's equations and finite elements with mass lumping for the acoustic wave equation.

The paper is organized as follows. In Sections 2.1 and 2.2, we introduce and analyze quasi-monotone evolution equations. For a unified treatment, we consider their space discretizations as differential equations in finite dimensional Hilbert spaces, as described in Section 2.3. In Section 2.4, we provide an overview of the tools and main ideas of the error analysis. Then, we prove a general error bound for stable space discretizations of quasi-monotone evolution equations in Section 2.5 and show a convergence result in the spirit of the Lax equivalence theorem in Section 2.6. The general error bound consists of data and discretization errors of the method. Under more specific assumptions on the structure of the wave equation and the numerical method, the discretization errors can be further analyzed against a sum of interpolation and conformity errors. The resulting error bounds then ultimately provide convergence rates. We discuss this for first-order wave-type equations in Section 3 and for second-order wave-type equations in Section 4. In order to provide a guideline for the reader who wants to find a concrete bound, or to prove convergence rates for a new application, we collected the most important assumptions, results and examples in two reference cards on the next page. In total, we show that our error

analysis is able to reproduce state-of-the-art convergence results for four applications. These examples are supplemented by a novel application presented in Section 5. There we use our a priori estimates to derive new convergence rates for an isoparametric bulk-surface finite element discretization of the wave equation with acoustic boundary conditions.

Reference card for first-order wave-type equations		
continuous	Assumption	The variational formulation of the PDE can be written as (3.1) and satisfies Assumption 3.1.
	<i>Well-posedness</i>	Follows from Theorem 2.4, see also (3.2)
	<i>Examples</i>	Advection equation in Example 2.6 and Maxwell's equation in Section 3.2
semi-discrete	Assumption	The space discretization is given by (3.3) and stable in the sense of Assumption 3.2.
		If $X_h^\ell \subset Y$:
		If $X_h^\ell \not\subset Y$:
	<i>Error bounds</i>	<i>Non-conforming</i> : Theorem 3.3 <i>Non-conforming</i> : Theorem 3.5 <i>Conforming</i> : Corollary 3.4 <i>Conforming</i> : Remark 3.6
	<i>Examples</i>	Edge elements for Maxwell in Section 3.2.1 DG method for Maxwell in Section 3.2.2

We emphasize that we focus on linear, inhomogeneous wave-type equations and error bounds in the energy norm. Error estimates in discrete norms, as derived for interior penalty discontinuous Galerkin discretizations in Grote *et al.* (2006), and convergence rates for parabolic problems, as given in Kovács & Lubich (2017) and Thomée (2006), are not covered. Moreover, we obtain sub-optimal convergence rates for discontinuous Galerkin discretizations stabilized with upwind fluxes as in Hochbruck & Pažur (2015). We further remark that the examples provided in this paper discuss finite element and discontinuous Galerkin methods, since they can deal with complex domains and provide high order approximations. However, we are convinced that our abstract estimates can also be used to derive convergence rates for other methods, e.g. finite difference or pseudospectral methods.

Reference card for second-order wave-type equations		
continuous	Assumption	The variational formulation of the PDE can be written as (4.1) and satisfies Assumption 4.1.
	<i>Well-posedness</i>	Follows from Theorem 4.3
	<i>Examples</i>	Acoustic wave equation with Dirichlet boundary conditions in Section 4.5 and with acoustic boundary conditions in Section 5
semi-discrete	Assumption	The space discretization is given by (4.8) and stable in the sense of Assumption 4.4.
	<i>Error bounds</i>	<i>Non-conforming</i> : Theorem 4.8 and Remark 4.9 <i>Conforming</i> : Corollary 4.10
	<i>Examples</i>	Lagrange elements with mass lumping in Section 4.5 and isoparametric bulk-surface finite elements in Section 5

Notation

In this section we collect the notation used throughout this paper. By C we denote a generic constant independent of time t and the space discretization parameter h . We consider problems on finite time intervals $[0, T]$, $T > 0$.

SPACES, NORMS, AND INNER PRODUCTS Let X, Y be two real Hilbert spaces with corresponding norms $\|\cdot\|_X, \|\cdot\|_Y$, respectively. By $\mathcal{L}(X, Y)$ we denote the space of all bounded linear operators from X to Y endowed with the operator norm

$$\|M\|_{Y \leftarrow X} := \sup_{\substack{x \in X \\ x \neq 0}} \frac{\|Mx\|_Y}{\|x\|_X} = \sup_{\substack{x \in X \\ \|x\|_X=1}} \|Mx\|_Y, \quad M \in \mathcal{L}(X, Y).$$

If $Y = \mathbb{R}$, then $X^* := \mathcal{L}(X, \mathbb{R})$ is the dual space of X and $\|\cdot\|_{X^*} := \|\cdot\|_{\mathbb{R} \leftarrow X}$. Moreover, for $\varphi \in X^*$ we define the duality pairing between X^* and X as

$$\langle \varphi, x \rangle_X := \varphi(x), \quad x \in X.$$

Let $b: Y \times X \rightarrow \mathbb{R}$ be a continuous bilinear form. Fixing the first argument of b yields an operator $b(y) := b(y, \cdot) \in X^*$ whose norm is given by

$$\|b(y)\|_{X^*} := \sup_{\substack{x \in X \\ \|x\|_X=1}} |b(y, x)|, \quad y \in Y.$$

Let $A: D(A) \rightarrow X$ be a linear operator defined on the subspace $D(A)$ of X . Then we denote by $[D(A)]$ the space $D(A)$ equipped with the graph norm of A (which is a Banach space if A is closed). In product spaces, we write

$$(u, v) := \begin{bmatrix} u \\ v \end{bmatrix} \in X^2 = X \times X.$$

The diagonal operator $\text{diag}(A_1, A_2): X_1 \times X_2 \rightarrow Y_1 \times Y_2$ for $A_i: X_i \rightarrow Y_i, i = 1, 2$ is defined by

$$\text{diag}(A_1, A_2) \begin{bmatrix} u \\ v \end{bmatrix} := \begin{bmatrix} A_1 u \\ A_2 v \end{bmatrix}.$$

Let $U \subset \mathbb{R}^d$ be a non-empty set. We define the supremum norm of $f: U \rightarrow X$ as

$$\|f\|_{\infty, U \rightarrow X} := \sup_{x \in U} \|f(x)\|_X$$

and use the short notation $\|f\|_{\infty, X} := \|f\|_{\infty, [0, T] \rightarrow X}$ for X -valued functions defined on $U = [0, T]$.

DOMAINS, BOUNDARIES, MESHES, AND DISCRETE SPACES The partial differential equations in this paper are considered in an open and bounded domain $\Omega \subset \mathbb{R}^d$. We denote its boundary by $\Gamma := \partial\Omega$ and the outer unit normal by $n: \Gamma \rightarrow \mathbb{R}^d$. For the scalar product in \mathbb{R}^d we write $x \cdot y$ for $x, y \in \mathbb{R}^d$ and $|x| := \sqrt{x \cdot x}$ denotes the Euclidean norm. We write $\gamma: H^1(\Omega) \rightarrow L^2(\Gamma)$ for the trace operator

and $\partial_n f: \Gamma \rightarrow \mathbb{R}$ is the normal derivative of $f: \Omega \rightarrow \mathbb{R}$. We use \mathcal{P}_k for the space of polynomials of maximal degree k . If not specified differently, we consider space discretizations based on an admissible mesh sequence $\mathcal{T}_{\mathcal{H}} = \{\mathcal{T}_h \mid h \in \mathcal{H}\}$ of Ω where the index h in \mathcal{T}_h denotes the maximal diameter of all the elements $K \in \mathcal{T}_h$ and $\Omega_h := \cup_{K \in \mathcal{T}_h} K$ is the computational domain. An admissible mesh sequence is shape-regular, contact-regular and satisfies an optimal polynomial approximation property, cf. (Di Pietro & Ern, 2012, Def. 1.57). We assume that \mathcal{T}_h consists of triangles or tetrahedra for $d = 2$ or $d = 3$, respectively, but our theory is not restricted to simplicial elements.

2. Evolution equations with linear monotone operators

We start the presentation of the unified framework by introducing evolution equations with linear monotone operators as an abstract formulation for wave-type equations. Such problems were already considered and analyzed by Showalter (1994), Showalter (1997) and Zeidler (1990a). After recalling conditions for their well-posedness, we then develop a theory for non-conforming space discretizations of evolution equations with linear monotone operators. For an overview of similar abstract approaches to space discretizations, we refer to Guidetti *et al.* (2004) and Ito & Kappel (2002).

2.1 Description of the continuous problem

Given a Gelfand triple of real Hilbert spaces

$$Y \hookrightarrow X \simeq X^* \hookrightarrow Y^*$$

we seek a solution $x: [0, T] \rightarrow Y$ of the evolution equation

$$x'(t) + \mathcal{S}x(t) = g(t) \quad \text{for } t \in [0, T], \quad (2.1a)$$

$$x(0) = x^0, \quad (2.1b)$$

where $g: [0, T] \rightarrow Y^*$ is a function and $\mathcal{S} \in \mathcal{L}(Y, Y^*)$ is a quasi-monotone operator.

DEFINITION 2.1 (Maximal and linear quasi-monotone operators) Let $W = Y^*$ or $W = X$.

- (i) An operator $\mathcal{S} \in \mathcal{L}(Y, W)$ is called *quasi-monotone* if there exists a constant $c_{\text{qm}} \geq 0$ s.t.

$$\langle \mathcal{S}y, y \rangle_Y + c_{\text{qm}} \|y\|_X^2 \geq 0 \quad \forall y \in Y. \quad (2.2a)$$

- (ii) A quasi-monotone operator $\mathcal{S} \in \mathcal{L}(Y, W)$ is called *maximal w.r.t. W* if there exists a $\lambda > c_{\text{qm}}$ s.t.

$$\text{range}(\lambda + \mathcal{S}) = W. \quad (2.2b)$$

REMARK 2.2 The theory of monotone operators is mostly used for non-linear functional problems. However, we feel that the term “quasi-monotone” is also suitable in our (linear) context, cf. also Showalter (1997) and Zeidler (1990a). A related notion can be found in ter Elst *et al.* (2015).

2.2 Well-posedness of the continuous problem

To apply semigroup theory, we restrict the operator \mathcal{S} to the Hilbert space X . The part of $\mathcal{S} \in \mathcal{L}(Y, Y^*)$ in X , as defined in Engel & Nagel (2000), is given by

$$S: D(S) \subset Y \rightarrow X, \quad y \mapsto Sy := \mathcal{S}y \quad \text{on } D(S) = \{y \in Y \mid \mathcal{S}y \in X\}. \quad (2.3)$$

The following lemma establishes a connection between quasi-monotone and dissipative operators. A similar result was shown in (Zeidler, 1990b, Sect. 31.4).

LEMMA 2.3. *Let $\mathcal{S} \in \mathcal{L}(Y, Y^*)$ and S be the part of \mathcal{S} in X as defined in (2.3).*

- (i) *If \mathcal{S} is quasi-monotone, then $-(S + c_{\text{qm}})$ is dissipative.*
- (ii) *If \mathcal{S} is quasi-monotone and maximal w.r.t. Y^* , then $\text{range}(\lambda + S) = X$ for all $\lambda > c_{\text{qm}}$ and $D(S)$ is dense in X .*

Proof. We only prove (ii), since (i) is obvious.

Let $f \in X$ be arbitrary. Since $X \xrightarrow{\text{d}} Y^*$, the maximality of \mathcal{S} ensures the existence of some $\lambda_0 > c_{\text{qm}}$ s.t. there is a $y \in Y$ which satisfies $(\lambda_0 + \mathcal{S})y = f$. Hence we have $\mathcal{S}y = f - \lambda_0 y \in X$, so that $y \in D(S)$ with $(\lambda_0 + S)y = f$. The surjectivity of $\lambda + S$ for all $\lambda > c_{\text{qm}}$ and the density of $D(S)$ follow from (Showalter, 1997, Prop. I.4.2). \square

To show the well-posedness of (2.1), we consider its corresponding abstract Cauchy problem in X .

THEOREM 2.4. *Let $W = Y^*$ or $W = X$ and assume that $\mathcal{S} \in \mathcal{L}(Y, W)$ is quasi-monotone and maximal w.r.t. W . If $x^0 \in D(S)$ and $g \in C([0, T]; [D(S)]) + C^1([0, T]; X)$, then (2.1) has a unique solution*

$$x \in C^1([0, T]; X) \cap C([0, T]; [D(S)])$$

which satisfies the stability estimate

$$\|x(t)\|_X \leq e^{c_{\text{qm}}t} \left(\|x^0\|_X + t \|g\|_{\infty, X} \right), \quad t \in [0, T]. \quad (2.4)$$

Proof. By Lemma 2.3 $-(S + c_{\text{qm}})$ is dissipative and satisfies the range condition. Hence it generates a contraction semigroup due to the Lumer-Philipp's theorem (Pazy, 1983, Sect. 1.3). This implies that $-S$ generates the C_0 -semigroup $(e^{-tS})_{t \geq 0}$ which satisfies

$$\|e^{-tS}\|_{X \leftarrow X} \leq e^{c_{\text{qm}}t}.$$

Under the assumptions on x^0 and g , the abstract Cauchy problem

$$x'(t) + Sx(t) = g(t), \quad t \in [0, T], \quad x(0) = x^0 \quad (2.5)$$

has a unique solution $x \in C^1([0, T]; X) \cap C([0, T]; [D(S)])$ which is given by Duhamel's formula

$$x(t) = e^{-tS}x^0 + \int_0^t e^{-(t-s)S}g(s) \, ds,$$

cf. (Pazy, 1983, Sect. 4.2). The stability estimate thus follows from

$$\|x(t)\| \leq e^{c_{\text{qm}}t} \|x^0\|_X + \int_0^t e^{c_{\text{qm}}(t-s)} \|g(s)\|_X \, ds \leq e^{c_{\text{qm}}t} \left(\|x^0\|_X + \|g\|_{\infty, X} \int_0^t 1 \, ds \right).$$

Finally, since $X \simeq X^* \xrightarrow{\text{d}} Y^*$ and $S = \mathcal{S}|_{D(S)}$, every solution of (2.5) also solves (2.1). \square

In the following, let $p: X \times X \rightarrow \mathbb{R}$ denote the inner product on X and $\langle \cdot, \cdot \rangle_Y$ the duality pairing between Y^* and Y . Then we have

$$p(z, y) = z(y) = \langle z, y \rangle_Y \quad \forall z \in X, y \in Y, \quad (2.6)$$

as an immediate consequence of the identification $X \simeq X^*$.

2.3 Space discretization

This section is dedicated to non-conforming space discretizations of (2.1). Such space discretizations seek to approximate the solution $x \in X$ in a finite dimensional Hilbert space X_h with inner product $p_h(\cdot, \cdot)$ and norm $\|\cdot\|_{X_h}$. Here, $h > 0$ corresponds to a discretization parameter of X_h , e.g. the maximal diameter of all elements of a mesh. We emphasize that in general

$$X_h \not\subset X.$$

A space discretization of (2.1) is a differential equation in X_h seeking $x_h: [0, T] \rightarrow X_h$ s.t.

$$x_h'(t) + S_h x_h(t) = g_h(t) \quad \text{for } t \in [0, T], \quad (2.7a)$$

$$x_h(0) = x_h^0 \in X_h, \quad (2.7b)$$

where $S_h \in \mathcal{L}(X_h, X_h)$ is a discretization of S , e.g. resulting from a finite element or dG method, and $g_h: [0, T] \rightarrow X_h$ is an approximation of g .

Following (Ciarlet, 2002, Chap. 4), we define conforming space discretizations. For that purpose, it is convenient to write the operators as bilinear forms. We denote the bilinear form associated with \mathcal{S} by

$$s(z, y) := \langle \mathcal{S}z, y \rangle_Y, \quad z, y \in Y, \quad (2.8)$$

and, analogously, the bilinear form associated with S_h by

$$s_h(z_h, y_h) := p_h(S_h z_h, y_h), \quad z_h, y_h \in X_h. \quad (2.9)$$

To motivate the criteria for a conforming method, we give the variational formulations of the continuous and the semi-discrete problem. Theorem 2.4 shows that the evolution equation (2.1) has, under suitable assumptions on the data, a solution x satisfying $x(t) \in D(S)$ and $x'(t) \in X$, $t \geq 0$. Considering (2.1) in variational form and using (2.6), we thus obtain that x solves

$$p(x'(t), y) + s(x(t), y) = p(g(t), y) \quad \forall y \in Y. \quad (2.10)$$

Analogously, the differential equation (2.7a) can be cast as

$$p_h(x_h'(t), y_h) + s_h(x_h(t), y_h) = p_h(g_h(t), y_h) \quad \forall y_h \in X_h.$$

DEFINITION 2.5 The space discretization (2.7) of the evolution equation (2.1) is called *conforming* if the following three conditions are satisfied.

- (i) $X_h \subset Y$,
- (ii) $p(z_h, y_h) = p_h(z_h, y_h)$ for all $z_h, y_h \in X_h$,
- (iii) $s(z_h, y_h) = s_h(z_h, y_h)$ for all $z_h, y_h \in X_h$.

Space discretizations which violate at least one of these conditions are called *non-conforming*.

Note that these conditions are not completely independent of each other: $X_h \subset X$ is needed for the second and $X_h \subset Y$ for the third condition. An overview of examples which fit into the unified framework and a classification of their non-conformity is given in Table 1.

Table 1. Overview and classification of non-conformity of examples from the unified framework

	$X_h \subset Y$	$p = p_h$	$s = s_h$	Discussed in
Advection eq. with Lagrange elements	✓	✓	✓	Example 2.6
Maxwell's eq. with Nédélec elements	✓	✓	✓	Section 3.2.1
Maxwell's eq. with discontinuous Galerkin	✗	✓	✗	Section 3.2.2
Heterogeneous multiscale method for Maxwell's eq.	✓	✗	✓	Hochbruck <i>et al.</i> (2017)
Wave eq. with Lagrange elements	✓	✓	✓	Section 4.5
Wave eq. with Lagrange elements with mass lumping	✓	✗	✓	Section 4.5
Wave eq. with acoustic bc. in smooth domains	✗	✗	✗	Section 5

EXAMPLE 2.6 To illustrate our exposition we consider the advection equation as a model problem, see, e.g. (Di Pietro & Ern, 2012, Chap. 2). Let $\Omega \subset \mathbb{R}^d$ be a bounded, polygonal, convex domain. We seek a function $x: [0, T] \times \Omega \rightarrow \mathbb{R}$ s.t.

$$x_t + \beta \cdot \nabla x + \mu x = f \quad \text{in } \Omega, \quad (2.11a)$$

$$x = 0 \quad \text{on } \Gamma^-, \quad (2.11b)$$

$$x(0) = x^0 \quad \text{in } \Omega. \quad (2.11c)$$

Here, $\mu \geq 0$, $\beta \in \mathbb{R}^d$, ∇x denotes the gradient of x and

$$\Gamma^- = \{x \in \Gamma \mid \beta \cdot n(x) < 0\} \quad (2.12)$$

denotes the inflow part of the boundary Γ .

Comparing the variational formulation of this problem with (2.10) shows that p is the $L^2(\Omega)$ inner product and

$$s(z, y) = \int_{\Omega} \mu zy + (\beta \cdot \nabla z) y \, dx. \quad (2.13)$$

Therefore, we choose $X = L^2(\Omega)$ and Y as the natural domain of the differential operator

$$Y = \{y \in L^2(\Omega) \mid \beta \cdot \nabla y \in L^2(\Omega), y|_{\Gamma^-} = 0\} \quad (2.14)$$

equipped with the graph norm

$$\|y\|_Y^2 = p(y, y) + p(\beta \cdot \nabla y, \beta \cdot \nabla y). \quad (2.15)$$

It is easy to see that Y is a dense subspace of X and that the associated operator $\mathcal{S} \in \mathcal{L}(Y, X)$ of s is monotone (i.e. $c_{\text{qm}} = 0$) and maximal w.r.t. X , see, e.g., (Di Pietro & Ern, 2012, Theorem 2.9). Thus the problem is well-posed for suitable initial values and source terms due to Theorem 2.4.

We consider a space discretization with linear finite elements on a triangulation \mathcal{T}_h of Ω . Hence X_h is the space of piecewise linear functions defined on \mathcal{T}_h equipped with the inner product $p_h = p$. We further have $\Omega_h = \Omega$ s.t. $X_h \subset Y$ and $s_h = s$, since the polygonal domain Ω is exactly triangulated. Therefore, the finite element discretization is conforming due to Definition 2.5.




FIG. 1. Overview of spaces and operators

2.4 Notation for spaces and operators

The approximation $x_h \in X_h$ obtained from a non-conforming space discretization with $X_h \not\subset X$ cannot be compared directly with the solution $x \in X$. Consider for example a finite element discretization of a partial differential equation in a smooth domain Ω where the computational domain $\Omega_h \neq \Omega$ only approximates Ω . In such a situation, we have $X_h \not\subset X$, since the finite element functions in X_h are defined in Ω_h and not in Ω . To deal with this issue, we assume there exists a linear operator

$$Q_h : X_h \rightarrow X \quad (2.16)$$

which reconstructs the approximation $Q_h x_h \approx x$ in X . We call Q_h the *lift operator*, as it “lifts” the approximation x_h to the lifted discrete space

$$X_h^\ell := Q_h(X_h).$$

For conforming methods, the lift operator can be chosen as $Q_h = I$ which implies $X_h^\ell = X_h$. Examples of non-trivial lift operators can be found in, e.g., Elliott & Ranner (2013), (Ciarlet, 2002, Chap. 4) and Cockburn *et al.* (2014). To map from continuous function spaces into the discrete space X_h , we introduce $J_h \in \mathcal{L}(Z, X_h)$ where Z is a Hilbert space that is continuously embedded in X . We call J_h the *reference operator*, since we base our error bounds on the following splitting of the error:

$$\|Q_h x_h - x\|_X \leq \|Q_h(x_h - J_h x)\|_X + \|(Q_h J_h - I)x\|_X.$$

To obtain optimal convergence rates, the choice of J_h has to fit to the application. For conforming methods, we choose the standard orthogonal projection onto X_h (w.r.t. p). However, for non-conforming methods, we will see below that a suitable interpolation operator $I_h : Z_{ip} \rightarrow X_h$ has to be used for J_h to prove optimal convergence rates. In this case, the space $Z = Z_{ip}$ is typically a higher order (broken) Sobolev space which ensures that the interpolation operator is continuous, i.e., $I_h \in \mathcal{L}(Z_{ip}, X_h)$.

The X -orthogonal projection onto the lifted discrete space X_h^ℓ is denoted by

$$\Pi_h : X \rightarrow X_h^\ell, \quad p((1 - \Pi_h)z, Q_h y_h) = 0 \quad \forall z \in X, y_h \in X_h. \quad (2.17a)$$

Moreover, we introduce the adjoint lift Q_h^* to map between these spaces via

$$Q_h^* : X \rightarrow X_h, \quad p_h(Q_h^* z, y_h) = p(z, Q_h y_h) \quad \forall z \in X, y_h \in X_h, \quad (2.17b)$$

and further set

$$P_h := Q_h Q_h^* : X \rightarrow X_h^\ell. \quad (2.17c)$$

An overview of all involved mappings and spaces is given in Figure 1.

REMARK ON CONFORMING METHODS For a conforming method, where $Q_h = I$ and $X_h^\ell = X_h$, many of these operators coincide. More precisely,

$$P_h = \Pi_h = Q_h^*$$

is just the p -orthogonal projection of X onto X_h .

Our error bounds will be given in terms of the remainder operator

$$R_h := Q_h^* S - S_h J_h : D(S) \cap Z \rightarrow X_h \quad (2.18a)$$

and in terms of conformity errors represented by the differences of the bilinear forms which are

$$\Delta p(z_h, y_h) := p(Q_h z_h, Q_h y_h) - p_h(z_h, y_h), \quad z_h, y_h \in X_h \quad (2.18b)$$

$$\Delta s(z_h, y_h) := s(Q_h z_h, Q_h y_h) - s_h(z_h, y_h), \quad z_h, y_h \in X_h. \quad (2.18c)$$

Note that the definition of Δs requires $X_h^\ell \subset Y$.

2.5 A priori error bounds

In applications where \mathcal{S} is a differential operator, $S_h \in \mathcal{L}(X_h, X_h)$ is not uniformly bounded w.r.t. h . Therefore, we assume the semi-discretization to be stable in the following sense.

DEFINITION 2.7 (Stability) We call the space discretization (2.7) *stable* if

- (i) the discrete operator $S_h \in \mathcal{L}(X_h, X_h)$ is quasi-monotone in X_h with $\widehat{c}_{\text{qm}} \geq 0$ s.t.

$$p_h(S_h y_h, y_h) + \widehat{c}_{\text{qm}} \|y_h\|_{X_h}^2 \geq 0 \quad \forall y_h \in X_h,$$

- (ii) the lift operator $Q_h : X_h \rightarrow X$ is continuous with $c_X \geq 0$ s.t.

$$\|Q_h y_h\|_X \leq c_X \|y_h\|_{X_h} \quad \forall y_h \in X_h. \quad (2.19)$$

REMARK 2.8 Under this assumption, Theorem 2.4 (with X replaced by X_h) shows that there exists a unique solution x_h of (2.7) with

$$\|x_h(t)\|_{X_h} \leq e^{\widehat{c}_{\text{qm}} t} (\|x_h^0\|_{X_h} + t \|g_h\|_{\infty, X_h}). \quad (2.20)$$

Note that our error analysis only makes use of (2.20) and not the quasi-monotonicity of S_h directly.

REMARK ON CONFORMING METHODS For conforming methods, the stability assumptions follow directly from the monotonicity of \mathcal{S} with $\widehat{c}_{\text{qm}} = c_{\text{qm}}$. Moreover, since the inner products of X and X_h coincide and $Q_h = I$, we have $c_X = 1$.

We now state the most general error bound of the unified error analysis.

THEOREM 2.9. *Let the assumptions of Theorem 2.4 be fulfilled and assume that the solution x of (2.1) satisfies $x \in C^1([0, T]; Z)$. If the space discretization (2.7) is stable, then the error of the semi-discrete approximation $Q_h x_h$ is bounded by*

$$\|Q_h x_h(t) - x(t)\|_X \leq C e^{\widehat{c}_{\text{qm}} t} \left(E_{\text{data}}(t) + t \|(Q_h^* - J_h)x'\|_{\infty, X_h} + t \|R_h x\|_{\infty, X_h} \right) + \|(I - Q_h J_h)x(t)\|_X \quad (2.21)$$

for $t \in [0, T]$, a constant C which is independent of h and t ,

$$E_{\text{data}}(t) := \|x_h^0 - J_h x^0\|_{X_h} + t \|g_h - Q_h^* g\|_{\infty, X_h} \quad (2.22)$$

and R_h defined in (2.18a).

Proof. Let $e_h := x_h - J_h x$ denote the discrete error. By splitting the error and using that $Q_h \in \mathcal{L}(X_h, X)$ for stable space discretizations, we obtain

$$\|Q_h x_h - x\|_X \leq \|Q_h e_h\|_X + \|(Q_h J_h - I)x\|_X \leq c_X \|e_h\|_{X_h} + \|(Q_h J_h - I)x\|_X. \quad (2.23)$$

Hence it is sufficient to bound the discrete error.

Since $x \in C^1([0, T]; Z)$ and $J_h \in \mathcal{L}(Z, X_h)$, we have $e_h \in C^1([0, T]; X_h)$ and

$$e_h' = x_h' - J_h x' = x_h' - Q_h^* x' + (Q_h^* - J_h)x'.$$

We rewrite the first part using (2.7a) and (2.5)

$$x_h' - Q_h^* x' = -S_h x_h + g_h - Q_h^* (-Sx + g) = -S_h e_h + g_h - Q_h^* g + (Q_h^* S - S_h J_h)x.$$

Inserting this into the previous equation shows that the discrete error e_h satisfies the differential equation

$$e_h' + S_h e_h = g_h - Q_h^* g + (Q_h^* S - S_h J_h)x + (Q_h^* - J_h)x'.$$

The discrete stability estimate (2.20) therefore yields

$$\|e_h(t)\|_{X_h} \leq e^{\hat{c}_{\text{qm}} t} \left(\|e_h(0)\|_{X_h} + t \left(\|g_h - Q_h^* g\|_{\infty, X_h} + \|(Q_h^* - J_h)x'\|_{\infty, X_h} + \|R_h x\|_{\infty, X_h} \right) \right).$$

Using this estimate in (2.23) completes the proof. \square

REMARK 2.10 For $J_h = Q_h^*$ the error bound (2.21) simplifies to

$$\|Q_h x_h(t) - x(t)\|_X \leq C e^{\hat{c}_{\text{qm}} t} \left(E_{\text{data}}(t) + t \|R_h x\|_{\infty, X_h} \right) + \|(I - P_h)x(t)\|_X,$$

where $R_h = Q_h^* S - S_h Q_h^*$. Since $Z = X$ in this case, the error bound is valid without further assumptions on the solution obtained by Theorem 2.4 .

2.6 Convergence

In the rest of this section, we show that $Q_h x_h$ converges to the exact solution for $h \rightarrow 0$, if the space discretization is stable and consistent in the following sense.

DEFINITION 2.11 (Consistency) We call the space discretization (2.7) of (2.1) *consistent* if

- (i) for all $y_h \in X_h$, we have $\|\Delta p(y_h)\|_{X_h^*} \rightarrow 0$, $h \rightarrow 0$,
- (ii) for all $z \in Z$, we have $\|(I - Q_h J_h)z\|_X \rightarrow 0$, $h \rightarrow 0$,
- (iii) for all $z \in D(S) \cap Z$, we have $\|R_h z\|_{X_h} \rightarrow 0$, $h \rightarrow 0$.

EXAMPLE 2.6 (continued) For the finite element discretization of the advection equation (2.11), we have $X = L^2(\Omega)$ and $\Delta p = 0$. Therefore, Assumption 2.11 (i) is fulfilled. If we choose J_h as the nodal interpolation operator $I_h: Z \rightarrow X_h$ with $Z = H^2(\Omega)$, then Assumption 2.11 (ii) follows from

$$\|(\mathbf{I} - I_h)z\|_{L^2(\Omega)} \leq Ch^2|z|_{H^2(\Omega)}, \quad z \in H^2(\Omega), \quad (2.24)$$

cf. (Brenner & Scott, 2008, Sect. 4.4). Hence only Assumption 2.11 (iii) remains to be verified. This will be done in the course of Section 3.

The following lemma provides a fundamental estimate which we will use frequently in the rest of this article. It bounds the difference between the adjoint lift operator and the reference operator by the sum of a reference error and a conformity error of the inner products.

LEMMA 2.12. *If $Q_h \in \mathcal{L}(X_h, X)$ satisfies (2.19), then*

$$\|(Q_h^* - J_h)z\|_{X_h} \leq c_X \|(\mathbf{I} - Q_h J_h)z\|_X + \|\Delta p(J_h z)\|_{X_h^*}, \quad z \in Z.$$

Proof. First observe that for all $z_h \in X_h$

$$\|z_h\|_{X_h} = \max_{\|y_h\|_{X_h}=1} p_h(z_h, y_h). \quad (2.25)$$

Therefore, we have for $z \in Z$ by (2.18b)

$$\begin{aligned} \|(Q_h^* - J_h)z\|_{X_h} &= \max_{\|y_h\|_{X_h}=1} p_h((Q_h^* - J_h)z, y_h) \\ &= \max_{\|y_h\|_{X_h}=1} p_h(Q_h^* z, y_h) - p_h(J_h z, y_h) \\ &= \max_{\|y_h\|_{X_h}=1} p(z, Q_h y_h) - p(Q_h J_h z, Q_h y_h) + p(Q_h J_h z, Q_h y_h) - p_h(J_h z, y_h) \\ &= \max_{\|y_h\|_{X_h}=1} p((\mathbf{I} - Q_h J_h)z, Q_h y_h) + \Delta p(J_h z, y_h) \\ &\leq c_X \|(\mathbf{I} - Q_h J_h)z\|_X + \|\Delta p(J_h z)\|_{X_h^*}. \end{aligned}$$

This was the claim. \square

COROLLARY 2.13. *Let the assumptions of Theorem 2.4 be fulfilled and assume that the unique solution x of (2.1) satisfies $x \in C^1([0, T]; Z)$.*

- (i) *If the space discretization (2.7) is stable, then the error of the semi-discrete approximation $Q_h x_h$ is bounded by*

$$\begin{aligned} \|Q_h x_h(t) - x(t)\|_X &\leq C e^{\hat{c}_{\text{qm}} t} (1+t) \left(E_{\text{data}}(1) + \|(\mathbf{I} - Q_h J_h)x\|_{\infty, X} + \|(\mathbf{I} - Q_h J_h)x'\|_{\infty, X} \right. \\ &\quad \left. + \|\Delta p(J_h x')\|_{\infty, X_h^*} + \|R_h x\|_{\infty, X_h} \right) \quad (2.26) \end{aligned}$$

for $t \in [0, T]$, a constant C which is independent of h and t , and E_{data} and R_h as defined in (2.22) and (2.18a), respectively.

(ii) If the space discretization (2.7) is stable and consistent, and $g(t) \in Z$, $t \in [0, T]$ with

$$\|x_h^0 - J_h x^0\|_{X_h} \rightarrow 0 \quad \text{and} \quad \|g_h - J_h g\|_{\infty, X_h} \rightarrow 0, \quad h \rightarrow 0,$$

then the semi-discrete approximation converges, i.e.,

$$\|Q_h x_h(t) - x(t)\|_X \rightarrow 0, \quad h \rightarrow 0,$$

for $t \in [0, T]$.

Proof. (i) The desired estimate follows directly from the general error bound (2.21) and Lemma 2.12.

(ii) Using Lemma 2.12 and Assumptions 2.11 (i)-(ii) to estimate the source term error in E_{data} , we obtain

$$\begin{aligned} \|g_h(t) - Q_h^* g(t)\|_{X_h} &\leq \|g_h(t) - J_h g(t)\|_{X_h} + \|(J_h - Q_h^*)g(t)\|_{X_h} \\ &\leq \|g_h(t) - J_h g(t)\|_{X_h} + c_X \|(I - Q_h J_h)g(t)\|_X + \|\Delta p(J_h g(t))\|_{X_h^*} \rightarrow 0 \end{aligned}$$

for $h \rightarrow 0$, since $g(t) \in Z$. Because the initial value converges by assumption, we thus showed $E_{\text{data}} \rightarrow 0$, $h \rightarrow 0$. All other terms in the upper bound of (2.26) vanish as $h \rightarrow 0$ due to the consistency of the method. This completes the proof. \square

For a specific application, the a priori error estimate (2.26) still needs to be complemented with a bound on the remainder term $\|R_h x\|_{X_h}$. In the following, we will show such bounds for space discretizations of first-order wave-type equations and second-order wave-type equations.

3. First-order wave-type equations

This section is devoted to the error analysis of non-conforming space discretizations of first-order wave-type equations. This class of wave equations comprises symmetric hyperbolic systems as defined in Benzoni-Gavage & Serre (2007) or Burazin & Erceg (2016), but also general dissipative first-order partial differential equations.

Let $Y \xrightarrow{d} X \simeq X^* \xrightarrow{d} Y^*$ be a Gelfand triple of real Hilbert spaces. We seek the solution $x: [0, T] \rightarrow Y$ of the first-order wave-type equation

$$p(x'(t), y) + s(x(t), y) = p(g(t), y) \quad \forall y \in Y, \quad (3.1a)$$

$$x(0) = x^0, \quad (3.1b)$$

where p denotes the inner product on X , $g: [0, T] \rightarrow X$ is a function and s is a bilinear form which satisfies the following assumption.

ASSUMPTION 3.1 (First-order wave-type equations)

(i) The bilinear form $s: Y \times X \rightarrow \mathbb{R}$ is continuous, i.e. there is a constant $c_s > 0$ s.t.

$$|s(z, y)| \leq c_s \|z\|_Y \|y\|_X \quad \forall z \in Y, y \in X.$$

(ii) The bilinear form s is quasi-monotone and maximal w.r.t. X , i.e. there is a constant $c_{\text{qm}} > 0$ s.t.

$$s(y, y) + c_{\text{qm}} \|y\|_X^2 \geq 0 \quad \forall y \in Y$$

and there exists a $\lambda > c_{\text{qm}}$ s.t. for every $f \in X$ there is a unique $z \in Y$ s.t.

$$\lambda p(z, y) + s(z, y) = p(f, y) \quad \forall y \in X.$$

Note that (3.1) is equivalent to the evolution equation (2.1) where \mathcal{S} is induced by s . As a consequence of Assumption 3.1, $\mathcal{S} \in \mathcal{L}(Y, X)$ is a maximal and quasi-monotone operator with $D(\mathcal{S}) = Y$ and $\|\mathcal{S}\|_{X \leftarrow Y} = c_s$. Therefore the first-order wave-type equation (3.1) has a unique solution

$$x \in C^1([0, T]; X) \cap C([0, T]; Y) \quad (3.2)$$

for initial values x^0 and source terms g which satisfy the conditions of Theorem 2.4 with $W = X$.

3.1 A priori error bounds

Next we consider space discretizations of first-order wave-type equations which yield an approximation $x_h: [0, T] \rightarrow X_h$ in the finite dimensional vector space X_h s.t.

$$p_h(x_h'(t), y_h) + s_h(x_h(t), y_h) = p_h(g_h(t), y_h) \quad \forall y_h \in X_h, \quad (3.3a)$$

$$x_h(0) = x_h^0. \quad (3.3b)$$

Here $g_h: [0, T] \rightarrow X_h$ is a function and p_h, s_h are bilinear forms which satisfy the following assumption.

ASSUMPTION 3.2 (Stability)

- (i) The bilinear form p_h is an inner product on X_h and induces the norm $\|y_h\|_{X_h}^2 := p_h(y_h, y_h)$.
- (ii) There is a constant $\widehat{c}_{\text{qm}} > 0$ s.t.

$$s_h(y_h, y_h) + \widehat{c}_{\text{qm}} \|y_h\|_{X_h}^2 \geq 0 \quad \forall y_h \in X_h.$$

- (iii) The lift operator $Q_h: X_h \rightarrow X$ is continuous with $\|Q_h y_h\|_X \leq c_X \|y_h\|_{X_h}$ for all $y_h \in X_h$.

The goal of this section is to derive a priori error estimates for $Q_h x_h$ in terms of interpolation and conformity errors. Since Assumption 3.2 guarantees that the evolution equation (2.7) associated with (3.3) is stable in the sense of Definition 2.7, we can employ the estimate from Corollary 2.13 (i). Therefore it only remains to estimate $\|R_h x\|_{X_h}$ in terms of errors of the interpolation operator $I_h \in \mathcal{L}(Z_{\text{ip}}, X_h)$, $Z_{\text{ip}} \xrightarrow{\text{cd}} Y$, and the conformity errors Δp and Δs as defined in (2.18).

Motivated by the applications presented in Section 3.2, we distinguish between two different cases. The finite element method leads to semi-discrete problems where $X_h^\ell \subset Y$ and the discontinuous Galerkin method to $X_h^\ell \not\subset Y$ but $X_h^\ell \subset X$.

3.1.1 *Space discretizations with $X_h^\ell \subset Y$.* In this section, we consider space discretizations where the lifted discrete space X_h^ℓ is not only contained in X but also in the smaller space Y .

THEOREM 3.3. *Let the assumptions of Theorem 2.4 be fulfilled and assume that the solution x of the first-order wave-type equation (3.1) satisfies $x \in C^1([0, T]; Z_{\text{ip}})$. If the space discretization (3.3) satisfies Assumption 3.2 and $X_h^\ell \subset Y$, then the error of the semi-discrete approximation $Q_h x_h$ is bounded by*

$$\begin{aligned} \|Q_h x_h(t) - x(t)\|_X \leq C e^{\widehat{c}_{\text{qm}} t} (1+t) & \left(e_{\text{data}} + \|(I - Q_h I_h) x'\|_{\infty, X} + \|(I - Q_h I_h) x\|_{\infty, Y} \right. \\ & \left. + \|\Delta p(I_h x')\|_{\infty, X_h^*} + \|\Delta s(I_h x)\|_{\infty, X_h^*} \right) \end{aligned}$$

for $t \in [0, T]$, a constant C which is independent of h and t , and

$$e_{\text{data}} := \|x_h^0 - I_h x^0\|_{X_h} + \|g_h - Q_h^* g\|_{\infty, X_h}. \quad (3.4)$$

Proof. By assumption we have $X_h^\ell \subset Y$ and $\mathcal{S} \in \mathcal{L}(Y, X)$. Therefore, we obtain for $y_h \in X_h$

$$\begin{aligned} p_h(R_h x, y_h) &= p_h((Q_h^* S - S_h J_h)x, y_h) = p(Sx, Q_h y_h) - p_h(S_h J_h x, y_h) \\ &= s(x, Q_h y_h) - s_h(J_h x, y_h) \\ &= s((I - Q_h J_h)x, Q_h y_h) + s(Q_h(J_h x), Q_h y_h) - s_h(J_h x, y_h) \\ &\leq c_s \|(I - Q_h J_h)x\|_Y \|Q_h y_h\|_X + \Delta s(J_h x, y_h) \\ &\leq c_s \|(I - Q_h J_h)x\|_Y c_X \|y_h\|_{X_h} + \Delta s(J_h x, y_h), \end{aligned}$$

where we used Assumption 3.2 (iii) for the last inequality. Thus it follows from (2.25) that

$$\|R_h x\|_{X_h} = \max_{\|y_h\|_{X_h}=1} p_h(R_h x, y_h) \leq c_X c_s \|(I - Q_h J_h)x\|_Y + \|\Delta s(J_h x)\|_{X_h^*}. \quad (3.5)$$

Finally, we choose $J_h = I_h$. The desired estimate then follows from $Y \stackrel{d}{\hookrightarrow} X$ and Corollary 2.13 (i). \square

For conforming methods, we obtain an error bound independent of x' if we choose $J_h = \Pi_h$. To prove this bound, we use that any two norms on the finite dimensional space X_h are equivalent. This implies that there exists a $\delta_h > 0$ s.t.

$$\delta_h \|y_h\|_Y \leq \|y_h\|_X, \quad y_h \in X_h. \quad (3.6)$$

In the context of finite element methods, such inequalities are called inverse estimates and we usually have $\delta_h \rightarrow 0$ as $h \rightarrow 0$.

COROLLARY 3.4. *Let the assumptions of Theorem 2.4 be fulfilled and let x be the unique solution of the first-order wave-type equation (3.1) satisfying $x(t) \in Z_{\text{ip}}$, $t \in [0, T]$. If the space discretization (3.3) is conforming due to Definition 2.5 and fulfills Assumption 3.2 for $Q_h = I$, then the error of the semi-discrete approximation x_h is bounded by*

$$\|x_h(t) - x(t)\|_X \leq C e^{c_{\text{qm}} t} (1+t) \left(e_{\text{data}} + \delta_h^{-1} \|(I - I_h)x\|_{\infty, X} + \|(I - I_h)x\|_{\infty, Y} \right)$$

for $t \in [0, T]$, a constant C which is independent of h and t , and e_{data} defined in (3.4).

Proof. First note that conforming methods are stable with $\widehat{c}_{\text{qm}} = c_{\text{qm}}$ and $c_X = 1$. For the error analysis of conforming methods, we choose $J_h = \Pi_h = Q_h^* \in \mathcal{L}(X, X_h)$ with $Z = X$ such that the simplified estimate from Remark 2.10 applies. Moreover, (3.5) and $\Delta s \equiv 0$ imply

$$\|R_h x\|_{X_h} \leq C \|(I - \Pi_h)x\|_Y.$$

To obtain an estimate in terms of interpolation errors, we apply (3.6) and use that Π_h is the best approximation w.r.t. the X -norm

$$\begin{aligned} \|(I - \Pi_h)x\|_Y &\leq \|(I - I_h)x\|_Y + \|(I_h - \Pi_h)x\|_Y \\ &\leq \|(I - I_h)x\|_Y + \delta_h^{-1} \|(I_h - \Pi_h)x\|_X \\ &\leq \|(I - I_h)x\|_Y + \delta_h^{-1} \left(\|(I_h - I)x\|_X + \|(I - \Pi_h)x\|_X \right), \\ &\leq \|(I - I_h)x\|_Y + 2\delta_h^{-1} \|(I - I_h)x\|_X. \end{aligned} \quad (3.7)$$

Collecting terms then yields the final estimate. \square

EXAMPLE 2.6 (continued) For the finite element discretization of the advection equation (2.11), there exists a δ_h s.t. $\delta_h^{-1} \leq Ch^{-1}$, cf. (Brenner & Scott, 2008, Lem. 4.5.3). Moreover, the interpolation error converges linearly in the Y -norm, since

$$\|(\mathbf{I} - I_h)z\|_Y \leq \|(\mathbf{I} - I_h)z\|_{H^1(\Omega)} \leq Ch|z|_{H^2(\Omega)}, \quad z \in H^2(\Omega),$$

and quadratically in $X = L^2(\Omega)$, as we showed in (2.24). Thus, we obtain from Corollary 3.4 that the error of the finite element approximation is bounded by

$$\|x_h(t) - x(t)\|_{L^2(\Omega)} \leq C(1+t)h\|x\|_{\infty, H^2(\Omega)},$$

if $x(t) \in H^2(\Omega)$, $t \in [0, T]$ and $x_h^0 = \Pi_h x^0$, $g_h(t) = \Pi_h g(t)$, $t \in [0, T]$. For similar results, we refer to Layton (1983) and Dunca (2017).

3.1.2 *Space discretizations with $X_h^\ell \not\subset Y$.* In this section, we consider space discretizations where

$$X_h^\ell \subset X \quad \text{and} \quad X_h^\ell \not\subset Y.$$

This situation appears e.g. for discontinuous Galerkin (dG) methods which approximate the solution by a function which may have discontinuities between elements of the mesh. A typical example is the discretization of an advection equation in the broken polynomial space $X_h = \mathcal{P}_k(\mathcal{T}_h)$ consisting of piecewise polynomials of degree k on a triangulation \mathcal{T}_h of Ω .

For our error analysis of such space discretizations it is necessary to insert the exact solution x into $s_h(\cdot, y_h)$ for $y_h \in X_h$. Thus we assume that $s_h: X_h \times X_h \rightarrow \mathbb{R}$ can be extended to

$$\overline{s}_h: (X_h + Z_{\text{ip}}) \times X_h \rightarrow \mathbb{R}. \quad (3.8a)$$

Furthermore, we define

$$\overline{\Delta s}(z, y_h) := s(z, Q_h y_h) - \overline{s}_h(z, y_h), \quad z \in Z_{\text{ip}} \cap Y, y_h \in X_h. \quad (3.8b)$$

In this setting we can show the following error bound.

THEOREM 3.5. *Let the assumptions of Theorem 2.4 be fulfilled and assume that the solution x of the first-order wave-type equation (3.1) satisfies $x \in C^1([0, T]; Z_{\text{ip}})$. If the space discretization (3.3) fulfills Assumption 3.2 and s_h can be extended to (3.8a), then the error of the semi-discrete approximation $Q_h x_h$ is bounded by*

$$\begin{aligned} \|Q_h x_h(t) - x(t)\|_X &\leq C e^{\widehat{c}_{\text{qm}} t} (1+t) \left(e_{\text{data}} + \|(\mathbf{I} - Q_h I_h)x'\|_{\infty, X} + \|(\mathbf{I} - Q_h I_h)x\|_{\infty, X} \right. \\ &\quad \left. + \|\overline{s}_h((\mathbf{I} - I_h)x)\|_{\infty, X_h^*} + \|\overline{\Delta s}(x)\|_{\infty, X_h^*} + \|\Delta p(I_h x')\|_{\infty, X_h^*} \right) \end{aligned}$$

for $t \in [0, T]$, a constant C which is independent of h and t , and e_{data} defined in (3.4).

Proof. Since the solution x belongs to $Z_{\text{ip}} \cap Y$, we find for $y_h \in X_h$

$$\begin{aligned} p_h(R_h x, y_h) &= s(x, Q_h y_h) - \overline{s}_h(J_h x, y_h) = s(x, Q_h y_h) - \overline{s}_h(x, y_h) + \overline{s}_h(x, y_h) - \overline{s}_h(J_h y, y_h) \\ &= \overline{\Delta s}(x, y_h) + \overline{s}_h((\mathbf{I} - J_h)x, y_h). \end{aligned}$$

By (2.25), taking the maximum over all y_h with $\|y_h\|_{X_h} = 1$ thus yields

$$\|R_h x\|_{X_h} \leq \|\overline{s}_h((I - J_h)x)\|_{X_h^*} + \|\overline{\Delta}s(x)\|_{X_h^*}.$$

The claim now follows from Corollary 2.13 (i) and setting $J_h = I_h$. \square

REMARK 3.6 If $Q_h = I$, $\Delta p \equiv 0$, and $\overline{\Delta}s \equiv 0$, and if the assumptions of Theorem 3.5 are satisfied, then similar arguments with $J_h = \Pi_h$ show

$$\|x_h(t) - x(t)\|_X \leq C e^{\widehat{c}_{\text{qm}} t} (1+t) \left(e_{\text{data}} + \|\overline{s}_h((I - \Pi_h)x')\|_{\infty, X_h^*} + \|(I - I_h)x\|_{\infty, X} \right), \quad (3.9)$$

cf. Corollary 3.4. Note that instead of $x \in C^1([0, T]; Z_{\text{ip}})$, we *only* need to assume $x(t) \in Z_{\text{ip}}$ for this estimate.

3.2 Examples: Maxwell's equations

As the prototype of a first-order wave-type equation we consider Maxwell's equations for linear isotropic materials with perfectly conducting boundary conditions, cf. Kirsch & Hettlich (2015).

Let $E: [0, T] \times \Omega \rightarrow \mathbb{R}^3$ be the electric field and $H: [0, T] \times \Omega \rightarrow \mathbb{R}^3$ be the magnetic field in a polyhedral domain $\Omega \subset \mathbb{R}^3$ given by

$$\begin{aligned} \mu H_t &= -\text{curl} E && \text{in } \Omega, \\ \varepsilon E_t &= \text{curl} H && \text{in } \Omega, \\ n \times E &= 0 && \text{on } \Gamma, \\ H(0) &= H^0, \quad E(0) = E^0 && \text{in } \Omega, \end{aligned}$$

where the permittivity and the permeability $\varepsilon, \mu \in L^\infty(\Omega)$ are uniformly positive. We assume that the initial values satisfy $\text{div}(\varepsilon E^0) = \text{div}(\mu H^0) = 0$ in Ω and $n \cdot (\mu H^0) = 0$ on Γ . Then $E(t)$ and $H(t)$ satisfy these conditions for all $t \geq 0$, cf. (Hochbruck *et al.*, 2015, Prop. 3.5).

The suitable functional analytic setting for $x = [H, E]^T$ is given by the Hilbert space $X := L^2(\Omega)^6$ endowed with a weighted inner product and $Y = H(\text{curl}, \Omega) \times H_0(\text{curl}, \Omega)$ which is densely and continuously embedded into X . Maxwell's equations are a first-order wave-type equation since the Maxwell operator $\mathcal{S} \in \mathcal{L}(Y, X)$ is skew-symmetric and maximal, cf., e.g. (Hochbruck *et al.*, 2015, Sect. 3.2). Hence Maxwell's equations are well-posed due to Theorem 2.4 for $x^0 \in Y$.

Since Ω is polyhedral in this application, we assume that the computational domain satisfies $\Omega_h = \Omega$ in the following examples. Moreover, we assume that $x_h^0 = I_h x^0$ s.t. $e_{\text{data}} = 0$.

3.2.1 *Edge element discretizations.* In this example, we consider a space discretization of Maxwell's equation using first order curl-conforming elements of Nédélec's second type on a quasi-uniform mesh \mathcal{T}_h , cf. Nédélec (1986). For such space discretizations, we have $X_h = V_h(\text{curl}) \times V_{h,0}(\text{curl})$ where

$$\begin{aligned} V_h(\text{curl}) &= \left\{ U_h \in H(\text{curl}, \Omega) \mid U_h|_K \in (\mathcal{P}_1)^3 \text{ for } K \in \mathcal{T}_h \right\}, \\ V_{h,0}(\text{curl}) &= \{ U_h \in V_h(\text{curl}) \mid \mathbf{v} \times U_h = 0 \text{ on } \Gamma \}, \end{aligned}$$

and the discrete inner product and differential form are given by $p_h = p$ and $s_h = s$. This is possible since $X_h \subset Y$ by construction. Moreover, there exists an interpolation operator $I_h: Z_{\text{ip}} \rightarrow X_h$, $Z_{\text{ip}} = H^2(\Omega)^6$

s.t.

$$\|(\mathbf{I} - I_h)z\|_X + h\|(\mathbf{I} - I_h)z\|_Y \leq Ch^2\|z\|_{H^2(\Omega)^6}, \quad z \in H^2(\Omega)^6,$$

cf. (Nédélec, 1986, Prop. 3), and the inverse estimate between $L^2(\Omega)$ and $H^1(\Omega)$ implies $\delta_h^{-1} \leq Ch^{-1}$ if $0 < h \leq 1$.

Therefore, we are in the situation of Section 3.1.1 and the a priori estimate from Corollary 3.4 applies. If $x = (H, E) \in C([0, T]; H^2(\Omega)^6)$, then the approximation properties of the interpolation imply that the semi-discrete solution $x_h = (H_h, E_h)$ converges linearly in h with

$$\|x_h(t) - x(t)\|_{L^2(\Omega)^6} \leq C(1+t)h.$$

A similar convergence result for elements of Nédélec's first type can be found in (Zhao, 2004, Thm. 4.1).

Observe that $x = (H, E) \in C([0, T]; H^2(\Omega)^6)$ can only be guaranteed under additional assumptions on x^0 , ε , μ and Ω , cf. for example (Hochbruck *et al.*, 2015, Lem. 3.7) for sufficient conditions if Ω is a cuboid.

3.2.2 Discontinuous Galerkin discretizations. Discontinuous Galerkin methods are a very competitive approach to approximate Maxwell's equation numerically. This example investigates a discontinuous Galerkin discretization where s_h stems from a central (also centered) fluxes dG discretization of the Maxwell operator, cf. Di Pietro & Ern (2012), and which seeks an approximation in the set of piecewise polynomials $X_h = \mathcal{P}_k(\mathcal{T}_h)^6$, $k \geq 0$ on \mathcal{T}_h . Then the extension of s_h to $(Y \cap H^1(\mathcal{T}_h)^6) \times X_h$ is consistent in the sense that $\overline{\Delta s} \equiv 0$. Moreover, we have by (Hochbruck & Sturm, 2016, (5.3) and (5.5))

$$\|(\mathbf{I} - I_h)z\|_X + h\|\overline{s_h}((\mathbf{I} - \Pi_h)z)\|_{X_h^*} \leq C \left(\sum_{K \in \mathcal{T}_h} h_K^{2k+2} |z|_{H^{k+1}(K)^6}^2 \right)^{1/2}, \quad z \in Y \cap H^{k+1}(\mathcal{T}_h)^6,$$

where $I_h: H^2(\mathcal{T}_h)^6 \rightarrow X_h$ is the piecewise nodal interpolation operator and $|x|_{H^{k+1}(K)^6}$ the $H^{k+1}(K)^6$ semi-norm of x . Hence $Z_{\text{ip}} = Y \cap H^2(\mathcal{T}_h)^6$ is a suitable choice for our setting.

The convergence result then follows from (3.9). If the solution x of Maxwell's equations belongs to $C([0, T]; H^2(\mathcal{T}_h)^6)$ then the dG approximation $x_h = (E_h, H_h)$ satisfies

$$\|x_h(t) - x(t)\|_{L^2(\Omega)^6} \leq C(t)h^k.$$

This result can for example be found in (Fezoui *et al.*, 2005, Thm. 3.5).

4. Second-order wave-type equations

In this section, we consider wave-problems formulated as second-order evolution equations. Our abstract formulation covers a wide range of problems including wave equations with dynamic boundary conditions and problems with damping or advection effects.

4.1 Description of the continuous problem

Let H and V be two Hilbert spaces with $V \xrightarrow{\text{d}} H$, i.e., there is a constant $C_{H,V} > 0$ s.t.

$$\|v\|_H \leq C_{H,V} \|v\|_V, \quad v \in V.$$

By $m: H \times H \rightarrow \mathbb{R}$ we denote the inner product of H and we identify $H \simeq H^*$ to form the Gelfand triple

$$V \xrightarrow{\text{d}} H \simeq H^* \xrightarrow{\text{d}} V^*.$$

The second-order wave-type equation then reads: Find $u: [0, T] \rightarrow V$ s.t.

$$\langle u''(t), v \rangle_V + b(u'(t), v) + a(u(t), v) = \langle f(t), v \rangle_V \quad \forall v \in V, \quad (4.1a)$$

$$u(0) = u_1^0, \quad u'(0) = u_2^0, \quad (4.1b)$$

where $f: [0, T] \rightarrow V^*$ is a given function and where the bilinear forms a and b satisfy the following assumption.

ASSUMPTION 4.1 (Second-order wave-type equations)

(i) The bilinear form $a: V \times V \rightarrow \mathbb{R}$ is continuous, symmetric and satisfies the Gårding inequality

$$a(v, v) + c_G \|v\|_H^2 \geq \alpha \|v\|_V^2, \quad v \in V, \quad (4.2)$$

for constants $c_G \geq 0$ and $\alpha > 0$.

(ii) The bilinear form $b: V \times V \rightarrow \mathbb{R}$ is continuous and there is a constant $\rho_{\text{qm}} \geq 0$ s.t. $b + \rho_{\text{qm}} m$ is monotone, i.e.,

$$b(v, v) + \rho_{\text{qm}} \|v\|_H^2 \geq 0, \quad v \in V.$$

Since the bilinear forms a and b induce operators $\mathcal{A}, \mathcal{B} \in \mathcal{L}(V, V^*)$, respectively, we can write (4.1) equivalently as the evolution equation

$$u'' + \mathcal{B}u' + \mathcal{A}u = f \quad \text{in } V^* \quad (4.3)$$

supplemented by initial conditions $u(0) = u_1^0$ and $u'(0) = u_2^0$.

Furthermore, we introduce the bilinear form

$$\tilde{a}(w, v) := a(w, v) + c_G m(w, v), \quad w, v \in V, \quad (4.4)$$

which is coercive on $V \times V$ due to (4.2), and define $\tilde{V} = (V, \tilde{a})$ as the Hilbert space equipped with \tilde{a} . Note that the Gårding inequality implies

$$\|v\|_H \leq C_{H,V} \|v\|_V \leq C_{H,V} \alpha^{-1/2} \|v\|_{\tilde{V}}, \quad v \in \tilde{V}. \quad (4.5)$$

4.2 Well-posedness of the continuous problem

Introducing $u_1 = u$ and the velocity $u_2 = u'$, the second order problem (4.3) can be written as a first-order in time problem (2.1) with

$$x(t) = \begin{bmatrix} u_1(t) \\ u_2(t) \end{bmatrix}, \quad \mathcal{S} = \begin{bmatrix} 0 & -\mathbf{I} \\ \mathcal{A} & \mathcal{B} \end{bmatrix}, \quad g(t) = \begin{bmatrix} 0 \\ f(t) \end{bmatrix}, \quad x^0 = \begin{bmatrix} u_1^0 \\ u_2^0 \end{bmatrix}. \quad (4.6a)$$

A suitable Gelfand triple for this evolution equation is given via

$$Y = \tilde{V} \times V \quad \text{and} \quad X = \tilde{V} \times H, \quad (4.6b)$$

equipped with their canonical inner products. In the following, we will refer to (2.1) with (4.6) as the first-order in time formulation of the second-order wave-type equation (4.3).

Variants of the following results can be found in the proof of (Showalter, 1994, Thm. VI.2.1). A complete proof is given in (Hipp, 2017, Lem. 6.2).

LEMMA 4.2. *If Assumption 4.1 is satisfied, then $\mathcal{S} \in \mathcal{L}(Y, Y^*)$ is quasi-monotone with*

$$c_{\text{qm}} = \frac{1}{2}c_G C_{H,V} \alpha^{-1/2} + \rho_{\text{qm}} \quad (4.7)$$

and maximal w.r.t. Y^* .

Expressing Theorem 2.4 in terms of (4.6) gives the following result.

THEOREM 4.3. *Let Assumption 4.1 be fulfilled. If the initial values $u_1^0, u_2^0 \in V$ satisfy $\mathcal{A}u_1^0 + \mathcal{B}u_2^0 \in H$ and $f \in C^1([0, T]; H)$ or $(f, \mathcal{B}f) \in C([0, T]; V \times H)$, then the second-order wave-type equation (4.3) has a unique solution $u \in C^2([0, T]; H) \cap C^1([0, T]; V)$ which satisfies $\mathcal{A}u + \mathcal{B}u' \in C([0, T]; H)$ and*

$$\left(\|u(t)\|_{\tilde{V}}^2 + \|u'(t)\|_H^2 \right)^{1/2} \leq e^{c_{\text{qm}}t} \left(\left(\|u_1^0\|_{\tilde{V}}^2 + \|u_2^0\|_H^2 \right)^{1/2} + t \|f\|_{\infty, H} \right), \quad t \in [0, T],$$

for c_{qm} from (4.7).

If further $\mathcal{B} \in \mathcal{L}(V, H)$, then we have $u \in C^2([0, T]; H) \cap C^1([0, T]; V) \cap C([0, T]; [D(A)])$ where $D(A) = \{v \in V \mid \mathcal{A}v \in H\}$ and $A = \mathcal{A}|_{D(A)}$.

Proof. The assumptions guarantee that \mathcal{S} , x^0 and g from (4.6) are such that Theorem 2.4 applies. More precisely, \mathcal{S} is quasi-monotone due to Lemma 4.2, $x^0 = (u_1^0, u_2^0) \in D(S)$, and, $g \in C^1([0, T]; X)$ or $Sg \in C([0, T]; X)$. By Theorem 2.4, (2.1) has the unique solution x and we obtain from (4.6) that

$$x \in C([0, T]; [D(S)]) \cap C^1([0, T]; \tilde{V} \times H),$$

where

$$D(S) = \{y \in Y \mid \mathcal{S}y \in X\} = \{(v_1, v_2) \in \tilde{V} \times V \mid \mathcal{A}v_1 + \mathcal{B}v_2 \in H\}.$$

Since $x = (u_1, u_2)$ and $u_1 = u$, $u_2 = u'$, the stability estimate follows from (2.4) and we have $u'' = u'_2 \in C([0, T]; H)$. Moreover, $Sx \in C([0, T]; X)$ implies $t \mapsto \mathcal{A}u(t) + \mathcal{B}u'(t) \in C([0, T]; H)$.

If $\mathcal{B} \in \mathcal{L}(V, H)$, then $D(S) = D(A) \times V$ and therefore $u \in C([0, T]; [D(A)])$, which gives the second claim. \square

4.3 Space discretization

The aim of this section is to derive a priori estimates for non-conforming space discretizations of (4.1) in the finite dimensional vector space V_h , which determine the approximation $u_h: [0, T] \rightarrow V_h$ as the solution of

$$m_h(u_h''(t), v_h) + b_h(u_h'(t), v_h) + a_h(u_h(t), v_h) = m_h(f_h(t), v_h) \quad \forall v_h \in V_h, \quad (4.8a)$$

$$u_h(0) = u_{h,1}^0, \quad u_h'(0) = u_{h,2}^0. \quad (4.8b)$$

Here $u_{h,1}^0, u_{h,2}^0 \in V_h$, $f_h: [0, T] \rightarrow V_h$ and $m_h, b_h, a_h: V_h \times V_h \rightarrow \mathbb{R}$ are the discrete counterparts of u_1^0, u_2^0 , f and m, b, a , respectively. Since we do not assume that $V_h \subset V$ this ansatz covers a wide range of non-conforming space discretizations. Analogously to Section 2.4, we assume that there exists a lift operator

$$Q_h^V: V_h \rightarrow V,$$

which yields the lifted approximation $Q_h^V u_h \in V$ of the exact solution u of (4.1).

STABILITY For the stability of the space discretization (4.8) and our error analysis, we assume that it satisfies the following properties.

ASSUMPTION 4.4 (Stability) The following conditions hold for $w_h, v_h \in V_h$.

(i) The bilinear form m_h is an inner product on V_h and induces the norm

$$\|v_h\|_{m_h}^2 := m_h(v_h, v_h).$$

(ii) The bilinear form a_h is monotone and symmetric and there is a constant $0 \leq \widehat{c}_G \leq 1$ s.t.

$$\widetilde{a}_h(w_h, v_h) := a_h(w_h, v_h) + \widehat{c}_G m_h(w_h, v_h) \quad (4.9a)$$

is positive definite and induces the norm

$$\|v_h\|_{\widetilde{a}_h}^2 := \widetilde{a}_h(v_h, v_h).$$

(iii) There is a constant $C_{m_h, \widetilde{a}_h} > 0$ independent of h s.t. $\|v_h\|_{m_h} \leq C_{m_h, \widetilde{a}_h} \|v_h\|_{\widetilde{a}_h}$.

(iv) There is a constant $\widehat{\rho}_{\text{qm}} \geq 0$ s.t. the bilinear form $b_h + \widehat{\rho}_{\text{qm}} m_h$ is monotone.

(v) The lift operator Q_h^V is continuous from $(V_h, \|\cdot\|_{m_h})$ into H with $c_H \geq 0$ s.t.

$$\|Q_h^V v_h\|_H \leq c_H \|v_h\|_{m_h}.$$

(vi) The lift operator Q_h^V is continuous from $(V_h, \|\cdot\|_{\widetilde{a}_h})$ into \widetilde{V} with $c_V \geq 0$ s.t.

$$\|Q_h^V v_h\|_{\widetilde{V}} \leq c_V \|v_h\|_{\widetilde{a}_h}.$$

If Assumption 4.4 is satisfied then we write $H_h := (V_h, m_h)$ and $\widetilde{V}_h := (V_h, \widetilde{a}_h)$ for V_h equipped with the inner product m_h and \widetilde{a}_h , respectively. Furthermore, note that $a_h: \widetilde{V}_h \times \widetilde{V}_h \rightarrow \mathbb{R}$ satisfies

$$|a_h(w_h, v_h)| \leq \|w_h\|_{\widetilde{a}_h} \|v_h\|_{\widetilde{a}_h}, \quad w_h, v_h \in \widetilde{V}_h, \quad (4.10)$$

since it is a monotone symmetric bilinear form.

REMARK 4.5 If we choose $\widehat{c}_G = 1$, then Assumption 4.4 (ii) and Assumption 4.4 (iii) with $C_{m_h, \widetilde{a}_h} = 1$ are always fulfilled. However, $\widehat{c}_G > 0$ leads to exponential growth of the constants in t , while equations where $\widehat{c}_G = \widehat{\rho}_{\text{qm}} = 0$, only exhibit linear growth, cf. Theorem 4.8. Therefore, smaller constants c_G are to be favored, since they lead to sharper bounds.

FORMULATION IN THE FRAMEWORK OF MONOTONE OPERATORS To write (4.8) as a differential equation, we define the operators

$$\begin{aligned} A_h: H_h &\rightarrow H_h, & m_h(A_h w_h, v_h) &= a_h(w_h, v_h), & w_h, v_h &\in V_h, \\ \text{and } B_h: H_h &\rightarrow H_h, & m_h(B_h w_h, v_h) &= b_h(w_h, v_h), & w_h, v_h &\in V_h. \end{aligned}$$

Then we can express the variational problem (4.8) as the second-order differential equation

$$u_h''(t) + B_h u_h'(t) + A_h u_h(t) = f_h(t), \quad u_h(0) = u_{h,1}^0, \quad u_h'(0) = u_{h,2}^0,$$

or, equivalently, as the first-order differential equation (2.7) with

$$x_h = \begin{bmatrix} u_h \\ u'_h \end{bmatrix}, \quad S_h = \begin{bmatrix} 0 & -I_{V_h} \\ A_h & B_h \end{bmatrix}, \quad g_h = \begin{bmatrix} 0 \\ f_h(t) \end{bmatrix}, \quad x_h^0 = \begin{bmatrix} u_{h,1}^0 \\ u_{h,2}^0 \end{bmatrix} \quad (4.11a)$$

in the Hilbert space

$$X_h = \widetilde{V}_h \times H_h \quad (4.11b)$$

endowed with the inner product

$$p_h((w_{h,1}, w_{h,2}), (v_{h,1}, v_{h,2})) := \widetilde{a}_h(w_{h,1}, v_{h,1}) + m_h(w_{h,2}, v_{h,2}). \quad (4.11c)$$

Finally, we define the lift operator $Q_h : X_h \rightarrow X$ as

$$Q_h \begin{bmatrix} w_{h,1} \\ w_{h,2} \end{bmatrix} := \begin{bmatrix} Q_h^V w_{h,1} \\ Q_h^V w_{h,2} \end{bmatrix}.$$

Note that one can also choose two different lifts for the components $w_{h,1}$ and $w_{h,2}$. For the ease of presentation, we refrain from investigating this here.

NOTATION To apply our results from Section 2 for the second-order wave equations, we need to write the operators from Section 2.4 componentwise. The components of the adjoint lift $Q_h^* = \text{diag}(Q_h^{V*}, Q_h^{H*})$ are characterized by

$$\begin{aligned} Q_h^{H*} : H &\rightarrow H_h, & m_h(Q_h^{H*} w, v_h) &= m(w, Q_h^V v_h), & w \in H, v_h \in V_h, \\ \text{and } Q_h^{V*} : V &\rightarrow \widetilde{V}_h, & \widetilde{a}_h(Q_h^{V*} w, v_h) &= \widetilde{a}(w, Q_h^V v_h), & w \in V, v_h \in V_h, \end{aligned}$$

and the components of the orthogonal projections $\Pi_h = \text{diag}(\Pi_h^V, \Pi_h^H)$ by

$$\begin{aligned} m((I - \Pi_h^H)w, Q_h^V v_h) &= 0, & w \in H, v_h \in V_h, \\ \text{and } \widetilde{a}((I - \Pi_h^V)w, Q_h^V v_h) &= 0, & w \in V, v_h \in V_h. \end{aligned}$$

The difference between the bilinear forms is denoted by

$$\begin{aligned} \Delta m(w_h, v_h) &:= m(Q_h^V w_h, Q_h^V v_h) - m_h(w_h, v_h), & w_h, v_h \in V_h, \\ \text{and } \Delta \widetilde{a}(w_h, v_h) &:= \widetilde{a}(Q_h^V w_h, Q_h^V v_h) - \widetilde{a}_h(w_h, v_h), & w_h, v_h \in V_h. \end{aligned}$$

We define the reference operator $J_h = \text{diag}(J_h^V, J_h^H)$ using the operators $J_h^V : Z_1 \rightarrow V_h$ and $J_h^H : Z_2 \rightarrow V_h$ on the Hilbert spaces $Z_1 \hookrightarrow V$ and $Z_2 \hookrightarrow H$. Finally, since norms on finite dimensional vector spaces are equivalent, there is an $\varepsilon_h > 0$ s.t.

$$\varepsilon_h \|v_h\|_{\widetilde{a}_h} \leq \|v_h\|_{m_h}, \quad v_h \in V_h. \quad (4.12)$$

REMARK 4.6 If X_h is a finite element space based on a mesh \mathcal{T}_h of Ω with mesh width h and the discretization satisfies $\|\cdot\|_{m_h} \sim \|\cdot\|_{L^2(\Omega)}$ and $\|\cdot\|_{\widetilde{a}_h} \sim \|\cdot\|_{H^1(\Omega)}$, then we have $\varepsilon_h^{-1} \leq Ch^{-1}$ due to the inverse estimate from (Brenner & Scott, 2008, Lem. 4.5.3).

4.4 A priori error bounds

As a first step towards an a priori error bound, we estimate the remainder operator R_h from (2.18a).

LEMMA 4.7. *Let $z = (w_1, w_2) \in (Z_1 \times Z_2) \cap (V \times V)$ s.t. $\mathcal{A}w_1 + \mathcal{B}w_2 \in H$. If Assumption 4.4 is satisfied, then the remainder term is bounded by*

$$\begin{aligned} \|R_h z\|_{X_h} \leq & C \left(\|\Delta \tilde{a}(J_h^H w_2)\|_{\tilde{V}_h^*} + \|\Delta \tilde{a}(J_h^H w_1)\|_{\tilde{V}_h^*} + \|\Delta m(J_h^H w_1)\|_{H_h^*} \right. \\ & + \|(I - Q_h^V J_h^H)w_1\|_{\tilde{V}} + \|(I - Q_h^V J_h^H)w_2\|_{\tilde{V}} + \varepsilon_h^{-1} \|(Q_h^{V^*} - J_h^V)w_1\|_{\tilde{a}_h} \\ & \left. + \max_{\|v_h\|_{m_h}=1} |b(w_2, Q_h^V v_h) - b_h(J_h^H w_2, v_h)| \right). \end{aligned}$$

Proof. Recall that p and p_h denote the inner products on X and X_h respectively, that

$$R_h \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} -(Q_h^{V^*} - J_h^H)w_2 \\ Q_h^{H^*}(\mathcal{A}w_1 + \mathcal{B}w_2) - (A_h J_h^V w_1 + B_h J_h^H w_2) \end{bmatrix}$$

by $R_h = Q_h^* S - S_h J_h$ with (4.6) and (4.11), and that

$$\|R_h z\|_{X_h} = \max_{\|y_h\|_{X_h}=1} p_h(R_h z, y_h), \quad z \in Z \cap Y. \quad (4.13)$$

Let $y_h = (v_{h,1}, v_{h,2}) \in X_h$ with $\|y_h\|_{X_h}^2 = \|v_{h,1}\|_{\tilde{a}_h}^2 + \|v_{h,2}\|_{m_h}^2 = 1$. Then we have for the right hand side

$$\begin{aligned} p_h(R_h z, y_h) &= -\tilde{a}_h((Q_h^{V^*} - J_h^H)w_2, v_{h,1}) + m_h(Q_h^{H^*}(\mathcal{A}w_1 + \mathcal{B}w_2) - (A_h J_h^V w_1 + B_h J_h^H w_2), v_{h,2}) \\ &= -\tilde{a}_h((Q_h^{V^*} - J_h^H)w_2, v_{h,1}) + \left(a(w_1, Q_h^V v_{h,2}) - a_h(J_h^V w_1, v_{h,2}) \right) \\ &\quad + \left(b(w_2, Q_h^V v_{h,2}) - b_h(J_h^H w_2, v_{h,2}) \right). \end{aligned}$$

For the first term, we use the Cauchy–Schwarz inequality for \tilde{a}_h and $\|v_{h,1}\|_{\tilde{a}_h} \leq 1$ to obtain

$$\tilde{a}_h((Q_h^{V^*} - J_h^H)w_2, v_h) \leq \|(Q_h^{V^*} - J_h^H)w_2\|_{\tilde{a}_h} \|v_h\|_{\tilde{a}_h} \leq c_V \|(I - Q_h J_h^H)w_2\|_{\tilde{V}} + \|\Delta \tilde{a}(J_h^H w_2)\|_{\tilde{V}_h^*},$$

where we applied Lemma 2.12 with $X_h = \tilde{V}_h$ in the second inequality.

For the upper bound of the second term, we first add and subtract $Q_h^{V^*}$ and then rewrite a and a_h in terms of \tilde{a} and \tilde{a}_h using (4.4) and (4.9a), respectively. The first difference then vanishes due to the definition of $Q_h^{V^*}$, while applying (4.10) and the Cauchy–Schwarz inequality for m_h to the remaining terms yields

$$\begin{aligned} & a(w_1, Q_h^V v_{h,2}) - a_h(J_h^V w_1, v_{h,2}) \\ & \leq a(w_1, Q_h^V v_{h,2}) - a_h(Q_h^{V^*} w_1, v_{h,2}) + a_h((Q_h^{V^*} - J_h^V)w_1, v_{h,2}) \\ & \leq \left(\tilde{a}(w_1, Q_h^V v_{h,2}) - \tilde{a}_h(Q_h^{V^*} w_1, v_{h,2}) \right) + \left(c_G m(w_1, Q_h^V v_{h,2}) - \widehat{c}_G m_h(Q_h^{V^*} w_1, v_{h,2}) \right) \\ & \quad + \|(Q_h^{V^*} - J_h^V)w_1\|_{\tilde{a}_h} \|v_{h,2}\|_{\tilde{a}_h} \\ & \leq \max\{c_G, \widehat{c}_G\} |m_h((Q_h^{V^*} - Q_h^{H^*})w_1, v_{h,2})| + \|(Q_h^{V^*} - J_h^V)w_1\|_{\tilde{a}_h} \|v_{h,2}\|_{\tilde{a}_h} \\ & \leq \max\{c_G, \widehat{c}_G\} \|(Q_h^{V^*} - Q_h^{H^*})w_1\|_{m_h} + \varepsilon_h^{-1} \|(Q_h^{V^*} - J_h^V)w_1\|_{\tilde{a}_h}. \end{aligned}$$

Here we used (4.12) and $\|v_{h,2}\|_{m_h} \leq 1$ in the last step. To further estimate the first term, we split it into two parts, use Assumption 4.4 (iii) for the first one, and then employ the estimate from Lemma 2.12 with $X_h = \tilde{V}_h$ and $X_h = H_h$. This yields

$$\begin{aligned}
& \|(\mathcal{Q}_h^{V*} - \mathcal{Q}_h^{H*})w_1\|_{m_h} \\
& \leq \|(\mathcal{Q}_h^{V*} - J_h^H)w_1\|_{m_h} + \|(J_h^H - \mathcal{Q}_h^{H*})w_1\|_{m_h} \\
& \leq C_{m_h, \tilde{a}_h} \|(\mathcal{Q}_h^{V*} - J_h^H)w_1\|_{\tilde{a}_h} + \|(J_h^H - \mathcal{Q}_h^{H*})w_1\|_{m_h} \\
& \leq C_{m_h, \tilde{a}_h} \left(c_V \|(\mathbb{I} - \mathcal{Q}_h^V J_h^H)w_1\|_{\tilde{V}} + \|\Delta \tilde{a}(J_h^H w_1)\|_{\tilde{V}^*} \right) + c_H \|(\mathbb{I} - \mathcal{Q}_h^V J_h^H)w_1\|_H + \|\Delta m(J_h^H w_1)\|_{H_h^*}, \\
& \leq \left(C_{m_h, \tilde{a}_h} c_V + C_{H,V} c_H \alpha^{-1/2} \right) \|(\mathbb{I} - \mathcal{Q}_h^V J_h^H)w_1\|_V + \|\Delta \tilde{a}(J_h^H w_1)\|_{\tilde{V}^*} + \|\Delta m(J_h^H w_1)\|_{H_h^*},
\end{aligned}$$

where the last estimate follows from (4.5).

Inserting the above estimates in (4.13) and collecting terms yields the desired bound. \square

To derive an a priori error bound for non-conforming space discretizations of second-order wave-type equations, we only need to combine the general error bound for monotone evolution equations from Theorem 2.21 with the estimate of the remainder term. Since the error is bounded solely in terms of data, interpolation, and conformity errors, it directly leads to convergence rates for concrete applications. For this purpose, we introduce the continuous interpolation operator $I_h: Z_{\text{ip}} \rightarrow V_h$ which is defined on a continuously embedded Hilbert space Z_{ip} in V .

THEOREM 4.8. *Let the assumptions of Theorem 4.3 be fulfilled and assume that the solution u of the second-order wave-type equation (4.1) satisfies $u \in C^2([0, T]; Z_{\text{ip}})$. If the space discretization (4.8) fulfills Assumption 4.4, then the error of the semi-discrete approximation $\mathcal{Q}_h^V u_h$ is bounded by*

$$\|\mathcal{Q}_h^V u_h(t) - u(t)\|_{\tilde{V}} + \|\mathcal{Q}_h^V u_h'(t) - u'(t)\|_H \leq C e^{\hat{c}_{\text{qm}} t} (1+t) (\varepsilon_{\text{data}} + \varepsilon_{\text{ip}} + \varepsilon_{\text{forms}} + \varepsilon_{\Delta b})$$

for $t \in [0, T]$, where C is independent of h and t , $\hat{c}_{\text{qm}} = \hat{c}_G C_{m_h, \tilde{a}_h} / 2 + \hat{\rho}_{\text{qm}}$, and

$$\varepsilon_{\text{data}} := \|u_{h,1}^0 - \mathcal{Q}_h^{V*} u_1^0\|_{\tilde{a}_h} + \|u_{h,2}^0 - I_h u_2^0\|_{m_h} + \|f_h - \mathcal{Q}_h^{H*} f\|_{\infty, H_h}, \quad (4.14a)$$

$$\varepsilon_{\text{ip}} := \|(\mathbb{I} - \mathcal{Q}_h^V I_h)u\|_{\infty, \tilde{V}} + \|(\mathbb{I} - \mathcal{Q}_h^V I_h)u'\|_{\infty, \tilde{V}} + \|(\mathbb{I} - \mathcal{Q}_h^V I_h)u''\|_{\infty, H}, \quad (4.14b)$$

$$\varepsilon_{\text{forms}} := \|\Delta \tilde{a}(I_h u)\|_{\infty, \tilde{V}_h^*} + \|\Delta m(I_h u)\|_{\infty, H_h^*} + \|\Delta \tilde{a}(I_h u')\|_{\infty, \tilde{V}_h^*} + \|\Delta m(I_h u'')\|_{\infty, H_h^*}, \quad (4.14c)$$

$$\varepsilon_{\Delta b} := \left\| \max_{\|v_h\|_{m_h}=1} |b(u', \mathcal{Q}_h^V v_h) - b_h(I_h u', v_h)| \right\|_{\infty, \mathbb{R}}. \quad (4.14d)$$

Proof. Theorem 2.9 applies since (2.7) with (4.11) is stable on $X_h = \tilde{V}_h \times H_h$ in the sense of Definition 2.7: By Assumption 4.4, m_h , b_h , and a_h have the same properties as their continuous counterparts. Hence, Lemma 4.2 applied to S_h yields that S_h is maximal and quasi-monotone with $\hat{c}_{\text{qm}} = \hat{c}_G C_{m_h, \tilde{a}_h} / 2 + \hat{\rho}_{\text{qm}}$. Moreover, Assumptions 4.4 (v) and 4.4 (vi) imply that the lift is continuous from X_h into X .

Thus the error bound (2.21) holds for $x = (u, u')$ and $x_h = (u_h, u_h')$ where u is the solution of (4.1) and u_h is the solution of (4.8). Since the left-hand side of (2.21) is bounded from below by

$$\|\mathcal{Q}_h^V u_h - u\|_{\tilde{V}} + \|\mathcal{Q}_h^V u_h' - u'\|_H \leq \sqrt{2} \|\mathcal{Q}_h x_h - x\|_X,$$

it remains to provide an upper bound for the terms on the right-hand side of (2.21). First, we choose $J_h^V = Q_h^{V*} \in \mathcal{L}(V, V_h)$ and $J_h^H = I_h \in \mathcal{L}(Z_{\text{ip}}, V_h)$. Then we have $E_{\text{data}} \leq \sqrt{2}\varepsilon_{\text{data}}$ for E_{data} defined in (2.22). Second, we obtain by Lemma 2.12 with $X = H$

$$\|(Q_h^* - J_h)x'\|_{X_h} = \|(Q_h^{H*} - I_h)u''\|_{m_h} \leq c_H \|(I - Q_h^V I_h)u''\|_H + \|\Delta m(I_h u'')\|_{H_h^*},$$

and therefore $\|(Q_h^* - J_h)x'\|_{X_h} \leq C(\varepsilon_{\text{ip}} + \varepsilon_{\text{forms}})$. Third, we apply Lemma 4.7 to $\|R_h x\|_{X_h}$. Since we chose $J_h^V = Q_h^{V*}$ and $J_h^H = I_h$, the estimate simplifies to

$$\begin{aligned} \|R_h x\|_{X_h} &\leq C \left(\|\Delta \tilde{a}(I_h u')\|_{\tilde{V}_h^*} + \|\Delta \tilde{a}(I_h u)\|_{\tilde{V}_h^*} + \|\Delta m(I_h u)\|_{H_h^*} \right. \\ &\quad \left. + \|(I - Q_h^V I_h)u\|_{\tilde{V}} + \|(I - Q_h^V I_h)u'\|_{\tilde{V}} + \max_{\|v_h\|_{m_h}=1} |b(u', Q_h^V v_h) - b_h(I_h u', v_h)| \right). \end{aligned}$$

Hence, $\|R_h x\|_{\infty, X_h} \leq C(\varepsilon_{\text{ip}} + \varepsilon_{\text{forms}} + \varepsilon_{\Delta b})$. Fourth, Assumption 4.4 (vi) and Lemma 2.12 with $X = \tilde{V}$ yield

$$\begin{aligned} \|(I - Q_h^V Q_h^{V*})u\|_{\tilde{V}} &\leq \|(I - Q_h^V I_h)u\|_{\tilde{V}} + c_V \|(Q_h^{V*} - I_h)u\|_{\tilde{a}_h} \\ &\leq (1 + c_V^2) \|(I - Q_h^V I_h)u\|_{\tilde{V}} + c_V \|\Delta \tilde{a}(I_h u)\|_{\tilde{V}_h^*}, \end{aligned}$$

which implies that the error of the reference solution $Q_h J_h x$ is bounded by

$$\|(I - Q_h J_h)x\|_X \leq \|(I - Q_h^V Q_h^{V*})u\|_{\tilde{V}} + \|(I - Q_h^V I_h)u'\|_H \leq C(\varepsilon_{\text{ip}} + \varepsilon_{\text{forms}}).$$

We obtain the final estimate after collecting terms. \square

REMARK 4.9 In some situations, it is more practical to further estimate some terms of the error bound.

- (i) To compare the discrete data with the interpolated exact data (instead of $Q_h^{V*} u_1^0$ and $Q_h^{H*} f$ as in $\varepsilon_{\text{data}}$), we apply Lemma 2.12 with $X_h = \tilde{V}_h$ which shows

$$\begin{aligned} \|u_{h,1}^0 - Q_h^{V*} u_1^0\|_{\tilde{a}_h} &\leq \|u_{h,1}^0 - I_h u_1^0\|_{\tilde{a}_h} + \|(I_h - Q_h^{V*})u_1^0\|_{\tilde{a}_h} \\ &\leq \|u_{h,1}^0 - I_h u_1^0\|_{\tilde{a}_h} + c_V \|(I - Q_h^V I_h)u_1^0\|_{\tilde{V}} + \|\Delta \tilde{a}(I_h u_1^0)\|_{\tilde{V}_h^*}, \quad u_1^0 \in Z_{\text{ip}}. \end{aligned}$$

Analogously, we obtain

$$\|f_h - Q_h^{H*} f\|_{\infty, H_h} \leq \|f_h - I_h f\|_{\infty, H_h} + c_H \|(I - Q_h^V I_h)f\|_{\infty, H} + \|\Delta m(I_h f)\|_{\infty, H_h^*},$$

if $f(t) \in Z_{\text{ip}}$ for $t \in [0, T]$.

- (ii) If $\mathcal{B} \in \mathcal{L}(\tilde{V}, H)$ then $\varepsilon_{\Delta b}$ is bounded by an interpolation and a geometric error which contains

$$\Delta b(w_h, v_h) := b(Q_h^V w_h, Q_h^V v_h) - b_h(w_h, v_h).$$

To see this, we apply Assumption 4.4 (v) and obtain for $u' \in \tilde{V}$ and $v_h \in V_h$

$$\begin{aligned} |b(u', Q_h^V v_h) - b_h(I_h u', v_h)| &\leq |b((I - Q_h^V I_h)u', Q_h^V v_h)| + |b(Q_h^V I_h u', Q_h^V v_h) - b_h(I_h u', v_h)| \\ &\leq |\langle \mathcal{B}(I - Q_h^V I_h)u', Q_h^V v_h \rangle_V| + |\Delta b(I_h u', v_h)| \\ &\leq \|\mathcal{B}\|_{H \leftarrow \tilde{V}} \|(I - Q_h^V I_h)u'\|_{\tilde{V}} \|Q_h^V v_h\|_H + |\Delta b(I_h u', v_h)| \\ &\leq \|\mathcal{B}\|_{H \leftarrow \tilde{V}} \|(I - Q_h^V I_h)u'\|_{\tilde{V}} c_H \|v_h\|_{m_h} + |\Delta b(I_h u', v_h)|. \end{aligned}$$

Therefore, $\varepsilon_{\Delta b}$ is bounded by

$$\varepsilon_{\Delta b} \leq C \left(\|(I - Q_h^V I_h)u'\|_{\infty, \tilde{V}} + \|\Delta b(I_h u')\|_{\infty, H_h^*} \right).$$

By Definition 2.5 the discretization (4.8) is conforming, if

$$V_h \subset V, \quad Q_h^V = I, \quad \Delta m = 0, \quad \Delta b = 0, \quad \Delta a = 0.$$

For conforming discretizations we state an error bound which is independent of u'' .

COROLLARY 4.10. *Let (4.8) be a conforming discretization and consider the situation from Theorem 4.8. Then the semi-discrete solution u_h satisfies*

$$\|u_h(t) - u(t)\|_{\tilde{V}} + \|u_h'(t) - u'(t)\|_H \leq C e^{c_{\text{qm}} t} (1+t) \left(\tilde{\varepsilon}_{\text{data}} + \tilde{\varepsilon}_{\text{ip}} + \tilde{\varepsilon}_b \right)$$

for $t \in [0, T]$, where C is independent of h and t , $c_{\text{qm}} = c_G C_{H,V} / 2 + \rho_{\text{qm}}$,

$$\begin{aligned} \tilde{\varepsilon}_{\text{data}} &:= \|u_{h,1}^0 - \Pi_h^V u_1^0\|_{\tilde{a}_h} + \|u_{h,2}^0 - \Pi_h^H u_2^0\|_{m_h} + \|f_h - \Pi_h^H f\|_{\infty, H_h}, \\ \tilde{\varepsilon}_{\text{ip}} &:= \|(I - I_h)u\|_{\infty, \tilde{V}} + \varepsilon_h^{-1} \|(I - I_h)u\|_{\infty, H} + \|(I - I_h)u'\|_{\infty, \tilde{V}} + \varepsilon_h^{-1} \|(I - I_h)u'\|_{\infty, H}, \\ \tilde{\varepsilon}_b &:= \|b((I - \Pi_h^H)u')\|_{\infty, H_h^*}, \end{aligned}$$

where ε_h is defined in (4.12).

Proof. In comparison to the previous proof, there are only three changes. First, we have $\hat{c}_{\text{qm}} = c_{\text{qm}}$, since $\Delta m = \Delta b = \Delta a = 0$. Second, we choose $J_h = \Pi_h = Q_h^*$ so that Remark 2.10 applies. Third, since $Q_h^{H^*} = \Pi_h^H$ and $Q_h^{V^*} = \Pi_h^V$, the estimate from Lemma 4.7 reads

$$\begin{aligned} \|R_h x\|_{X_h} &\leq C \left(\|(I - \Pi_h^H)u\|_{\tilde{V}} + \|(I - \Pi_h^H)u'\|_{\tilde{V}} + \max_{\|v_h\|_{m_h}=1} |b(u', v_h) - b_h(\Pi_h^H u', v_h)| \right) \\ &\leq C \left(\|(I - \Pi_h^H)u\|_{\tilde{V}} + \|(I - \Pi_h^H)u'\|_{\tilde{V}} + \max_{\|v_h\|_{m_h}=1} |b((I - \Pi_h^H)u', v_h)| \right). \end{aligned}$$

To further bound the two terms with H -orthogonal projection errors in the \tilde{V} -norm, we use (3.7) with $X = H$ and $Y = \tilde{V}$ and the best approximation property of Π_h^H . This yields

$$\|(I - \Pi_h^H)w\|_{\tilde{V}} \leq \|(I - I_h)w\|_{\tilde{V}} + 2\varepsilon_h^{-1} \|(I - I_h)w\|_H, \quad w \in Z_{\text{ip}}.$$

We apply this estimate for $w = u$ and $w = u'$ in the above bound of the remainder term. For the final bound, we estimate the orthogonal projection error $\|(I - \Pi_h)x\|_X$ by interpolation errors and collect terms. \square

4.5 Example: Finite elements for the acoustic wave equation

In this example, we consider the acoustic wave equation with homogeneous Dirichlet boundary conditions and its space discretization using linear Lagrange finite elements with mass lumping, cf. (Cohen, 2002, Chapters 11–13). The aim is to show that our general analysis provides the same order of convergence as in the literature (Dupont (1973), Baker (1976), and Baker & Dougalis (1976)). However, our

analysis allows to account for errors resulting from numerical quadrature for mass lumping in a simple way.

We seek the solution $u : [0, T] \times \Omega \rightarrow \mathbb{R}$ of

$$u_{tt} - \operatorname{div}(c_\Omega \nabla u) = f \quad \text{in } \Omega, \quad (4.15a)$$

$$u(t) = 0 \quad \text{on } \Gamma, \quad (4.15b)$$

$$u(0) = u_1^0, \quad u_t(0) = u_2^0 \quad \text{in } \Omega. \quad (4.15c)$$

Here, f is a given source term, $c_\Omega \in L^\infty(\Omega)^{d \times d}$ models the wave speed, and Ω is polygonal. We assume that $c_\Omega(x)$, $x \in \Omega$ is symmetric, and that there are constants $c_\Omega^+ \geq c_\Omega^- > 0$ s.t.

$$c_\Omega^- \|\xi\|^2 \leq c_\Omega(x) \xi \cdot \xi \leq c_\Omega^+ \|\xi\|^2 \quad \text{for a.e. } x \in \Omega \text{ and all } \xi \in \mathbb{R}^d.$$

We can write the variational formulation of (4.15) in the form of (4.1) by making the following identifications. For the function spaces we set $V = H_0^1(\Omega)$ and $H = L^2(\Omega)$. As usual, m is the standard $L^2(\Omega)$ inner product, the bilinear form a is given by

$$a(u, v) := \int_\Omega c_\Omega \nabla u \cdot \nabla v \, dx, \quad u, v \in V,$$

and b vanishes. Due to the Poincaré inequality, Assumption 4.1 holds with $c_G = 0$ and we have $\tilde{a} = a$. Hence we can apply Theorem 4.3 which yields for suitable u_1^0 , u_2^0 , and f the existence of a unique solution of (4.15) with

$$u \in C^2([0, T]; L^2(\Omega)) \cap C^1([0, T]; H_0^1(\Omega)) \cap C([0, T]; [D(A)]),$$

where

$$D(A) = \left\{ u \in H_0^1(\Omega) \mid \operatorname{div}(c_\Omega \nabla u) \in L^2(\Omega) \right\}.$$

For the spatial discretization we restrict us to linear finite elements for this exposition. However, higher order elements can also be treated without further difficulties. Assume that the mesh \mathcal{T}_h is a triangulation of Ω s.t. the computational domain satisfies $\Omega_h = \Omega$. Then the space of linear finite elements V_h on \mathcal{T}_h is a subspace of V and the lift operator $Q_h^V = \mathbf{I}$ is trivial.

First, we study finite elements with exact integration. This means that we choose $m_h = m$ and $a_h = a$. Hence Assumption 4.4 holds trivially since $\Delta m = \Delta \tilde{a} = 0$ and Corollary 4.10 implies

$$\begin{aligned} & \|u_h(t) - u(t)\|_{\tilde{V}} + \|u_h'(t) - u'(t)\|_H \\ & \leq C(1+t) \left(\|(I - I_h)u\|_{\infty, \tilde{V}} + \varepsilon_h^{-1} \|(I - I_h)u\|_{\infty, H} + \|(I - I_h)u'\|_{\infty, \tilde{V}} + \varepsilon_h^{-1} \|(I - I_h)u'\|_{\infty, H} \right), \end{aligned}$$

if $\tilde{\varepsilon}_{\text{data}} = 0$. To obtain a convergence rate, we use that the error of the nodal interpolation operator I_h is bounded by

$$\|(I - I_h)\varphi\|_H + h\|(I - I_h)\varphi\|_{\tilde{V}} \leq Ch^2 |\varphi|_{H^2(\Omega)}, \quad \varphi \in H^2(\Omega),$$

cf. (Brenner & Scott, 2008, Sect. 4.4), and that $\varepsilon_h^{-1} \leq Ch^{-1}$ by inverse inequalities. Overall, we find that the difference in the energy norm between the exact solution u of (4.15) and its corresponding FEM approximation u_h scales like h , if $u \in C^1([0, T]; H^2(\Omega))$.

We next study the effect of numerical integration. The main difference to exact integration is, that now m_h and a_h differ from m and a , respectively. In this case, we apply Theorem 4.8 which yields

$$\begin{aligned} & \|u_h(t) - u(t)\|_{\tilde{V}} + \|u'_h(t) - u'(t)\|_H \\ & \leq C(1+t) \left(\|(I - I_h)u\|_{\infty, \tilde{V}} + \|(I - I_h)u'\|_{\infty, \tilde{V}} + \|(I - I_h)u''\|_{\infty, H} \right. \\ & \quad \left. + \|\Delta \tilde{a}(I_h u)\|_{\infty, \tilde{V}_h^*} + \|\Delta m(I_h u)\|_{\infty, H_h^*} + \|\Delta \tilde{a}(I_h u')\|_{\infty, \tilde{V}_h^*} + \|\Delta m(I_h u'')\|_{\infty, H_h^*} \right), \end{aligned}$$

if $\varepsilon_{\text{data}} = 0$. Hence it remains to quantify the differences in the bilinear form. For example in the above setting one can use the d -dimensional trapezoidal rule to approximate the integrals. More precisely, let $\{x_{K,j}\}_{j=1}^{d+1}$ be vertices of the element $K \in \mathcal{T}_h$. Then m_h and a_h are given by the quadrature formulas

$$m_h(v, w) = \sum_{K \in \mathcal{T}_h} \sum_{j=1}^{d+1} \frac{|K|}{d+1} v(x_{K,j}) w(x_{K,j})$$

and respectively

$$\tilde{a}_h(v, w) = \sum_{K \in \mathcal{T}_h} \sum_{j=1}^{d+1} \frac{|K|}{d+1} c_{\Omega}(x_{K,j}) \nabla v(x_{K,j}) \cdot \nabla w(x_{K,j}).$$

Under appropriate regularity assumptions on the wave speed c_{Ω} , it is known that $\|\Delta \tilde{a}(v_h)\|_{\tilde{V}_h^*} \in \mathcal{O}(h)$ and $\|\Delta m(v_h)\|_{H_h^*} \in \mathcal{O}(h)$ for all $v_h \in V_h$, see e.g. (Ciarlet, 2002, Section 4.1). Inserting this into the a priori bound above shows that numerical quadrature does not lead to order reduction, if $u \in C^2([0, T]; H^2(\Omega))$.

5. Application: Finite elements for the wave equation with acoustic boundary conditions

In this section, we use the a priori error bound from Theorem 4.8 to show new convergence rates for an isoparametric bulk-surface finite element discretization of the wave equation with acoustic boundary conditions while factoring in non-conforming error sources due to domain approximation.

We are interested in the solutions of the wave equation with acoustic boundary conditions. It models the propagation of sound waves in a fluid at rest filling a tank Ω , whose walls on Γ are subject to small oscillations in normal direction and elastic effects in tangential direction. Here u describes the acoustic velocity potential and δ the displacement of Γ in normal direction. The first well-posedness analysis was already given in Beale (1976), but acoustic boundary conditions continue to be a topic of research, see e.g. Gal *et al.* (2003), Mugnolo (2006), Frota *et al.* (2011), Graber (2012) and Vedurmundi *et al.* (2016).

For the space discretization we consider an isoparametric bulk-surface finite element method. Such finite element methods are non-conforming, since the computational domain is in general not exact, i.e. $\Omega_h \neq \Omega$. Therefore the error analysis requires a non-trivial lift operation. To the best of our knowledge, such an analysis has not been considered so far for hyperbolic problems. For parabolic problems, it was recently presented in Kovács & Lubich (2017). Our general framework and our abstract results allows us to derive convergence rates almost as easy as in the previous example for a conforming discretization by using the a priori estimate from Theorem 4.8.

The problem can be stated as follows. Let $\Omega \subset \mathbb{R}^d$, where $d = 2$ or $d = 3$ and assume that $\Gamma \in C^{k+1}$

for some $k \in \mathbb{N}$. We seek $u: [0, T] \times \Omega \rightarrow \mathbb{R}$ and $\delta: [0, T] \times \Gamma \rightarrow \mathbb{R}$ s.t.

$$u_{tt} + a_\Omega u - c_\Omega \Delta u = f_\Omega \quad \text{in } \Omega, \quad (5.1a)$$

$$\mu_\Gamma \delta_{tt} + k_\Gamma \delta - c_\Gamma \Delta_\Gamma \delta + c_\Omega u_t = f_\Gamma \quad \text{on } \Gamma, \quad (5.1b)$$

$$\delta_t = \partial_n u \quad \text{on } \Gamma, \quad (5.1c)$$

where we assume that $c_\Gamma, c_\Omega, \mu_\Gamma > 0$ and $a_\Omega, k_\Gamma \geq 0$ are constants and that u and δ take initial values $u(0) = u_1^0$, $u_t(0) = u_2^0$, $\delta(0) = \delta^0$, $\delta_t(0) = \vartheta^0$. By Δ_Γ we denote the Laplace–Beltrami operator.

Well-posedness

Assume that u and δ are sufficiently smooth solutions of (5.1). Multiplying (5.1a) with $v \in C^\infty(\overline{\Omega})$, integrating over Ω , applying Gauss' Theorem and inserting the boundary condition (5.1c) yields

$$\int_\Omega u_{tt} v + a_\Omega u v + c_\Omega \nabla u \cdot \nabla v \, dx - \int_\Gamma c_\Omega \delta_t v \, ds = \int_\Omega f_\Omega v \, dx. \quad (5.2a)$$

Analogously, we multiply (5.1b) with $\vartheta \in C^2(\Gamma)$, integrate over Γ , and use Gauss' Theorem on surfaces to find

$$\int_\Gamma \mu_\Gamma \delta_{tt} \vartheta + k_\Gamma \delta \vartheta + c_\Gamma \nabla_\Gamma \delta \cdot \nabla_\Gamma \vartheta + c_\Omega u_t \vartheta \, ds = \int_\Gamma f_\Gamma \vartheta \, ds. \quad (5.2b)$$

To obtain the variational problem for $\vec{u}(t) := (u(t), \delta(t))$, we add (5.2b) to (5.2a). Altogether, we showed that any classical solution $\vec{u} \in C^2(\overline{\Omega} \times [0, T]) \times C^2(\Gamma \times [0, T])$ of (5.1) satisfies the variational problem

$$m(\vec{u}''(t), \vec{v}) + b(\vec{u}'(t), \vec{v}) + a(\vec{u}(t), \vec{v}) = \langle f(t), \vec{v} \rangle \quad (5.3)$$

for all $\vec{v} = (v, \vartheta) \in C^\infty(\overline{\Omega}) \times C^2(\Gamma)$ and where, for $\vec{w} = (w, \omega)$ and $\vec{v} = (v, \vartheta)$, the bilinear forms are given by

$$\begin{aligned} m(\vec{w}, \vec{v}) &:= \int_\Omega w v \, dx + \int_\Gamma \mu_\Gamma \omega \vartheta \, ds, \\ b(\vec{w}, \vec{v}) &:= c_\Omega \int_\Gamma \gamma(w) \vartheta - \omega \gamma(v) \, ds, \\ a(\vec{w}, \vec{v}) &:= \int_\Omega a_\Omega w v + c_\Omega \nabla w \cdot \nabla v \, dx + \int_\Gamma k_\Gamma \omega \vartheta + c_\Gamma \nabla_\Gamma \omega \cdot \nabla_\Gamma \vartheta \, ds, \\ \langle f(t), \vec{\varphi} \rangle &:= \int_\Omega f_\Omega(t) v \, dx + \int_\Gamma f_\Gamma(t) \vartheta \, ds, \quad t \in [0, T]. \end{aligned}$$

For the corresponding abstract formulation, we choose $H = \mathbb{H}^0$ and $V = \mathbb{H}^1$, where

$$\mathbb{H}^0 := L^2(\Omega) \times L^2(\Gamma), \quad \text{and} \quad \mathbb{H}^r := H^r(\Omega) \times H^r(\Gamma), \quad r \in \mathbb{N},$$

and consider the continuous extensions of m and a , b to $H \times H$ and $V \times V$, respectively. Since V is a dense subspace of H and Assumption 4.1 is fulfilled with $c_G = \alpha = \min\{c_\Omega, c_\Gamma\} > 0$ and $\rho_{\text{qm}} = 0$, the abstract interpretation of (5.3) is a second-order wave-type equation (4.1). Thus, if the data satisfies

$$\begin{aligned} &u_1^0, u_2^0 \in V \quad \text{s.t.} \quad \mathcal{A}u_1^0 + \mathcal{B}u_2^0 \in H \\ \text{and} \quad &f \in C^1([0, T]; H) \quad \text{or} \quad (f, \mathcal{B}f) \in C([0, T]; V \times H), \end{aligned}$$

then Theorem 4.3 implies that (4.1) has a unique solution \vec{u} .

The following Lemma states the well-posedness result in terms of Sobolev spaces. Since our convergence result for finite elements will require higher regularity, we do not use it in the error analysis and refer to (Hipp, 2017, Cor. 6.9 (i)) for the proof. A related result can be found in Beale (1976).

LEMMA 5.1. *Let the above assumptions be satisfied. If the initial values $(u_1^0, \delta^0), (u_2^0, \vartheta^0) \in \mathbb{H}^1$ satisfy $(\Delta u_1^0, \Delta_\Gamma \delta^0) \in \mathbb{H}^0$ and $\vartheta^0 = \partial_n u_1^0$, and $(f_\Omega, f_\Gamma) \in C^1([0, T]; \mathbb{H}^0)$ or $f_\Omega \in C([0, T]; H^1(\Omega))$ with $f_\Gamma = 0$, then (5.1) has a unique solution*

$$(u, \delta) \in C^2([0, T]; \mathbb{H}^0) \cap C^1([0, T]; \mathbb{H}^1), \quad (\Delta u, \Delta_\Gamma \delta) \in C([0, T]; \mathbb{H}^0).$$

The bulk-surface finite element method

In this section, we consider the bulk-surface finite element method from Elliott & Ranner (2013) which was developed and analyzed for coupled bulk-surface partial differential equations of elliptic type.

COMPUTATIONAL DOMAIN Let \mathcal{T}_h be a mesh consisting of isoparametric elements K of degree p , where h denotes the mesh parameter, see (Elliott & Ranner, 2013, Sect. 4.1.2) for details on the construction. We denote the computational domain by

$$\Omega_h := \bigcup_{K \in \mathcal{T}_h} K \approx \Omega$$

and refer to $\Gamma_h := \partial\Omega_h$ as the computational surface. The construction admits quasi-uniform triangulations \mathcal{T}_h and $\mathcal{T}_h|_{\Gamma_h}$ of Ω_h and Γ_h , respectively.

FINITE ELEMENT SPACES Let $\mathcal{P}_p(\widehat{K})$ denote the space of polynomials of degree p on the reference triangle \widehat{K} , and let F_K be the transformation from \widehat{K} to $K \in \mathcal{T}_h$. For the bulk and the surface finite element functions of degree $p \geq 1$, we introduce

$$\begin{aligned} V_{h,p}^\Omega &:= \left\{ v_h \in C(\Omega_h) \mid v_h|_K = \widehat{v}_h \circ (F_K)^{-1} \text{ with } \widehat{v}_h \in \mathcal{P}_p(\widehat{K}) \text{ for all } K \in \mathcal{T}_h \right\}, \\ V_{h,p}^\Gamma &:= \left\{ \vartheta_h \in C(\Gamma_h) \mid \vartheta_h = v_h|_{\Gamma_h}, v_h \in V_{h,p}^\Omega \right\}, \end{aligned}$$

respectively, cf. (Elliott & Ranner, 2013, Sect. 5.1). An important property of this construction is the relation

$$\gamma(V_{h,p}^\Omega) = V_{h,p}^\Gamma. \quad (5.4)$$

LIFT OPERATION In general, the finite element approximation is defined in the computational domain $\Omega_h \neq \Omega$ and its boundary $\Gamma_h \neq \Gamma$. To compare it with the exact solution, we transform the finite element solution s.t. it is defined on Ω and Γ , respectively. A major advantage of this approach over extension or restriction strategies is that it does not depend on whether or not $\Omega_h \subset \Omega$.

The transformation is done via the elementwise smooth homeomorphism G_h from (Elliott & Ranner, 2013, Sect. 4.2) with

$$G_h: \Omega_h \rightarrow \Omega, \quad G_h|_K \in C^{p+1}(K) \quad \text{for } p \leq k \text{ and } K \in \mathcal{T}_h.$$

We shortly sketch its construction. The authors start from a base triangulation of Ω consisting of triangles only. It is assumed that each element has at most one boundary face and that all boundary faces lie within a band about Γ s.t. each point x in this band has a unique normal projection $p(x) \in \Gamma$. Note that such a band always exists since $\Gamma \in C^2$. The isoparametric mesh \mathcal{T}_h is then obtained by transforming each triangle from the boundary layer s.t. its boundary face interpolates the corresponding segment of Γ with order p . While $G_h = I$ in the interior elements, boundary elements are handled in two steps. First, they are mapped to their counterparts in the base triangulation. Then, by a smooth extension of the normal projection p , these elements are transformed s.t. their boundary face lies on Γ while their other faces remain invariant. For details we refer to (Elliott & Ranner, 2013, Sect. 4).

Given $v_h \in V_{h,p}^\Omega$ and $\vartheta_h \in V_{h,p}^\Gamma$, we define their lifted counterparts as

$$v_h^\ell(x) := v_h(G_h^{-1}(x)), \quad x \in \Omega, \quad (5.5a)$$

$$\vartheta_h^\ell(x) := \vartheta_h(G_h^{-1}(x)), \quad x \in \Gamma. \quad (5.5b)$$

Note that the bulk lifting complies with the surface lifting in the sense that

$$\gamma(v_h^\ell) = \gamma(v_h)^\ell, \quad v_h \in V_{h,p}^\Omega, \quad (5.6)$$

where we overload the notation with $\gamma: H^1(\Omega) \rightarrow L^2(\Gamma)$ on the left-hand side and $\gamma: H^1(\Omega_h) \rightarrow L^2(\Gamma_h)$ on the right-hand side.

REMARK 5.2 Actually, our definition of lifted surface functions differs from (Elliott & Ranner, 2013, Def. 4.12) where a closest point mapping from Γ_h to Γ is used. However, it follows from Demlow (2009) that the surface functions lifted by (5.5b) have the same properties and satisfy (5.6) in addition.

INTERPOLATION As the exact solution has two components u and δ , we introduce two interpolation operators. The nodal interpolation operator $I_h^\Omega: H^2(\Omega) \rightarrow V_{h,p}^\Omega$ for bulk functions $v \in H^{r+1}(\Omega)$ satisfies

$$\|v - (I_h^\Omega v)^\ell\|_{L^2(\Omega)} + h\|v - (I_h^\Omega v)^\ell\|_{H^1(\Omega)} \leq Ch^{r+1}|v|_{H^{r+1}(\Omega)}, \quad 1 \leq r \leq p. \quad (5.7a)$$

Analogously, we write $I_h^\Gamma: H^2(\Gamma) \rightarrow V_{h,p}^\Gamma$ for the nodal interpolation on the surface and the interpolation error of a function $\vartheta \in H^{r+1}(\Gamma)$ is bounded by

$$\|\vartheta - (I_h^\Gamma \vartheta)^\ell\|_{L^2(\Gamma)} + h\|\vartheta - (I_h^\Gamma \vartheta)^\ell\|_{H^1(\Gamma)} \leq Ch^{r+1}|\vartheta|_{H^{r+1}(\Gamma)}, \quad 1 \leq r \leq \min\{p, k\}. \quad (5.7b)$$

Since the nodes in the bulk and on the surface coincide by construction, it follows from (5.4) that we have $I_h^\Gamma \gamma(v) = \gamma(I_h^\Omega v)$ for any $v \in H^2(\Omega)$ with $\gamma(v) \in H^2(\Gamma)$.

A priori error bounds for the wave equation with acoustic boundary conditions

Applying the bulk-surface finite element method to (5.1) yields a differential equation of the form (4.8). The semi-discrete problem is to find the $V_h := V_{h,p}^\Omega \times V_{h,p}^\Gamma$ -valued function $\vec{u}_h := (u_h, \delta_h): [0, T] \rightarrow V_h$ s.t.

$$\begin{aligned} m_h(u_h''(t), \vec{v}_h) + b_h(u_h'(t), \vec{v}_h) + a(u_h(t), \vec{v}_h) &= m_h((I_h^\Omega f_\Omega(t), I_h^\Gamma f_\Gamma(t)), \vec{v}_h) \quad \forall \vec{v}_h \in V_h, \\ \vec{u}_h(0) &= (I_h^\Omega u_1^0, I_h^\Gamma \delta^0), \quad \vec{u}_h'(0) = (I_h^\Omega u_2^0, I_h^\Gamma \vartheta^0), \end{aligned}$$

where, for $\vec{w}_h = (w_h, \omega_h)$, $\vec{v}_h = (v_h, \vartheta_h) \in V_{h,p}^\Omega \times V_{h,p}^\Gamma$, the bilinear forms are defined as

$$\begin{aligned} m_h(\vec{w}_h, \vec{v}_h) &:= \int_{\Omega_h} w_h v_h \, dx + \int_{\Gamma_h} \mu_\Gamma \omega_h \vartheta_h \, ds, \\ b_h(\vec{w}_h, \vec{v}_h) &:= c_\Omega \int_{\Gamma_h} \gamma(w_h) \vartheta_h - \omega_h \gamma(v_h) \, ds, \\ a_h(\vec{w}_h, \vec{v}_h) &:= \int_{\Omega_h} a_\Omega w_h v_h + c_\Omega \nabla w_h \cdot \nabla v_h \, dx + \int_{\Gamma_h} k_\Gamma \omega_h \vartheta_h + c_\Gamma \nabla_{\Gamma_h} \omega_h \cdot \nabla_{\Gamma_h} \vartheta_h \, ds. \end{aligned}$$

Due to the abstract error bound from Theorem 4.8, we can derive convergence rates for the above discretization in a few simple steps. Basically, we only have to insert the approximation properties of the finite element method.

THEOREM 5.3. *Let $\Gamma \in C^{k+1}$ for some $k \in \mathbb{N}$ and $(f_\Omega(t), f_\Gamma(t)) \in \mathbb{H}^2$, $t \in [0, T]$. Let the assumptions of Lemma 5.1 be fulfilled and assume that the solution of the wave equation with acoustic boundary conditions (5.1) satisfies $(u, \delta) \in C^2([0, T]; \mathbb{H}^2)$. Moreover, let $\vec{u}_h = (u_h, \delta_h)$ be the finite element solution as described above where $1 \leq p \leq k$ and $0 < h \leq 1$. Then the error of lifted finite element approximation $(u_h^\ell, \delta_h^\ell)$ is bounded by*

$$\|(u_h^\ell - u, \delta_h^\ell - \delta)(t)\|_{\mathbb{H}^1} + \|((u_h^\ell)^\ell - u', (\delta_h^\ell)^\ell - \delta')(t)\|_{\mathbb{H}^0} \leq CK_p(u, f) e^{\widehat{c}_{\text{qm}} t} (1+t) h^p$$

for $t \in [0, T]$, a constant C which is independent of h and t , $\widehat{c}_{\text{qm}} = \min\{c_\Omega, c_\Gamma\}^{1/2}/2$ and

$$K_p(u, f) = \|(u, \delta)\|_{\infty, \mathbb{H}^{p+1}} + \|(u', \delta')\|_{\infty, \mathbb{H}^{p+1}} + \|\gamma(u')\|_{\infty, H^2(\Gamma)} + \|(u'', \delta'')\|_{\infty, \mathbb{H}^q} + \|(f_\Omega, f_\Gamma)\|_{\infty, \mathbb{H}^q}$$

for $q = \max\{2, p\}$.

REMARK 5.4 Note that the conditions from Lemma 5.1 on Ω and the data are in general not sufficient for $(u, \delta) \in C^2([0, T]; \mathbb{H}^2)$ and the proof requires further considerations, cf. (Beale, 1976, Thm. 2.2).

Proof. Assumptions 4.4 (i)-4.4 (iv) are fulfilled with $\widehat{c}_G = \min\{c_\Omega, c_\Gamma\}$ and $\widehat{\rho}_{\text{qm}} = 0$ as in the continuous case. To compare the finite element approximation with the exact solution, we choose the lift operator

$$Q_h^V \vec{v}_h := (v_h^\ell, \vartheta_h^\ell). \quad (5.8)$$

Since the coefficients in $\|\cdot\|_H$ and $\|\cdot\|_{\widehat{V}}$ are constant, (Elliott & Ranner, 2013, Prop. 4.9 and 4.13) imply that $Q_h^V: V_h \rightarrow V$ satisfies Assumptions 4.4 (v) and 4.4 (vi). Altogether the space discretization is stable in the sense of Assumption 4.4 and the error estimate from Theorem 4.8 applies. Since $\|\cdot\|_{\mathbb{H}^0} \sim \|\cdot\|_H$ and $\|\cdot\|_{\mathbb{H}^1} \sim \|\cdot\|_{\widehat{V}}$, it remains to bound $\varepsilon_{\text{data}} + \varepsilon_{\text{ip}} + \varepsilon_{\text{forms}} + \varepsilon_{\Delta b}$.

(ε_{ip}) We choose the interpolation operator as $I_h := \text{diag}(I_h^\Omega, I_h^\Gamma)$ s.t. $I_h \in \mathcal{L}(Z_{\text{ip}}, V_h)$ for $Z_{\text{ip}} = \mathbb{H}^2$. Using (5.7), we have for $\vec{v} = (v, \vartheta) \in \mathbb{H}^{r+1}$ and $1 \leq r \leq p$

$$\begin{aligned} \|(I - Q_h^V I_h) \vec{v}\|_H + h \|(I - Q_h^V I_h) \vec{v}\|_{\widehat{V}} &\leq \|v - (I_h^\Omega v)^\ell\|_{L^2(\Omega)} + \mu_\Gamma \|\vartheta - (I_h^\Gamma \vartheta)^\ell\|_{L^2(\Gamma)} \\ &\quad + \max\left\{\sqrt{a_\Omega + \widehat{c}_G}, \sqrt{c_\Omega}\right\} h \|v - (I_h^\Omega v)^\ell\|_{H^1(\Omega)} \\ &\quad + \max\left\{\sqrt{k_\Gamma + \mu_\Gamma \widehat{c}_G}, \sqrt{c_\Gamma}\right\} h \|\vartheta - (I_h^\Gamma \vartheta)^\ell\|_{H^1(\Gamma)} \\ &\leq Ch^{r+1} |\vec{v}|_{H^{r+1}(\Omega) \times H^{r+1}(\Gamma)}. \end{aligned} \quad (5.9)$$

Applying this bound on the interpolation errors ε_{ip} , yields

$$\varepsilon_{\text{ip}} \leq Ch^p \left(\|(u, \delta)\|_{\infty, \mathbb{H}^{p+1}} + \|(u', \delta')\|_{\infty, \mathbb{H}^{p+1}} + \|(u'', \delta'')\|_{\infty, \mathbb{H}^q} \right) \leq Ch^p K_p(u, f).$$

($\varepsilon_{\text{forms}}$) To estimate the geometric errors, we use (Elliott & Ranner, 2013, Lem. 6.2) which yields

$$\begin{aligned} \left| \int_{\Omega} w_h^\ell v_h^\ell \, dx - \int_{\Omega_h} w_h v_h \, dx \right| &\leq Ch^p \|w_h\|_{L^2(\Omega_h)} \|v_h\|_{L^2(\Omega_h)}, \\ \left| \int_{\Omega} \nabla w_h^\ell \cdot \nabla v_h^\ell \, dx - \int_{\Omega_h} \nabla w_h \cdot \nabla v_h \, dx \right| &\leq Ch^p \|\nabla w_h\|_{L^2(\Omega_h)} \|\nabla v_h\|_{L^2(\Omega_h)}, \\ \left| \int_{\Gamma} \omega_h^\ell \vartheta_h^\ell \, dx - \int_{\Gamma_h} \omega_h \vartheta_h \, dx \right| &\leq Ch^{p+1} \|\omega_h\|_{L^2(\Gamma_h)} \|\vartheta_h\|_{L^2(\Gamma_h)}, \\ \left| \int_{\Gamma} \nabla_{\Gamma} \omega_h^\ell \cdot \nabla_{\Gamma} \vartheta_h^\ell \, dx - \int_{\Gamma_h} \nabla_{\Gamma_h} \omega_h \cdot \nabla_{\Gamma_h} \vartheta_h \, dx \right| &\leq Ch^{p+1} \|\nabla_{\Gamma_h} \omega_h\|_{L^2(\Gamma_h)} \|\nabla_{\Gamma_h} \vartheta_h\|_{L^2(\Omega_h)}. \end{aligned} \quad (5.10)$$

Now consider $\Delta \tilde{a}$ and let $\vec{w}_h = (w_h, \omega_h)$, $\vec{v}_h = (v_h, \vartheta_h) \in V_{h,p}^{\Omega} \times V_{h,p}^{\Gamma}$. From the above geometric error bounds, we deduce

$$\begin{aligned} |\Delta \tilde{a}(\vec{w}_h, \vec{v}_h)| &\leq C \max\{c_{\Omega}, a_{\Gamma}, c_{\Gamma}, k_{\Gamma}\} \left(h^p \|w_h\|_{H^1(\Omega_h)} \|v_h\|_{H^1(\Omega_h)} + h^{p+1} \|\omega_h\|_{H^1(\Gamma_h)} \|\vartheta_h\|_{H^1(\Gamma_h)} \right) \\ &\leq Ch^p \|\vec{w}_h\|_{\tilde{a}_h} \|\vec{v}_h\|_{\tilde{a}_h}. \end{aligned}$$

Since $I_h \in \mathcal{L}(\mathbb{H}^2, \tilde{V}_h)$, we obtain

$$\|\Delta \tilde{a}(I_h \vec{w})\|_{\tilde{V}_h^*} = \max_{\|\vec{v}_h\|_{\tilde{a}_h} = 1} |\Delta \tilde{a}(I_h \vec{w}, \vec{v}_h)| \leq Ch^p \|\vec{w}\|_{\mathbb{H}^2}, \quad \vec{w} \in \mathbb{H}^2.$$

Analogously, it holds $\|\Delta m(I_h \vec{w})\|_{H_h^*} \leq Ch^p \|\vec{w}\|_{\mathbb{H}^2}$ for $\vec{w} \in \mathbb{H}^2$. Since $q, p+1 \geq 2$, we therefore showed that

$$\varepsilon_{\text{forms}} \leq Ch^p \left(\|(u, \delta)\|_{\infty, \mathbb{H}^2} + \|(u', \delta')\|_{\infty, \mathbb{H}^2} + \|(u'', \delta'')\|_{\infty, \mathbb{H}^2} \right) \leq Ch^p K_p(u, f).$$

($\varepsilon_{\text{data}}$) Since $u_{h,1}^0 = I_h u_1^0$ and $f_h = I_h f$, we can apply the estimates from Remark 4.9 (i) and obtain, together with the shown bounds for the interpolation and consistency errors, that

$$\varepsilon_{\text{data}} \leq Ch^p \left(\|u_1^0\|_{\mathbb{H}^q} + \|f\|_{\infty, \mathbb{H}^q} \right) \leq Ch^p K_p(u, f).$$

($\varepsilon_{\Delta b}$) First, note that b can be written as

$$b(\vec{w}, \vec{v}) = \frac{c_{\Omega}}{\mu_{\Gamma}} \left(m((0, \gamma(w)), (0, \vartheta)) - m((0, \omega), (0, \gamma(v))) \right). \quad (5.11)$$

Hence, we obtain with (5.6) and (5.8)

$$\begin{aligned} \frac{\mu_{\Gamma}}{c_{\Omega}} b(\vec{w}, \mathcal{Q}_h^V \vec{v}_h) &= m((0, \gamma(w)), (0, \vartheta_h^\ell)) - m((0, \omega), (0, \gamma(v_h^\ell))) \\ &= m((0, \gamma(w)), (0, \vartheta_h^\ell)) - m((0, \omega), (0, \gamma(v_h^\ell))) \\ &= m((0, \gamma(w)), \mathcal{Q}_h^V(0, \vartheta_h)) - m((0, \omega), \mathcal{Q}_h^V(0, \gamma(v_h))) \\ &= m_h(\mathcal{Q}_h^{H^*}(0, \gamma(w)), (0, \vartheta_h)) - m_h(\mathcal{Q}_h^{H^*}(0, \omega), (0, \gamma(v_h))). \end{aligned}$$

Since b_h satisfies a representation analogous to (5.11), we have with $\gamma(I_h^\Omega v) = I_h^\Gamma \gamma(v)$

$$\begin{aligned} b_h(I_h \vec{w}, \vec{v}_h) &= \frac{c_\Omega}{\mu_\Gamma} \left(m_h((0, \gamma(I_h^\Omega w)), (0, \vartheta_h)) - m_h((0, I_h^\Gamma \omega), (0, \gamma(v_h))) \right) \\ &= \frac{c_\Omega}{\mu_\Gamma} \left(m_h((0, I_h^\Gamma \gamma(w)), (0, \vartheta_h)) - m_h((0, I_h^\Gamma \omega), (0, \gamma(v_h))) \right) \\ &= \frac{c_\Omega}{\mu_\Gamma} \left(m_h(I_h(0, \gamma(w)), (0, \vartheta_h)) - m_h(I_h(0, \omega), (0, \gamma(v_h))) \right). \end{aligned}$$

Thus the difference in $\varepsilon_{\Delta b}$ is bounded by

$$\begin{aligned} &|b(\vec{w}, \mathcal{Q}_h^V \vec{v}_h) - b_h(I_h \vec{w}, \vec{v}_h)| \\ &= \left| \frac{c_\Omega}{\mu_\Gamma} \left(m_h((\mathcal{Q}_h^{H^*} - I_h)(0, \gamma(w)), (0, \vartheta_h)) - m_h((\mathcal{Q}_h^{H^*} - I_h)(0, \omega), (0, \gamma(v_h))) \right) \right| \\ &\leq \frac{c_\Omega}{\mu_\Gamma} \left(\|(\mathcal{Q}_h^{H^*} - I_h)(0, \gamma(w))\|_{m_h} \| (0, \vartheta_h) \|_{m_h} + \|(\mathcal{Q}_h^{H^*} - I_h)(0, \omega)\|_{m_h} \| (0, \gamma(v_h)) \|_{m_h} \right), \end{aligned}$$

where we applied the Cauchy–Schwarz inequality for m_h in the last step. To deal with the last term, we use the continuity of the trace operator and the inverse inequality from (Brenner & Scott, 2008, Lem. 4.5.3), which yields

$$\| (0, \gamma(v_h)) \|_{m_h} = \sqrt{\mu_\Gamma} \|\gamma(v_h)\|_{L^2(\Gamma_h)} \leq \|\gamma\|_{L^2(\Gamma_h) \leftarrow H^1(\Omega_h)} \|v_h\|_{H^1(\Omega_h)} \leq Ch^{-1} \|v_h\|_{L^2(\Omega_h)}.$$

Thus, we showed that

$$\begin{aligned} \varepsilon_{\Delta b} &= \max_{\|\vec{v}_h\|_{m_h}=1} |b(\vec{u}', \mathcal{Q}_h^V \vec{v}_h) - b_h(I_h \vec{u}', \vec{v}_h)| \\ &\leq C \left(\|(\mathcal{Q}_h^{H^*} - I_h)(0, \gamma(u'))\|_{m_h} + h^{-1} \|(\mathcal{Q}_h^{H^*} - I_h)(0, \delta')\|_{m_h} \right). \end{aligned} \quad (5.12)$$

Using Lemma 2.12 with $X = H$, the interpolation error estimate (5.9) and the geometric error bound (5.10), we are able to bound $\mathcal{Q}_h^{H^*} - I_h$ for surface functions $\omega \in H^{r+1}(\Gamma)$, $1 \leq r \leq p$ by

$$\begin{aligned} \|(\mathcal{Q}_h^{H^*} - I_h)(0, \omega)\|_{m_h} &\leq C(\|(\mathbf{I} - \mathcal{Q}_h^V I_h)(0, \omega)\|_{m_h} + \|\Delta m(I_h(0, \omega))\|_{H_h^*}) \\ &\leq C \left(h^{r+1} |\omega|_{H^{r+1}(\Gamma)} + h^{p+1} \|\omega\|_{H^2(\Gamma)} \right). \end{aligned}$$

If we insert this estimate (with $r = p - 1$, $\omega = \gamma(u')$ and $r = p$, $\omega = \delta'$) into (5.12), we obtain

$$\begin{aligned} \varepsilon_{\Delta b} &\leq Ch^p |\gamma(u')|_{H^p(\Gamma)} + Ch^{p+1} \|\gamma(u')\|_{H^2(\Gamma)} + Ch^{-1} \left(h^{p+1} |\delta'|_{H^{p+1}(\Gamma)} + h^{p+1} \|\delta'\|_{H^2(\Gamma)} \right) \\ &\leq Ch^p \|u'\|_{H^{p+1}(\Omega)} + Ch^{p+1} \|\gamma(u')\|_{H^2(\Gamma)} + Ch^p \|\delta'\|_{H^{p+1}(\Gamma)} \\ &\leq Ch^p K_p(u, f), \end{aligned}$$

since $\gamma \in \mathcal{L}(H^{r+1}(\Omega), H^r(\Gamma))$ for $r \geq 1$, cf. (Atkinson & Han, 2009, Thm. 7.3.11). This completes the proof. \square

ACKNOWLEDGEMENTS We thank Jan Leibold, Bernhard Maier and Roland Schnaubelt for their careful reading of this manuscript and helpful discussions.

FUNDING Deutsche Forschungsgemeinschaft (DFG) through the collaborative research center CRC 1173 »Wave phenomena: analysis and numerics«, Klaus Tschira Stiftung.

REFERENCES

- ATKINSON, K. & HAN, W. (2009) *Theoretical numerical analysis*. Texts in Applied Mathematics, vol. 39, third edn. Springer, Dordrecht, pp. xvi+625. A functional analysis framework.
- BAKER, G. A. (1976) Error estimates for finite element methods for second order hyperbolic equations. *SIAM J. Numer. Anal.*, **13**, 564–576.
- BAKER, G. A. & DOUGALIS, V. A. (1976) The effect of quadrature errors on finite element approximations for second order hyperbolic equations. *SIAM J. Numer. Anal.*, **13**, 577–598.
- BÁTKAI, A., CSOMÓS, P., FARKAS, B. & NICKEL, G. (2012) Operator splitting with spatial-temporal discretization. *Spectral theory, mathematical system theory, evolution equations, differential and difference equations*. Oper. Theory Adv. Appl., vol. 221. Birkhäuser/Springer Basel AG, Basel, pp. 161–171.
- BEALE, J. T. (1976) Spectral properties of an acoustic boundary condition. *Indiana Univ. Math. J.*, **25**, 895–917.
- BENZONI-GAVAGE, S. & SERRE, D. (2007) *Multidimensional hyperbolic partial differential equations*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, Oxford, pp. xxvi+508.
- BRENNER, P., CROUZEIX, M. & THOMÉE, V. (1982) Single-step methods for inhomogeneous linear differential equations in Banach space. *RAIRO Anal. Numér.*, **16**, 5–26.
- BRENNER, S. C. & SCOTT, L. R. (2008) *The mathematical theory of finite element methods*. Texts in Applied Mathematics, vol. 15, third edn. Springer, New York, pp. xviii+397.
- BURAZIN, K. S. & ERCEG, M. (2016) Non-stationary abstract Friedrichs systems. *Mediterr. J. Math.*, **13**, 3777–3796.
- BURMAN, E., ERN, A. & FERNÁNDEZ, M. A. (2010) Explicit Runge-Kutta schemes and finite elements with symmetric stabilization for first-order linear PDE systems. *SIAM J. Numer. Anal.*, **48**, 2019–2042.
- CIARLET, P. G. (2002) *The finite element method for elliptic problems*. Classics in Applied Mathematics, vol. 40. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, pp. xxviii+530.
- COCKBURN, B., QIU, W. & SOLANO, M. (2014) A priori error analysis for HDG methods using extensions from subdomains to achieve boundary conformity. *Math. Comp.*, **83**, 665–699.
- COHEN, G. C. (2002) *Higher-order numerical methods for transient wave equations*. Scientific Computation. Springer-Verlag, Berlin, pp. xviii+348.
- COHEN, G. & PERNET, S. (2017) *Finite element and discontinuous Galerkin methods for transient wave equations*. Scientific Computation. Springer, Dordrecht, pp. xvii+381. With a foreword by Patrick Joly.
- DEMLOW, A. (2009) Higher-order finite element methods and pointwise error estimates for elliptic problems on surfaces. *SIAM J. Numer. Anal.*, **47**, 805–827.
- DI PIETRO, D. A. & ERN, A. (2012) *Mathematical aspects of discontinuous Galerkin methods*. Mathématiques & Applications (Berlin) [Mathematics & Applications], vol. 69. Heidelberg: Springer, pp. xviii+384.
- DRONIOU, J., EYMARD, R., GALLOUËT, T., GUICHARD, C. & HERBIN, R. (2017) The gradient discretisation method. <https://hal.archives-ouvertes.fr/hal-01382358>.
- DUNCA, A. A. (2017) On an optimal finite element scheme for the advection equation. *J. Comput. Appl. Math.*, **311**, 522–528.
- DUPONT, T. (1973) L^2 -estimates for Galerkin methods for second order hyperbolic equations. *SIAM J. Numer. Anal.*, **10**, 880–889.
- ELLIOTT, C. M. & RANNER, T. (2013) Finite element analysis for a coupled bulk-surface partial differential equation. *IMA J. Numer. Anal.*, **33**, 377–402.
- ENGEL, K.-J. & NAGEL, R. (2000) *One-parameter semigroups for linear evolution equations*. Graduate Texts in Mathematics, vol. 194. Springer-Verlag, New York, pp. xxii+586. With contributions by S. Brendle, M. Campiti, T. Hahn, G. Metafuno, G. Nickel, D. Pallara, C. Perazzoli, A. Rhandi, S. Romanelli and R.

Schnaubelt.

- FEZOU, L., LANteri, S., LOHRENGEL, S. & PIPERNO, S. (2005) Convergence and stability of a discontinuous Galerkin time-domain method for the 3D heterogeneous Maxwell equations on unstructured meshes. *M2AN Math. Model. Numer. Anal.*, **39**, 1149–1176.
- FROTA, C. L., MEDEIROS, L. A. & VICENTE, A. (2011) Wave equation in domains with non-locally reacting boundary. *Differential Integral Equations*, **24**, 1001–1020.
- FUJITA, H., SAITO, N. & SUZUKI, T. (2001) *Operator theory and numerical methods*. Studies in Mathematics and its Applications, vol. 30. North-Holland Publishing Co., Amsterdam, pp. viii+309.
- GAL, C. G., GOLDSTEIN, G. R. & GOLDSTEIN, J. A. (2003) Oscillatory boundary conditions for acoustic wave equations. *J. Evol. Equ.*, **3**, 623–635.
- GRABER, P. J. (2012) The Wave Equation with Generalized Nonlinear Acoustic Boundary Conditions. *Ph.D. thesis*, University of Virginia.
- GROTE, M. J., SCHNEEBELI, A. & SCHÖTZAU, D. (2006) Discontinuous Galerkin finite element method for the wave equation. *SIAM J. Numer. Anal.*, **44**, 2408–2431.
- GUIDETTI, D., KARASÖZEN, B. & PISKAREV, S. (2004) Approximation of abstract differential equations. *J. Math. Sci. (N. Y.)*, **122**, 3013–3054. Functional analysis.
- HIPPE, D. (2017) A unified error analysis for spatial discretizations of wave-type equations with applications to dynamic boundary conditions. *Ph.D. thesis*, Karlsruhe Institute of Technology.
- HOCHBRUCK, M., JAHNKE, T. & SCHNAUBELT, R. (2015) Convergence of an ADI splitting for Maxwell's equations. *Numer. Math.*, **129**, 535–561.
- HOCHBRUCK, M., MAIER, B. & STÖHRER, C. (2017) Heterogeneous multiscale method for Maxwell's equations. To be published.
- HOCHBRUCK, M. & PAŽUR, T. (2015) Implicit Runge-Kutta methods and discontinuous Galerkin discretizations for linear Maxwell's equations. *SIAM J. Numer. Anal.*, **53**, 485–507.
- HOCHBRUCK, M. & STURM, A. (2016) Error analysis of a second-order locally implicit method for linear Maxwell's equations. *SIAM J. Numer. Anal.*, **54**, 3167–3191.
- HUSSEIN, A. E.-R. A. & KAPPEL, F. (2002) Weak stability and the Trotter-Kato theorem. *Ital. J. Pure Appl. Math.*, 147–158.
- ITO, K. & KAPPEL, F. (2002) *Evolution equations and approximations*. Series on Advances in Mathematics for Applied Sciences, vol. 61. World Scientific Publishing Co., Inc., River Edge, NJ, pp. xiv+498.
- JOLY, P. (2003) Variational methods for time-dependent wave propagation problems. *Topics in computational wave propagation*. Lect. Notes Comput. Sci. Eng., vol. 31. Springer, Berlin, pp. 201–264.
- KIRSCH, A. & HETTLICH, F. (2015) *The mathematical theory of time-harmonic Maxwell's equations*. Applied Mathematical Sciences, vol. 190. Springer, Cham, pp. xiv+337. Expansion-, integral-, and variational methods.
- KOVÁCS, B. & LUBICH, C. (2017) Numerical analysis of parabolic problems with dynamic boundary conditions. *IMA J. Numer. Anal.*, **37**, 1–39.
- LAYTON, W. J. (1983) Stable Galerkin methods for hyperbolic systems. *SIAM J. Numer. Anal.*, **20**, 221–233.
- MUGNOLO, D. (2006) Abstract wave equations with acoustic boundary conditions. *Math. Nachr.*, **279**, 299–318.
- NÉDÉLEC, J.-C. (1986) A new family of mixed finite elements in \mathbf{R}^3 . *Numer. Math.*, **50**, 57–81.
- PAZY, A. (1983) *Semigroups of linear operators and applications to partial differential equations*. Applied Mathematical Sciences, vol. 44. Springer-Verlag, New York, pp. viii+279.
- SHOWALTER, R. E. (1994) *Hilbert space methods for partial differential equations*. Electronic Monographs in Differential Equations, San Marcos, TX, pp. iii+242 pp. Electronic reprint of the 1977 original.
- SHOWALTER, R. E. (1997) *Monotone operators in Banach space and nonlinear partial differential equations*. Mathematical Surveys and Monographs, vol. 49. American Mathematical Society, Providence, RI, pp. xiv+278.

- TER ELST, A. F. M., SAUTER, M. & VOGT, H. (2015) A generalisation of the form method for accretive forms and operators. *J. Funct. Anal.*, **269**, 705–744.
- THOMÉE, V. (2006) *Galerkin finite element methods for parabolic problems*. Springer Series in Computational Mathematics, vol. 25, second edn. Springer-Verlag, Berlin, pp. xii+370.
- VEDURMUDI, A. P., GOULET, J., CHRISTENSEN-DALSGAARD, J., YOUNG, B. A., WILLIAMS, R. & VAN HEMMEN, J. L. (2016) How Internally Coupled Ears Generate Temporal and Amplitude Cues for Sound Localization. *Physical Review Letters*, **116**, 028101.
- ZEIDLER, E. (1990a) *Nonlinear functional analysis and its applications. II/A*. Springer-Verlag, New York, pp. xviii+467.
- ZEIDLER, E. (1990b) *Nonlinear functional analysis and its applications. II/B*. Springer-Verlag, New York, pp. i–xvi and 469–1202.
- ZHAO, J. (2004) Analysis of finite element approximation for time-dependent Maxwell problems. *Math. Comp.*, **73**, 1089–1105.