



Space-time discontinuous Petrov-Galerkin methods for linear wave equations in heterogeneous media

Johannes Ernesti, Christian Wieners

CRC Preprint 2018/13 (revised), February 2019

KARLSRUHE INSTITUTE OF TECHNOLOGY





Participating universities





Funded by



ISSN 2365-662X

SPACE-TIME DISCONTINUOUS PETROV-GALERKIN METHODS FOR LINEAR WAVE EQUATIONS IN HETEROGENEOUS MEDIA

JOHANNES ERNESTI AND CHRISTIAN WIENERS¹

Abstract. We establish an abstract space-time DPG framework for the approximation of linear waves in heterogeneous media. The estimates are based on a suitable variational setting in the energy space. The analysis combines the approaches for acoustic waves in Gopalakrishnan / Sepulveda (A space-time DPG method for acoustic waves, arXiv 2017) and in Ernesti / Wieners (RICCAM proceedings, submitted 2017) and is based on the abstract definition of traces on the skeleton of the time-space sub-structuring. The method is evaluated by large-scale parallel computations motivated from applications in seismic imaging, where the computational domain can be restricted substantially to a subset of the full space-time cylinder.

1991 Mathematics Subject Classification. 65N30.

February 12, 2019.

1. Introduction

Space-time finite elements aim for a unified analysis of discretization and solution methods in space and time. In particular, they allow for an efficient combined error control and for scaling of the solution scheme to the next generation of massively parallel computers.

The discontinuous Petrov-Galerkin method (DPG) is a well-suited finite element class for space-time applications which provides robust a-priori estimates, reliable error control, and the efficient hybridization to a symmetric positive definite Schur complement system. This is attractive for hyperbolic systems and allows to transfer features of discretizations for elliptic problems to wave-type equations. The long-term goal is, as it is discussed in [4] for the transport equation, to establish optimality of the solution process and of adaptive schemes. For a general discussion on the DPG technology we refer to [9].

First results of space-time DPG methods are established in [10] for the Schrödinger equations and in [14, 15, 17] for acoustic waves. Here, we show that the analysis transfers to general wave equations in heterogeneous media and provides robust estimated in the energy norm. Therefore, we recall in Lem. 3 and Lem. 4 the abstract DPG analysis based on the technique introduced in [10, 13, 17] which avoids explicit traces. Then, following the arguments in [4] we show that a test space exists which guarantees discrete inf-sup stability for general wave equations, and we extend the analysis for the simplified DPG method with nonconforming traces as in [15] to this more general setting. Finally, we apply a Strang-type argument to estimate the consistency error of the DPG method due to inexact quadrature in heterogeneous media.

Keywords and phrases: space-time methods, discontinuous Petrov-Galerkin finite elements, linear hyperbolic systems, heterogenous media

¹ Institut für Angewandte und Numerische Mathematik, KIT, Karlsruhe, Germany. Email: christian.wieners@kit.edu

The analysis is complemented by numerical results for wave propagation in heterogeneous media. Here we discuss an application scenario motivated from seismic measurements, were the wave signal is initiated by a point source and the results are only measured at selected points. In this application class, the finite propagation speed of wave solutions results into an a priori information about the region of interest within the space-time cylinder and which allows to truncate the computational domain substantially.

2. Linear hyperbolic systems

A semigroup framework. On a Lipschitz domain Ω_0 , we consider the evolution equation

$$M\partial_t u + Au = f$$
 in $(0,T) \times \Omega_0 \subset \mathbb{R} \times \mathbb{R}^d$

subject to homogeneous initial conditions u(0) = 0 in Ω_0 , where $f \in L_2((0,T) \times \Omega_0; \mathbb{R}^m)$ is a source function, and with

- a) a symmetric positive definite operator M in $L_2(\Omega_0; \mathbb{R}^m)$ represented by $M \in L_\infty(\Omega_0; \mathbb{R}^{m \times m}_{sym})$;
- b) a hyperbolic operator A with domain $\mathcal{D}(A) \subset L_2(\Omega_0; \mathbb{R}^m)$ such that

$$(Av, z)_{0,\Omega_0} = -(v, Az)_{0,\Omega_0}, \qquad v, z \in \mathcal{D}(A)$$

$$\tag{1}$$

and such that M + A is surjective, i.e.,

$$(M+A)(\mathcal{D}(A)) = \mathcal{L}_2(\Omega_0; \mathbb{R}^m).$$
⁽²⁾

In this setting, the Lumer-Phillips theorem [21, Thm. 12.22], or more general, the Hille-Yoshida theorem [8, Thm. XVII.3.2.1] applies: the operator $M^{-1}A$ generates a semigroup in $L_2(\Omega_0; \mathbb{R}^m)$ and for $f \in C^0((0,T); \mathcal{D}(A))$ the solution is given by

$$u(t) = \int_0^t \exp\left((t-s)M^{-1}A\right)M^{-1}f(s)\,\mathrm{d}s\,.$$
(3)

The solution belongs to the Banach space

$$\mathcal{V} = \left\{ v \in \mathrm{C}^1([0,T]; \mathrm{L}_2(\Omega_0; \mathbb{R}^m)) \cap \mathrm{C}^0([0,T]; \mathcal{D}(A)) : v(0) = 0 \right\}.$$

In particular, we obtain for the range of the space-time operator $L = M\partial_t + A$

$$C^{1}_{c}((0,T) \times \Omega_{0}; \mathbb{R}^{m}) \subset C^{0}((0,T); \mathcal{D}(A)) \subset L(\mathcal{V}) \subset L_{2}((0,T) \times \Omega_{0}; \mathbb{R}^{m})$$

$$\tag{4}$$

so that $L(\mathcal{V})$ is dense in $L_2((0,T) \times \Omega_0; \mathbb{R}^m)$, cf. [21, Thm. 12.16].

Linear wave equations. Our basic example is the acoustic wave equation for velocity and pressure with $(v, p) \in \mathcal{D}(A) = \mathrm{H}(\mathrm{div}, \Omega_0) \times \mathrm{H}^1_0(\Omega_0), \ M(v, p) = (\rho v, \kappa^{-1}p), \text{ and } A(v, p) = (\nabla p, \nabla \cdot v) \text{ where } \rho, \kappa \in \mathrm{L}_{\infty}(\Omega_0)$ model the spatially varying mass density and the bulk modulus of the material.

Note that in this example the definition of the domain $\mathcal{D}(A)$ includes homogeneous Dirichlet boundary conditions for the pressure p; more general boundary conditions can be included into the domain of the operator A, provided the conditions (1) and (2) are satisfied (see also [18, Sect. 2.2] for various examples).

This framework also applies to linear elastic waves described by $(v, \sigma) \in \mathcal{D}(A) = \mathrm{H}_{0}^{1}(\Omega_{0}; \mathbb{R}^{d}) \times \mathrm{H}(\mathrm{div}, \Omega_{0}; \mathbb{R}_{\mathrm{sym}}^{d \times d}),$ $M(v, \sigma) = (\rho v, \mathbb{C}^{-1}\sigma), \text{ and } A(v, \sigma) = (\mathrm{div}\,\sigma, \varepsilon(v)), \text{ and to electro-magnetic waves described by Maxwell's equations with <math>(E, H) \in \mathrm{H}_{0}(\mathrm{curl}, \Omega_{0}) \times \mathrm{H}(\mathrm{curl}, \Omega_{0}), M(E, H) = (\varepsilon E, \mu H), \text{ and } A(E, H) = (-\nabla \times H, \nabla \times E).$ Note that in all cases $\frac{1}{2} (Mu(t), u(t))_{\Omega_0}$ is the free energy, i.e., for elastic waves the kinetic and potential energy, and for the Maxwell case the electro-magnetic energy.

A main property of the linear wave equation is the finite speed of propagation $c_{\text{max}} > 0$, which allows – in case of local support of the source function f – to restrict the computation to the cone illustrated in Fig. 1

$$\mathcal{C}_{+}(\operatorname{supp} f) = \{(t, x) \in (0, T) \times \Omega_{0} \colon |x - x_{0}| \le c_{\max}(t - t_{0}) \text{ for all } (t_{0}, x_{0}) \in \operatorname{supp} f\},$$
(5)

i.e., for the solution $u \in \mathcal{V}$ of Lu = f holds $\operatorname{supp} u \in \mathcal{C}_+(\operatorname{supp} f)$, cf. [16, Chap. 7.2.4], [1]. The maximal wave speed can be determined by the equivalent formulation as a symmetric hyperbolic system, i.e., by using the representation of the linear operator of the form $Av = \sum_{j=1}^{d} A_j \partial_j v$ with symmetric matrices $A_j \in \mathbb{R}^{m \times m}_{\text{sym}}$. Then, the maximal speed of propagation in heterogeneous media is given by

$$c_{\max} = \|c\|_{\infty,\Omega_0}, \qquad c(x) = \max_{|n|=1} \max_{|w|=1} \frac{w^{\top} A_n w}{w^{\top} M(x) w}, \qquad A_n = \sum_{j=1}^d n_j A_j.$$
(6)

E.g., in the acoustic case we have $c(x) = \sqrt{\kappa(x)/\rho(x)}$.



FIGURE 1. The grey area depicts the cone of dependence $C_+(\{P_0\}) \subset (0,T) \times \Omega_0$ in 1D for a single point source at position P_0 . Due to the limited wave speed, information originating from P_0 only affects this space-time region, resulting in $C_+(\operatorname{supp} f) = \bigcup_{P_0 \in \operatorname{supp} f} C_+(\{P_0\})$ for a right-hand side f.

The adjoint equation. Let A^* be the adjoint operator of A with domain $\mathcal{D}(A^*)$, and let L^* be the adjoint operator for the wave equation backward in time. We assume that $M^{-1}A^*$ generates a semigroup such that for

$$\mathcal{V}^* = \left\{ z \in \mathcal{C}^1([0,T]; \mathcal{L}_2(\Omega_0; \mathbb{R}^m)) \cap \mathcal{C}^0([0,T]; \mathcal{D}(A^*)) : z(T) = 0 \right\},\$$

 $L^*(\mathcal{V}^*) \subset L_2(\Omega; \mathbb{R}^m)$ is dense by (4). E.g., this assumption is fulfilled for homogeneous Dirichlet or Neumann boundary conditions with $\mathcal{D}(A) = \mathcal{D}(A^*)$ and $A^* = -A$. We have $L^*v = -Lv$ for $v \in C^1_c((0,T) \times \Omega_0; \mathbb{R}^m)$, and

$$(Lv, z)_{0,(0,T)\times\Omega_0} = (v, L^*z)_{0,(0,T)\times\Omega_0}, \qquad v \in \mathcal{V}, \ z \in \mathcal{V}^*.$$
(7)

The corresponding backward cone for a domain of interest $\overline{\omega} \subset [0,T] \times \Omega_0$ is given by

$$\mathcal{C}_{-}(\overline{\omega}) = \left\{ (t,x) \in (0,T) \times \Omega_0 \colon |x - x_0| \le c_{\max}(t_0 - t) \text{ for all } (t_0, x_0) \in \overline{\omega} \right\}.$$
(8)

In the following, we consider applications where the source f has local support, and where the solution is evaluated only in the domain of interest $\overline{\omega}$, so that the solution process can be restricted to $C_+(\operatorname{supp} f) \cap C_-(\overline{\omega})$.

Subsets of the space-time cylinder. We consider $\Omega \subset (0,T) \times \Omega_0 \subset \mathbb{R} \times \mathbb{R}^d$ combining time slices of the form $\overline{\Omega} = \bigcup_{n=1}^{N} [t_{n-1}, t_n] \times \overline{\Omega}_n$ with open subsets $\Omega_n \subset \Omega_0$ for $n = 1, \ldots, N$ and a decomposition of the time interval $0 = t_0 < t_1 < \cdots < t_N = T$. In every time slice we select $\mathcal{D}(A; \Omega_n) \subset \mathcal{D}(A)$ such that the conditions (1) and (2) are satisfied in $L_2(\Omega_n; \mathbb{R}^m)$. This defines $\mathcal{V}_n = C^1([t_{n-1}, t_n]; L_2(\Omega_n; \mathbb{R}^m)) \cap C^0([t_{n-1}, t_n]; \mathcal{D}(A; \Omega_n))$. Using $v_n \in C^0([t_{n-1}, t_n]; L_2(\Omega_n; \mathbb{R}^m))$ for $v_n \in \mathcal{V}_n$ we define

$$\mathcal{V}(\Omega) = \left\{ v \in \mathcal{L}_2(\Omega; \mathbb{R}^m) \colon v_n \in \mathcal{V}_n \,, v(0) = 0 \,, v_n(t_n) = v_{n+1}(t_n) \text{ on } \Omega_n \cap \Omega_{n+1} \text{ and } v_{n+1}(t_n) = 0 \text{ on } \Omega_{n+1} \setminus \Omega_n \right\},$$
$$\mathcal{V}^*(\Omega) = \left\{ z \in \mathcal{L}_2(\Omega; \mathbb{R}^m) \colon z_n \in \mathcal{V}_n \,, z(T) = 0 \,, z_n(t_n) = z_{n+1}(t_n) \text{ on } \Omega_n \cap \Omega_{n+1} \text{ and } z_n(t_n) = 0 \text{ on } \Omega_n \setminus \Omega_{n+1} \right\}$$

with $v_n = v|_{(t_{n-1}, t_n)}$. By construction, we have

$$(Lv, z)_{0,\Omega} = (v, L^* z)_{0,\Omega}, \qquad v \in \mathcal{V}(\Omega), \ z \in \mathcal{V}^*(\Omega),$$
(9)

and the ranges $L(\mathcal{V}(\Omega))$, $L^*(\mathcal{V}^*(\Omega))$ are dense in the energy space $W = L_2(\Omega; \mathbb{R}^m)$. The corresponding energy norm in space and time is given by $||w||_W = \sqrt{(Mw, w)_{0,\Omega}}$ for $w \in W$.

Lemma 1. We have for $u \in \mathcal{V}(\Omega)$ and f = Lu

$$||u||_W \le \frac{T}{\sqrt{2}} ||M^{-1}f||_W.$$

This is a Poincaré type estimate since it relies on the initial condition u(0) = 0.

Proof. The estimate relies on the representation (3) of the solution in every slice $[t_{n-1}, t_n] \times \Omega_n$. In every slice, define $W_n = L_2(\Omega_n; \mathbb{R}^m)$ and we use the energy inner product $(v_n, w_n)_{W_n} = (Mv_n, w_n)_{0,\Omega_n}$ for $v_n, w_n \in W_n$. Since the operator $M^{-1}A$ is skew-adjoint in W_n , i.e., $(M^{-1}Av_n, v_n)_{W_n} = (Av_n, v_n)_{0,\Omega_n} = 0$ for $v_n \in \mathcal{D}(A; \Omega_n)$, the spectrum is contained in i \mathbb{R} which yields $\|\exp(sM^{-1}A)v_n\|_{W_n} \le \|v_n\|_{W_n}$ for all $s \in \mathbb{R}$. Now, inserting

$$u(t) = \exp\left((t - t_{n-1})M^{-1}A\right)u(t_{n-1}) + \int_{t_{n-1}}^{t} \exp\left((t - s)M^{-1}A\right)M^{-1}f(s)\,\mathrm{d}s\,, \qquad t \in (t_{n-1}, t_n]\,.$$

yields recursively for $t \in (t_{n-1}, t_n]$

$$\begin{aligned} \|u(t)\|_{W_{n}} &\leq \|u(t_{n-1})\|_{W_{n}} + \int_{t_{n-1}}^{t} \|M^{-1}f(s)\|_{W_{n}} \,\mathrm{d}s \\ &\leq \|u(t_{n-1})\|_{W_{n}} + \sqrt{t - t_{n-1}} \left(\int_{t_{n-1}}^{t} \|M^{-1}f(s)\|_{W_{k}}^{2} \,\mathrm{d}s \right)^{1/2} \\ &\leq \sum_{k=1}^{n-1} \sqrt{t_{k} - t_{k-1}} \left(\int_{t_{k-1}}^{t_{k}} \|M^{-1}f(s)\|_{W_{k}}^{2} \,\mathrm{d}s \right)^{1/2} + \sqrt{t - t_{n-1}} \left(\int_{t_{n-1}}^{t} \|M^{-1}f(s)\|_{W_{n}}^{2} \,\mathrm{d}s \right)^{1/2} \\ &\leq \sqrt{t} \left(\sum_{k=1}^{n-1} \int_{t_{k-1}}^{t_{k}} \|M^{-1}f(s)\|_{W_{k}}^{2} \,\mathrm{d}s + \int_{t_{n-1}}^{t} \|M^{-1}f(s)\|_{W_{n}}^{2} \,\mathrm{d}s \right)^{1/2}. \end{aligned}$$

Together, this yields

$$\|u\|_{W}^{2} = \sum_{n=1}^{N} \int_{t_{n-1}}^{t_{n}} \|u(t)\|_{W_{n}}^{2} dt \leq \int_{0}^{T} t \left(\sum_{k=1}^{N} \int_{t_{k-1}}^{t_{k}} \|M^{-1}f(s)\|_{W_{k}}^{2} ds \right) dt = \frac{1}{2} T^{2} \|M^{-1}f\|_{W}^{2}.$$

A variational setting. We extend the operator L in $\mathcal{V}(\Omega) \subset W$ to a suitable Hilbert space defined by

$$\begin{aligned} \mathrm{H}(L,\Omega) &= \{ v \in W \colon Lv \in W \} \\ &= \{ v \in W \colon w \in W \text{ exists such that } (w,z)_{0,\Omega} = (v,L^*z)_{0,\Omega} \text{ for all } z \in \mathrm{C}^1_{\mathrm{c}}(\Omega;\mathbb{R}^m) \} . \end{aligned}$$

We use the weighted graph norm $\|v\|_{\mathrm{H}(L,\Omega)} = \sqrt{\|v\|_W^2 + \|M^{-1}Lv\|_W^2} = \sqrt{(Mv,v)_{0,\Omega} + (M^{-1}Lv,Lv)_{0,\Omega}}$. The closure of $\mathrm{C}^1_{\mathrm{c}}(\Omega,\mathbb{R}^m)$ with respect to $\|\cdot\|_{\mathrm{H}(L,\Omega)}$ is denoted by $\mathrm{H}_0(L,\Omega)$, cf. [11, Sect. 5.5.2].

In the hyperbolic case we have $H(L, \Omega) = H(L^*, \Omega)$. Nevertheless, since L is associated to the forward problem and L^* to the backward problem, we will need different subspaces in the following arguments.

Since $C_c^1(\Omega, \mathbb{R}^m)$ is dense in $L_2(\Omega, \mathbb{R}^m)$, this defines $L \in \mathcal{L}(H(L, \Omega), W)$ and $L^* \in \mathcal{L}(H(L^*, \Omega), W)$ which are consistent extensions of the operators L in $\mathcal{V}(\Omega)$ and L^* in $\mathcal{V}^*(\Omega)$.

Let V be the closure of $\mathcal{V}(\Omega)$ in $\mathcal{H}(L,\Omega)$, and let V^* be the closure of $\mathcal{V}^*(\Omega)$ in $\mathcal{H}(L^*,\Omega)$. Then, Lem. 1 yields

$$\|v\|_{W} \leq C_{L} \|M^{-1}Lv\|_{W}, \quad v \in V$$
(10)

with $C_L = T/\sqrt{2}$, and, correspondingly, $||z||_W \leq C_L ||M^{-1}L^*z||_W$ for $z \in V^*$ for the adjoint problem backwards in time. Moreover, (10) implies that $L(V) \subset W$ is closed, and since $L(\mathcal{V}) \subset W$ is dense, we obtain L(V) = W, so that $L \in \mathcal{L}(V, W)$ is a bijection. Furthermore, (9) extends to

$$(Lv, z)_{0,\Omega} = (v, L^* z)_{0,\Omega}, \qquad v \in V, \ z \in V^*.$$
 (11)

The solution in the restricted space-time domain $\Omega \subset (0,T) \times \Omega_0$ is now compared with the solution in the full space-time cylinder $(0,T) \times \Omega_0$. Therefore, let V_0 and V_0^* be the closures of $\mathcal{V} = \mathcal{V}((0,T) \times \Omega_0)$ and $\mathcal{V}^* = \mathcal{V}^*((0,T) \times \Omega_0)$. Furthermore, set $W_0 = L_2((0,T) \times \Omega_0; \mathbb{R}^m)$.

Lemma 2. Let $f_0 \in W_0$ be a source function and let $u_0 \in V_0$ be the unique solution of $Lu_0 = f_0$ in $(0, T) \times \Omega_0$. Assume that $\Omega \supset C_+(\text{supp } f_0) \cap C_-(\overline{\omega})$ for a domain of interest $\overline{\omega} \subset [0, T] \times \overline{\Omega}_0$. Then, we have for the unique solution $u \in V$ of Lu = f with restricted source function $f = f_0|_{\Omega} \in W$ consistency with u_0 in ω , i.e.,

$$u|_{\omega} = u_0|_{\omega} \,.$$

Proof. Let $z_0 \in V_0^*$ be the dual solution with respect to the goal functional induced by $(u-u_0)|_{\omega}$, i.e.,

$$(L^*z_0, w_0)_{0,(0,T)\times\Omega} = (u - u_0, w_0)_{0,\omega}, \qquad w_0 \in W_0.$$

Then, we observe supp $z_0 \subset \mathcal{C}_{-}(\overline{\omega})$.

Let $\chi \in C_c^1((0,T) \times \Omega_0)$ be a function with $\operatorname{supp} \chi \subset \Omega$ and $\chi \equiv 1$ in $\mathcal{C} = \mathcal{C}_+(\operatorname{supp} f_0) \cap \mathcal{C}_-(\overline{\omega})$. This extends $u \in V$ to $\chi u \in V_0$ with u(t,x) = 0 for $(t,x) \in (0,T) \times \Omega_0 \setminus \Omega$. Using $\operatorname{supp} u \in \mathcal{C}_+(\operatorname{supp} f_0)$ and (11) we obtain

$$(L^*z_0, \chi u)_{0,\Omega} = (L^*z_0, \chi u)_{0,(0,T) \times \Omega_0} = (z_0, L\chi u)_{0,(0,T) \times \Omega_0} = (z_0, Lu)_{0,\mathcal{C}} = (z_0, Lu)_{0,\Omega}$$

and $(L^*z_0, \chi u_0)_{0,(0,T)\times\Omega_0} = (z_0, Lu_0)_{0,(0,T)\times\Omega_0}$. This yields the assertion by

$$\begin{aligned} \|u - u_0\|_{0,\omega}^2 &= (u - u_0, \chi(u - u_0))_{0,\omega} \\ &= (L^* z_0, \chi u)_{0,\Omega} - (L^* z_0, \chi u_0)_{0,(0,T) \times \Omega_0} \\ &= (z_0, L u)_{0,\Omega} - (z_0, L u_0)_{0,(0,T) \times \Omega_0} \\ &= (z_0, f)_{0,\Omega} - (z_0, f_0)_{0,(0,T) \times \Omega_0} = (z_0, f)_{0,\Omega} - (z_0, f_0)_{0,\mathcal{C}} = 0. \end{aligned}$$

Integration by parts defines the operators $D_{\Omega} \in \mathcal{L}(\mathrm{H}(L,\Omega),\mathrm{H}(L^*,\Omega)')$ and $D'_{\Omega} \in \mathcal{L}(\mathrm{H}(L^*,\Omega),\mathrm{H}(L,\Omega)')$ with

$$\langle D_{\Omega}v, z \rangle = (Lv, z)_{0,\Omega} - (v, L^*z)_{0,\Omega} = \langle D'_{\Omega}z, v \rangle, \qquad v \in \mathcal{H}(L, \Omega), \ z \in \mathcal{H}(L^*, \Omega)$$

The kernel of D_{Ω} is denoted by $\mathcal{N}(D_{\Omega}) = \{y \in \mathrm{H}(L,\Omega) : \langle D_{\Omega}y, z \rangle = 0 \text{ for all } z \in \mathrm{H}(L^*,\Omega)\} \supset \mathrm{H}_0(L,\Omega)$. In fact, the boundary conditions in $\mathrm{H}_0(L,\Omega)$ and in V are characterized by duality. Following [13, Lem. 2.4] and [6], this is shown using properties of polar sets, cf. [22, Sect. 4.5], applied to

$$C^{1}_{c}(\Omega; \mathbb{R}^{m})^{\perp} = \left\{ \eta \in H(L, \Omega)' : \langle \eta, v \rangle = 0 \text{ for all } v \in C^{1}_{c}(\Omega; \mathbb{R}^{m}) \right\},$$

$${}^{\perp}D'_{\Omega}(\mathcal{V}^{*}(\Omega)) = \left\{ v \in H(L, \Omega) : \langle D_{\Omega}v, z \rangle = 0 \text{ for all } z \in \mathcal{V}^{*}(\Omega) \right\}.$$

Lemma 3. We have $H_0(L, \Omega) = \mathcal{N}(D_\Omega)$ and $V = {}^{\perp}D'_{\Omega}(\mathcal{V}^*(\Omega))$.

Proof. For a given functional $\ell \in C^1_c(\Omega; \mathbb{R}^m)^{\perp}$ we define $u_\ell \in H(L, \Omega)$ solving

$$\left(M^{-1}Lu_{\ell}, Lv\right)_{0,\Omega} + \left(Mu_{\ell}, v\right)_{0,\Omega} = \langle \ell, v \rangle, \quad v \in \mathcal{H}(L,\Omega).$$

$$(12)$$

Then, $(Mu_{\ell}, v)_{0,\Omega} = -(M^{-1}Lu_{\ell}, Lv)_{0,\Omega}$ for all $v \in C_c^1(\Omega; \mathbb{R}^m)$, which yields $z_{\ell} = M^{-1}Lu_{\ell} \in H(L^*, \Omega)$ and $L^*z_{\ell} = -Mu_{\ell}$. This is inserted in (12), and we obtain for the adjoint operator D'_{Ω}

$$\left\langle D'_{\Omega} z_{\ell}, v \right\rangle = \left\langle D_{\Omega} v, z_{\ell} \right\rangle = \left(L v, z_{\ell} \right)_{0,\Omega} - \left(v, L^* z_{\ell} \right)_{0,\Omega} = \left(L v, M^{-1} L u_{\ell} \right)_{0,\Omega} + \left(v, M u_{\ell} \right)_{0,\Omega} = \left\langle \ell, v \right\rangle, \quad v \in \mathcal{H}(L,\Omega),$$

i.e., $D'_{\Omega} z_{\ell} = \ell$. This proves $C^1_c(\Omega; \mathbb{R}^m)^{\perp} \subset D'_{\Omega}(H(L^*, \Omega))$, and by duality we conclude the first assertion

$$\mathcal{N}(D_{\Omega}) = {}^{\perp}D'_{\Omega}(\mathrm{H}(L^*, \Omega)) \subset {}^{\perp}(\mathrm{C}^{1}_{\mathrm{c}}(\Omega; \mathbb{R}^m)^{\perp}) = \mathrm{H}_{0}(L, \Omega).$$

Since ${}^{\perp}D'_{\Omega}(\mathcal{V}^*(\Omega))$ is closed in $\mathrm{H}(L,\Omega)$ and $\mathcal{V}(\Omega) \subset {}^{\perp}D'_{\Omega}(\mathcal{V}^*(\Omega))$ by (9), we have $V \subset {}^{\perp}D'_{\Omega}(\mathcal{V}^*(\Omega))$. On the other hand, for $v \in {}^{\perp}D'_{\Omega}(\mathcal{V}^*(\Omega)) \subset \mathrm{H}(L,\Omega)$ set f = Lv and let $u \in V$ be the unique solution of Lu = f. Together, we have by construction $u - v \in {}^{\perp}D'_{\Omega}(\mathcal{V}^*(\Omega))$ and L(u - v) = 0, so that

$$0 = \left\langle D'_{\Omega} z, u - v \right\rangle = \left(L(u - v), z \right)_{0,\Omega} - \left(u - v, L^* z \right)_{0,\Omega} = \left(v - u, L^* z \right)_{0,\Omega}, \qquad z \in \mathcal{V}^*(\Omega).$$

Since $L^*(\mathcal{V}^*(\Omega)) \subset W$ is dense, we obtain $u = v \in V$.

3. Space-time substructuring

Let $\Omega_h = \bigcup K \subset \Omega$ be a decomposition into open convex subsets K with skeleton $\partial \Omega_h$ satisfying $\overline{\Omega} = \Omega_h \cup \partial \Omega_h$. We consider the broken space $Z = \mathrm{H}(L^*, \Omega_h) = \prod_K \mathrm{H}(L^*, K)$ and its dual space $Z' = \mathrm{H}(L^*, \Omega_h)'$, again using the weighted norm $||z||_Z = \sqrt{||z||_W + ||M^{-1}L^*z||_W}$ with $||z||_W = \sqrt{(Mz, z)_{0,\Omega_h}}$ evaluated in Ω_h .

Integration by parts defines the operators
$$D_K \in \mathcal{L}(\mathrm{H}(L,K),\mathrm{H}(L^*,K))$$
 and $D_{\Omega_h} \in \mathcal{L}(\mathrm{H}(L,\Omega_h),Z')$ by
 $\langle D_{\Omega_h}v,z \rangle = (Lv,z)_{0,\Omega_h} - (v,L^*z)_{0,\Omega_h} = \sum_K (Lv,z)_{0,K} - (v,L^*z)_{0,K} = \sum_K \langle D_Kv|_K,z|_K \rangle, \ v \in \mathrm{H}(L,\Omega_h), \ z \in Z.$

Lem. 3 yields $H_0(L, K) = \mathcal{N}(D_K)$ and thus $H_0(L, \Omega_h) = \prod_K H_0(L, K) = \mathcal{N}(D_{\Omega_h})$. This allows to identify traces on the skeleton $\partial \Omega_h$ with functionals $D_{\Omega_h}(H(L, \Omega_h)) \subset Z'$. By construction, we have

$$|\langle D_{\Omega_h} v, z \rangle| = \left| (M^{-1}Lv, Mz)_{0,\Omega_h} - (Mv, M^{-1}L^*z)_{0,\Omega_h} \right| \le ||v||_{\mathcal{H}(L,\Omega_h)} ||z||_Z, \qquad v \in \mathcal{H}(L,\Omega_h), \ z \in \mathbb{Z},$$

i.e., $||D_{\Omega_h}v||_{Z'} \le ||v||_{\mathcal{H}(L,\Omega_h)}$.

 $\mathbf{6}$

For given $f \in W$ let $u \in V$ be the unique solution of Lu = f. Then, we observe

$$(u, L^*z)_{0,\Omega_h} + \langle D_{\Omega_h}u, z \rangle = (Lu, z)_{0,\Omega_h} = (f, z)_{0,\Omega_h}, \qquad z \in \mathbb{Z}.$$

This allows for a weak characterization in the energy space W and the abstract trace space $\hat{V} = D_{\Omega_h}(V) \subset Z'$. Therefore, we define the bilinear form in $(W \times Z') \times Z$ by

$$b(w, \hat{v}; z) = (v, L^* z)_{0,\Omega_h} + \langle \hat{v}, z \rangle, \qquad (w, \hat{v}) \in W \times Z', \ z \in Z.$$

We now show inf-sup stability of the bilinear form $b(\cdot, \cdot)$, which implies that a unique weak solution $(u, \hat{u}) \in W \times \hat{V}$ exists with $b(u, \hat{u}; z) = (f, z)_{0,\Omega_h}$ for all $z \in Z$.

In order to obtain improved optimal estimates, we use in $W \times Z'$ the norm $\|(w, \hat{v})\|_{W \times Z'} = \max\{\|w\|_W, \|\hat{v}\|_{Z'}\}$. Then, the bilinear form $b(\cdot; \cdot)$ is continuous with $|b(w, \hat{v}; z)| \leq \|(w, \hat{v})\|_{W \times Z'} \|z\|_Z$, and we obtain the following estimate for the inf-sup stability for the ideal DPG method, cf. [6].

Lemma 4. The bilinear form $b(\cdot; \cdot)$ is injective on $W \times \hat{V}$ and inf-sup stable with $\beta = (2C_L^2 + 2)^{-1/2}$, i.e.,

$$\sup_{z \in Z} \frac{b(w, \hat{v}; z)}{\|z\|_{Z}} \ge \beta \,\|(w, \hat{v})\|_{W \times Z'} \,, \qquad (w, \hat{v}) \in W \times \hat{V} \,. \tag{13}$$

Proof. Let $(w, \hat{v}) \in W \times \hat{V}$ be in the kernel of b, i.e, $b(w, \hat{v}; z) = 0$ for $z \in Z$. Then we have

$$0 = (w, L^*z)_{0,\Omega_h} + \langle \hat{v}, z \rangle = (w, L^*z)_{0,\Omega_h}, \qquad z \in \mathcal{C}^1_{\mathsf{c}}(\Omega_h, \mathbb{R}^m),$$

i.e., $w \in H(L, \Omega_h)$ and Lw = 0 in Ω_h . This yields for all $v \in V$ with $D_{\Omega_h}v = \hat{v}$

$$\left\langle D_{\Omega_h}(w-v), z \right\rangle = \left(Lw, z \right)_{0,\Omega_h} - \left(w, L^* z \right)_{0,\Omega_h} - \left\langle \hat{v}, z \right\rangle = 0 - b(w, \hat{v}; z) = 0, \qquad z \in \mathcal{H}(L^*, \Omega_h),$$

i.e., $w - v \in \mathcal{N}(D_{\Omega_h}) = \mathrm{H}_0(L, \Omega_h)$. Thus we have $w \in v + \mathrm{H}_0(L, \Omega_h) \subset V$, and together with Lw = 0 and using (10) we obtain w = 0. Thus, b is injective on $W \times \hat{V}$.

In the second step we show inf-sup stability. For given $z \in Z \subset W$ we select $u_z \in V$ with $Lu_z = Mz$, which yields $b(u_z, D_{\Omega_h} u_z; z) = (u_z, L^*z)_{0,\Omega_h} + (Lu_z, z)_{0,\Omega_h} - (u_z, L^*z)_{0,\Omega_h} = (Lu_z, z)_{0,\Omega_h} = ||z||_W^2$.

Inserting (10) we obtain

$$\begin{aligned} \|(u_z, D_{\Omega_h} u_z)\|_{W \times Z'} &= \max \left\{ \|u_z\|_W, \|D_{\Omega_h} u_z\|_{Z'} \right\} \le \max \left\{ \|u_z\|_W, \|u_z\|_{\mathcal{H}(L,\Omega_h)} \right\} \\ &\le \|u_z\|_{\mathcal{H}(L,\Omega_h)} \le \sqrt{C_L^2 + 1} \|M^{-1} L u_z\|_W = \sqrt{C_L^2 + 1} \|z\|_W. \end{aligned}$$

This establishes for all $z \in Z$ by selecting $(u_z, D_{\Omega_h} u_z)$ and $(M^{-1}L^*z, 0)$ in $W \times Z'$

$$\sup_{(v,\hat{v})\in W\times\hat{V}} \frac{b(v,\hat{v};z)}{\|(v,\hat{v})\|_{W\times Z'}} \ge \max\left\{\frac{b(u_z, D_{\Omega_h}u_z;z)}{\|(u_z, D_{\Omega_h}u_z)\|_{W\times Z'}}, \frac{b(M^{-1}L^*z, 0;z)}{\|(M^{-1}L^*z, 0)\|_{W\times Z'}}\right\}$$
(14)
$$\ge \max\left\{\frac{\|z\|_W^2}{\sqrt{C_L^2 + 1}\|z\|_W}, \frac{\|M^{-1}L^*z\|_W^2}{\|M^{-1}L^*z\|_W}\right\}$$
$$\ge \frac{\max\left\{\|z\|_W, \|M^{-1}L^*z\|_W\right\}}{\sqrt{C_L^2 + 1}} \ge \frac{\|z\|_Z}{\sqrt{2\sqrt{C_L^2 + 1}}}.$$

Since $b(\cdot; \cdot)$ is injective in $W \times \hat{V}$, we obtain (13) by duality [2, Lem. 4.4.2].

4. Petrov-Galerkin estimates

In Lem. 4 we established a variational space-time setting for weak solutions in the product space $Y = W \times \hat{V}$ with a broken test space Z. Introducing the trial-to-test operator T_Z , see Tab. 1, we observe

$$||T_Z y||_Z = \sup_{z \in Z} \frac{b(y; z)}{||z||_Z} \ge \beta ||y||_Y, \quad y \in Y,$$

i.e., inf-sup stability of the bilinear form $b(\cdot; \cdot)$ in $Y \times Z$ implies ellipticity $||T_Z y||_Z^2 = \langle Sy, y \rangle \ge \beta^2 ||y||_Y^2$ of the corresponding symmetric Schur complement problem in Y.

Now we select discrete spaces $W_h \subset W$ and $V_h \subset V$, and we set $\hat{V}_h = D_{\Omega_h}(V_h) \subset Z'$ and $Y_h = W_h \times \hat{V}_h \subset Y$. An appropriate discrete test space $Z_h \subset Z$ always exists (see, e.g., [4, Thm. 4.8] for the transport equation and [26, Sect. 6]).

TABLE 1. Operators for the Petrov-Galerkin analysis.

	$B \in \mathcal{L}(Y, Z')$	$\langle By,z\rangle=b(y;z)$	$y\in Y,\ z\in Z$
Riesz operator	$A_Z \in \mathcal{L}(Z, Z')$	$\langle A_Z z, \psi \rangle = (z, \psi)_Z$	$z,\psi\in Z$
trial-to-test operator	$T_Z = A_Z^{-1}B \in \mathcal{L}(Y, Z)$	$(T_Z y, z)_Z = b(y; z)$	$y\in Y,\ z\in Z$
Schur complement	$S = B' A_Z^{-1} B \in \mathcal{L}(Y, Y')$		
natural embedding	$E_{Y_h} \in \mathcal{L}(Y_h, Y)$	$E_{Y_h}y_h = y_h$	$y_h \in Y_h$
	$E_{Z_h} \in \mathcal{L}(Z_h, Z)$	$E_{Z_h} z_h = z_h$	$z_h \in Z_h$
	$A_{Z_h} = E'_{Z_h} A_Z E_{Z_h} \in \mathcal{L}(Z_h, Z'_h)$	$\langle A_{Z_h} z_h, \psi_h \rangle = (z_h, \psi_h)_Z$	$z_h, \psi_h \in Z_h$
Galerkin projection	$P_{Z_h} = A_{Z_h}^{-1} E'_{Z_h} A_Z \in \mathcal{L}(Z, Z_h)$	$(P_{Z_h}z,\psi_h)_Z = (z,\psi_h)_Z$	$z \in Z, \ \psi_h \in Z_h$
	$B_h = E'_{Z_h} B E'_{Y_h} \in \mathcal{L}(Y_h, Z'_h)$	$\langle B_h y_h, z_h \rangle = b(y_h; z_h)$	$y_h \in Y_h, z_h \in Z_h$
	$T_{Z_h} = A_{Z_h}^{-1} B_h = P_{Z_h} T_Z E_{Y_h} \in \mathcal{L}(Y_h, Z_h)$	$(T_{Z_h}y_h, z_h)_Z = b(y_h; z_h)$	$y_h \in Y_h, z_h \in Z_h$
	$S_h = B'_h A_{Z_h}^{-1} B_h \in \mathcal{L}(Y_h, Y'_h)$		

Lemma 5. For given $\beta_h \in (0, \beta)$ a discrete test space $Z_h \subset Z$ exists such that

$$\|T_{Z_h}y_h\|_Z = \sup_{z_h \in Z_h} \frac{b(y_h; z_h)}{\|z_h\|_Z} \ge \beta_h \|y_h\|_Y, \qquad y_h \in Y_h.$$
(15)

Again this implies ellipticity $\langle S_h y_h, y_h \rangle \ge \beta_h^2 \|y_h\|_Y^2$ of the corresponding discrete Schur complement problem.

Proof. Let $Z_h^{\text{opt}} = T_Z E_{Y_h}(Y_h)$ be the optimal test space, and let $Z_{h,k}, k \in \mathbb{N}$ be a dense family of discrete spaces so that $z = \lim_{k \to \infty} P_{Z_{h,k}} z$ for all $z \in Z$. Since Z_h^{opt} is discrete, for all $\varepsilon \in (0,1)$ some $k = k(\varepsilon) \in \mathbb{N}$ exists such that $\|P_{Z_{h,k}} z_h - z_h\|_Z \le \varepsilon \|z_h\|_Z$ for all $z_h \in Z_h^{\text{opt}}$. This yields

$$\|P_{Z_{h,k}}z_h\|_Z^2 = \|z_h\|_Z^2 - \|P_{Z_{h,k}}z_h - z_h\|_Z^2 \ge \|z_h\|_Z^2 - \varepsilon^2 \|z_h\|_Z^2 = (1 - \varepsilon^2) \|z_h\|_Z^2, \qquad z_h \in Z_h^{\text{opt}}.$$

Now, for given $\beta_h \in (0, \beta)$, select $\varepsilon = \sqrt{1 - \beta_h^2/\beta^2} > 0$ and set $Z_h = Z_{h,k(\varepsilon)}$. This yields for all $y_h \in Y_h$

$$\sup_{z_h \in Z_h} \frac{b(y_h; z_h)}{\|z_h\|_Z} = \|T_{Z_h} y_h\|_Z = \|P_{Z_h} T_Z E_{Y_h} y_h\|_Z \ge \sqrt{1 - \varepsilon^2} \|T_Z E_{Y_h} y_h\|_Z \ge \sqrt{1 - \varepsilon^2} \beta \|y_h\|_Y = \beta_h \|y_h\|_Y. \quad \Box$$

TABLE 2. Operators and bilinear forms for the local Petrov-Galerkin analysis with restricted spaces $Y_K = L_2(K; \mathbb{R}^m) \times H(K, L)$ and $Z_K = H(K, L^*)$ for $K \subset \Omega_h$, and local finite element spaces $W_{K,h} = W_h|_K$, $V_{K,h} = V_h|_K$, $Y_{K,h} = W_{K,h} \times V_{K,h}$, and $Z_{K,h} = Z_h|_K$.

$$\begin{array}{ll} B_{K} \in \mathcal{L}(Y_{K}, Z'_{K}) & \langle B_{K}y_{K}, z_{K} \rangle = b_{K}(y_{K}; z_{K}) & y_{K} \in Y_{K}, \ z_{K} \in Z_{K} \\ B_{K,h} \in \mathcal{L}(Y_{K,h}, Z'_{K,h}) & \langle B_{K,h}y_{K,h}, z_{K,h} \rangle = b_{K}(y_{K}; z_{K}) & y_{K,h} \in Y_{K,h}, \ z_{K,h} \in Z_{K,h} \\ \hline A_{Z_{K,h}} \in \mathcal{L}(Z_{K,h}, Z'_{K,h}) & \langle A_{Z_{K,h}}z_{K,h}, \psi_{K,h} \rangle = (z_{K,h}, \psi_{K,h})z_{K} & z_{K,h}, \psi_{K,h} \in Z_{K,h} \\ \hline A_{Y_{K,h}} \in \mathcal{L}(Y_{K,h}, Y'_{K,h}) & \langle A_{Y_{K,h}}y_{K,h}, \phi_{K,h} \rangle = (y_{K,h}, \phi_{K,h})y_{K} & y_{K,h}, \phi_{K,h} \in Y_{K,h} \\ \hline E_{Y_{K,h}} \in \mathcal{L}(Y_{K,h}, Y_{K}) & E_{Y_{K,h}}y_{K,h} = y_{K,h} & y_{K,h} \in Y_{K,h} \\ \hline S_{K,h} = B'_{K,h}A^{-1}_{Z_{K,h}}B_{K,h} \in \mathcal{L}(Y_{K,h}, Y'_{K,h}) \end{array}$$

Since the optimal test space Z_h^{opt} is not accessible, the proof in Lem. 5 is not constructive, and the norm in $Y_h = W_h \times \hat{V}_h \subset W \times Z'$ cannot be evaluated. Nevertheless, for broken test spaces $Z_h = \prod Z_{K,h}$ the well-posedness of the discrete problem can be tested by a local criterion, and norm estimates can be evaluated in $W_{K,h} \times V_{K,h} = (W_h \times V_h)|_K$ using the norm in $V_{K,h} \subset H(L,K)$, so that for computable estimates it is not required to evaluate the norm in the trace space $\hat{V}_{K,h} = D_K(V_{K,h}) \subset Z'_{K,h}$.

Therefore we introduce local operators (see Tab. 2), the local bilinear form $b_K(\cdot; \cdot)$ defined by

$$b_K(w_K, v_K; z_K) = (w_K, L^* z_K)_{0,K} + \langle D_K v_K, z_K \rangle, \qquad (w_K, v_K) \in Y_K, \ z_K \in Z_K,$$

and for all $z_K \in Z_K$ the local affine spaces

$$Z_{K,h}(z_K) = \left\{ z_{K,h} \in Z_{K,h} : b_K(y_{K,h}, z_K - z_{K,h}) = 0 \text{ for all } y_{K,h} \in Y_{K,h} \right\}.$$

Lemma 6. If $Z_{K,h}(z_K)$ is not empty for all $z_K \in Z_K$ and $K \subset \Omega_h$, the operator B_h is injective in Y_h and a Fortin operator exists, i.e., a projection $\Pi_h \in \mathcal{L}(Z, Z_h)$ with $b(y_h; z - \Pi_h z) = 0$ for all $y_h \in Y_h$ and $z \in Z$.

This provides discrete stability and the estimate $\beta_h \geq \beta / ||\Pi_h||_Z$ for the inf-sup constant [3, Prop. II.2.8].

Proof. For $z_K \in Z_K$ and $K \subset \Omega_h$ we define $\Pi_{K,h} z_K \in Z_{K,h}(z_K)$ as the element with minimal norm. An explicit representation is derived as follows. We compute a critical point $(z_{K,h}, y_{K,h}) \in Z_{K,h} \times Y_{K,h}$ of the corresponding local Lagrange functional

$$\begin{aligned} F_{K,h}(z_{K,h}, y_{K,h}) &= \frac{1}{2} \| z_{K,h} \|_{Z_K}^2 + b_K(y_{K,h}; z_K - z_{K,h}) \\ &= \frac{1}{2} \langle A_{Z_{K,h}} z_{K,h}, z_{K,h} \rangle + \langle B_K E_{Y_{K,h}} y_{K,h}, z_K \rangle - \langle B_{K,h} y_{K,h}, z_{K,h} \rangle \,, \end{aligned}$$

i.e., $A_{Z_{K,h}} z_{K,h} = B_{K,h} y_{K,h}$ and $B'_{K,h} z_{K,h} = E'_{Y_{K,h}} B'_K z_K$. This yields

$$S_{K,h}y_{K,h} = B'_{K,h}A_{Z_{K,h}}^{-1}B_{K,h}y_{K,h} = E'_{Y_{K,h}}B'_{K}z_{K}.$$
(16)

We have

$$\langle E'_{Y_{K,h}}B'_K z_K, \psi_{K,h} \rangle = b_K(\psi_{K,h}; z_K) = \langle B_{K,h}\psi_{K,h}, E_{Z_{K,h}}z_K \rangle = 0, \qquad \psi_{K,h} \in \mathcal{N}(B_{K,h}),$$

i.e., $E'_{Y_{K,h}}B'_K z_K \in \mathcal{N}(B_{K,h})^{\perp}$. Since $S_{K,h}$ is self-adjoint and $\mathcal{N}(S_{K,h}) = \mathcal{N}(B_{K,h})$, this shows that a Lagrange multiplier $y_{K,h} \in Y_{K,h}$ solving the local Schur complement problem (16) exists, but in general, the solution is not unique. Thus we select the solution with minimal norm in $Y_{K,h}$. This is determined by the pseuso-inverse with respect to the topology in Y_K

$$S_{K,h}^{+} = \lim_{\delta \to 0} \left(S_{K,h} A_{Y_{K,h}}^{-1} S_{K,h} + \delta A_{Y_{K,h}} \right)^{-1} S_{K,h} A_{Y_{K,h}}^{-1} \in \mathcal{L}(Y_{K,h}', Y_{K,h}).$$

Then, $y_{K,h} = S_{K,h}^+ E'_{Y_{K,h}} B'_K z_K$ is a Lagrange multiplier and the minimizer is given by $z_{K,h} = \prod_{K,h} z_K$ with

$$\Pi_{K,h} = A_{Z_{K,h}}^{-1} B_{K,h} S_{K,h}^{+} E_{Y_{K,h}}' B_{K}' \in \mathcal{L}(Z, Z_{K,h})$$

This defines $\Pi_h z = \left(\sum_{K,h} E_{Z_{K,h}} \Pi_{K,h} z|_K\right)_K$, and thus $b(y_h, z - \Pi_h z) = \sum_K b_K(y_h|_K, z|_K - \Pi_{K,h} z|_K) = 0$ for all $y_h \in Y_h$, i.e., Π_h is a Fortin operator. Then, for all $y_h \in \mathcal{N}(B_h)$

$$0 = b(y_h; \Pi_h z) = b(y_h; z) = \langle B E_{Y_h} y_h, z \rangle, \qquad z \in \mathbb{Z}$$

Since B is injective in Y, this implies $y_h = 0$ and thus the assertion $\mathcal{N}(B_h) = \{0\}$.

Remark 7. If B_h is injective, a Fortin operator exists (see, e.g., [12] for a general Banach space case and [7] for the application to DPG). An optimal Fortin operator Π_h^{opt} can be determined as follows: for given $z \in Z$ find $z_h \in Z_h$ with minimal norm $||z_h||_Z$ subject to the constraint $b(y_h; z - z_h) = 0$ for all $y_h \in Y_h$. Again, this can be computed from a critical point $(z_h, y_h) \in Z_h \times Y_h$ of the corresponding Lagrange functional

$$F(z_h, y_h) = \frac{1}{2} ||z_h||_Z^2 + b(y_h; z - z_h),$$

i.e., $A_{Z_h}z_h = B_h y_h$, $B'_h z_h = E'_{Y_h}B'z$, and thus $z_h = A_{Z_h}^{-1}B_h y_h$ and $S_h y_h = B'_h A_{Z_h}^{-1}B_h y_h = B'_h z_h = E'_{Y_h}B'z$. This yields $z_h = \prod_h^{\text{opt}} z$ with

$$\Pi_h^{\text{opt}} = A_{Z_h}^{-1} B_h S_h^{-1} E'_{Y_h} B' \,.$$

Bounding Π_h^{opt} involves an estimate for S_h^{-1} which requires a bound for the inf-sup constant. The construction in Lem. 6 requires only local estimates for $S_{K,h}^+$ which can be computed by local discrete symmetric eigenvalue problems

$$A_{Y_{K,h}}S^+_{K,h}A_{Y_{K,h}}y_{K,h} = \lambda_{K,h}A_{Y_{K,h}}y_{K,h}, \qquad (\lambda_{K,h}, y_{K,h}) \in [0,\infty) \times Y_{K,h}.$$
(17)

Let $||(w_K, v_K)||_{Y_K} = \max \{ ||w_K||_{W_K}, ||v_K||_{V_K} \}$ be the norm in $Y_K = W_K \times V_K$. Lemma 8. If the constants $\alpha_{K,h} > 0$ satisfy

$$\|S_{K,h}^{+}\ell_{K,h}\|_{Y_{K}} \leq \alpha_{K,h} \|A_{Y_{K,h}}^{-1}\ell_{K,h}\|_{Y_{K}}, \qquad \ell_{K,h} \in Y_{K,h}', \ K \subset \Omega_{h},$$
(18)

the Fortin operator constructed in Lem. 6 is bounded by $\|\Pi_h\|_Z \leq \max \alpha_{K,h}$.

Solving the eigenvalue problem (17) yields the estimate $\alpha_{K,h} = \sqrt{2} \max \lambda_{K,h}$.

Proof. For $z_K \in Z_K$ we define $\ell_{K,h} = E'_{Y_{K,h}} B'_K z_K$ and $y_{K,h} = S^+_{K,h} \ell_{K,h}$, so that

$$\Pi_{K,h} z_{K} = A_{Z_{K,h}}^{-1} B_{K,h} S_{K,h}^{+} E'_{Y_{K,h}} B'_{K} z_{K}$$

$$= A_{Z_{K,h}}^{-1} B_{K,h} S_{K,h}^{+} \ell_{K,h}$$

$$= A_{Z_{K,h}}^{-1} B_{K,h} y_{K,h} .$$

The definition of the norm in Y_K yields $|b_K(y_{K,h}; z_{K,h})| = |\langle B_{K,h}y_{K,h}, z_{K,h}\rangle| \le ||y_{K,h}||_{Y_K} ||z_{K,h}||_{Z_K}, z_{K,h} \in Z_{K,h}$, which gives

$$\|\Pi_{K,h} z_K\|_{Z_K} = \|A_{Z_{K,h}}^{-1} B_{K,h} y_{K,h}\|_{Z_K} \le \|y_{K,h}\|_{Y_K}.$$

10

From (18) we we obtain $\|y_{K,h}\|_{Y_K} = \|S_{K,h}^+ \ell_{K,h}\|_{Y_K} \le \alpha_{K,h} \|A_{Y_{K,h}}^{-1} \ell_{K,h}\|_{Y_K}$, and finally it holds

$$\|A_{Y_{K,h}}^{-1}\ell_{K,h}\|_{Y_{K}} = \|A_{Y_{K,h}}^{-1}E'_{Y_{K,h}}B'_{K}z_{K}\|_{Y_{K}} \le \|z_{K}\|_{Z_{K}}.$$

Together this yields $\|\Pi_{K,h}z\|_{Z_{K,h}} \leq \alpha_{K,h}\|z\|_{Z_K}$. Then, $\|\Pi_h z\|_Z^2 \leq \sum \alpha_{K,h}^2 \|z|_K\|_{Z_K}^2 \leq \max \alpha_{K,h}^2 \|z\|_Z^2$ yields the assertion.

Remark 9. Scaling arguments on uniformly shape regular meshes in [20] and [15, Sect. 5.2] show that a bound for $\alpha_{K,h}$ can be estimated on the reference cell. Results for the acoustic wave equation on tensor-product space-time cells are presented in [15, Sect. 7.1]

5. The realization of the DPG method

In heterogeneous materials, the finite element error also depends on the approximation error of the PDE, and the realization of the DPG method uses an approximation $M_h \in L_{\infty}(\Omega_0; \mathbb{R}_{sym}^{m \times m})$ of M; then we set $L_h = M_h \partial_t + A$ and $L_h^* z = -L_h z$ for $z \in C_c^1(\Omega_h; \mathbb{R}^m)$.

The estimates for the DPG analysis use functionals in $D_{\Omega_h}(V) \subset Z'$. In the implementation, we use representations of these functionals in $\widetilde{W} = L_2(\partial \Omega_h; \mathbb{R}^m)$. For this purpose we introduce trace mappings for sufficiently smooth functions.

Since $\mathcal{N}(D_K) = \mathrm{H}_0(L, K) \supset \mathrm{C}^1_{\mathrm{c}}(K; \mathbb{R}^m)$, we can define trace mappings $\mathrm{tr}_K, \mathrm{tr}_K^* \in \mathcal{L}(\mathrm{C}^1(K; \mathbb{R}^m), \mathrm{L}_2(\partial K; \mathbb{R}^m))$ such that

$$(L_h v_K, z_K)_{0,K} - (L_h^* v_K, z_K)_{0,K} = (\operatorname{tr}_K v_K, \operatorname{tr}_K^* z_K)_{0,\partial K}, \qquad v_K, z_K \in \operatorname{C}^1(K; \mathbb{R}^m).$$

This extends to $\operatorname{tr}_h \in \mathcal{L}(\operatorname{C}^1(\Omega_h; \mathbb{R}^m) \cap V, \widetilde{W})$ and $\operatorname{tr}_h^* \in \mathcal{L}(\operatorname{C}^1(\Omega_h; \mathbb{R}^m), \widetilde{W})$ such that

$$(L_h v, z)_{0,\Omega_h} - (L_h^* v, z)_{0,\Omega_h} = (\operatorname{tr}_h v, \operatorname{tr}_h^* z)_{0,\partial\Omega_h}, \qquad v \in \mathrm{C}^1(\Omega_h; \mathbb{R}^m) \cap V, \ z \in \mathrm{C}^1(\Omega_h; \mathbb{R}^m)$$

by the selection of an orientation on inner faces $\overline{F} = \partial K \cap \partial K_F$ and the restriction $\operatorname{tr}_h v|_F = \operatorname{tr}_K v|_F$ for conforming functions, and the jump term $\operatorname{tr}_h^* z|_F = \operatorname{tr}_K z|_F - \operatorname{tr}_{K_F} z|_F$ for functions in the broken space (see [15, Sect. 6.1] for the acoustic case).

The trace defines the approximation of the bilinear form $b(\cdot; \cdot)$

$$b_h(w_h, \tilde{v}_h; z_h) = (w_h, L_h^* z_h)_{0,\Omega_h} + (\tilde{v}_h, \operatorname{tr}_h^* z_h)_{0,\partial\Omega_h}, \qquad w_h \in W, \ \tilde{v}_h \in \widetilde{W}, \ z_h \in Z_h.$$

Now we select $W_h \subset W$, $\widetilde{V}_h \subset \widetilde{W}$, and $Z_h \subset C^1(\Omega_h; \mathbb{R}^m)$ such that

$$\|\tilde{v}_{h}\|_{Z'_{h}} = \sup_{z_{h}\in Z_{h}} \frac{(\tilde{v}_{h}, \operatorname{tr}_{h}^{*} z_{h})_{0,\partial\Omega_{h}}}{\|z_{h}\|_{Z_{h}}}, \quad v_{h}\in \widetilde{V}_{h}$$

is a norm in \tilde{V}_h and such that the bilinear form $b_h(\cdot; \cdot)$ is inf-sup stable, i.e., $\beta_0 > 0$ exists such that

$$\sup_{z_h \in Z_h} \frac{b_h(w_h, \tilde{v}_h; z_h)}{\|z_h\|_{Z_h}} \ge \beta_0 \|(w_h, \tilde{v}_h)\|_{W_h \times Z'_h}, \qquad (w_h, \tilde{v}_h) \in W_h \times \widetilde{V}_h,$$
(19)

with respect to the norms for $w \in W$, $z \in Z$, and $(w_h, \tilde{v}_h) \in W_h \times \tilde{V}_h$ given by

$$\|w\|_{W_h} = \sqrt{(M_h w, w)_{0,\Omega_0}}, \quad \|z\|_{Z_h} = \sqrt{\|z\|_{W_h}^2 + \|M_h^{-1}L^* z\|_{W_h}^2}, \quad \|(w_h, \tilde{v}_h)\|_{W \times Z'_h} = \max\{\|w_h\|_W, \|\tilde{v}_h\|_{Z'_h}\}.$$

For the error analysis we construct a conforming extension $V_h \subset V$ of the trace space \tilde{V}_h so that for all $\tilde{v}_h \in \tilde{V}_h$ a reconstruction $v_h \in V_h$ exist satisfying

$$(\operatorname{tr}_{h} v_{h}, \operatorname{tr}_{h}^{*} z_{h})_{0,\partial\Omega_{h}} = (\tilde{v}_{h}, \operatorname{tr}_{h}^{*} z_{h})_{0,\partial\Omega_{h}}, \qquad z_{h} \in Z_{h}.$$

$$(20)$$

Then, we denote by $\widehat{\operatorname{tr}}_h v_h \in Z'_h$ the corresponding functional defined by

$$\langle \widehat{\operatorname{tr}}_h v_h, z_h \rangle = (\operatorname{tr}_h v_h, \operatorname{tr}_h^* z_h)_{0,\partial\Omega_h}, \qquad z_h \in Z_h,$$
(21)

and we set $\hat{V}_h = \hat{\mathrm{tr}}_h(V_h) \subset Z'_h$.

The DPG approximation $(u_h, \tilde{u}_h) \in W_h \times \tilde{V}_h$ for a given right-hand side $f \in W$ minimizes the residual

$$(u_h, \tilde{u}_h) = \arg\min_{(w_h, \tilde{v}_h) \in W_h \times \tilde{V}_h} \sup_{z_h \in Z_h} \frac{b_h(w_h, \tilde{v}_h; z_h) - (f, z_h)_{0,\Omega}}{\|z_h\|_{Z_h}}$$

It is computed by

$$b_h(u_h, \tilde{u}_h; z_h) = (f, z_h)_{0,\Omega}, \qquad z_h \in \tilde{T}_h(W_h \times \tilde{V}_h),$$
(22)

where the discrete trial-to-test operator $\tilde{T}_h \in \mathcal{L}(W_h \times \tilde{V}_h, Z_h)$ is defined by

$$\left(\tilde{T}_h(w_h,\tilde{v}_h),z_h\right)_Z = b_h(w_h,\tilde{v}_h;z_h), \qquad (w_h,\tilde{v}_h) \in W_h \times \widetilde{V}_h, \ z_h \in Z_h.$$

Now, the first Strang Lemma [11, Lem. 2.27] takes the following form.

Theorem 10. For $f \in W$ let $u \in V$ be the solution of Lu = f, and let $(u_h, \tilde{u}_h) \in W_h \times \tilde{V}_h$ be the DPG approximation solving (22). We assume that a conforming reconstruction space $V_h \subset V$ satisfying (20) and $\hat{u}_h \in \hat{V}_h$ exist with

$$\langle \hat{u}_h, z_h \rangle = (\tilde{u}_h, \operatorname{tr}_h^* z_h)_{0,\partial\Omega_h}, \qquad z_h \in Z_h.$$

$$(23)$$

Let $\widehat{\operatorname{tr}}_h u \in Z'_h$ be defined by

$$\langle \widehat{\mathrm{tr}}_h u, z_h \rangle = (L_h u, z_h)_{0,\Omega_h} - (u, L_h^* z_h)_{0,\Omega_h}, \qquad z_h \in Z_h,$$
(24)

and we set $\hat{u} = D_{\Omega_h} u \in Z'$. Then, the error is bounded by

$$\begin{aligned} \|(u - u_h, \widehat{\operatorname{tr}}_h u - \widehat{u}_h)\|_{W_h \times Z'_h} &\leq \inf_{(w_h, v_h) \in W_h \times V_h} \left(\|(u - w_h, \widehat{\operatorname{tr}}_h u - \widehat{\operatorname{tr}}_h v_h)\|_{W_h \times Z'_h} \\ &+ \beta_0^{-1} \sup_{z_h \in Z_h} \frac{b_h(w_h, \operatorname{tr}_h v_h; z_h) - b(u, \widehat{u}; z_h)}{\|z_h\|_{Z_h}} \right). \end{aligned}$$

Furthermore, if $u \in \mathcal{V}$, the error is bounded by

$$\begin{aligned} \|(u-u_h,\widehat{\mathrm{tr}}_h u - \widehat{u}_h)\|_{W_h \times Z'_h} &\leq (1+\beta_0^{-1}) \inf_{(w_h,v_h) \in W_h \times V_h} \|(u-w_h,\widehat{\mathrm{tr}}_h u - \widehat{\mathrm{tr}}_h v_h)\|_{W_h \times Z'_h} \\ &+ \beta_0^{-1} \|M_h^{1/2} (M_h - M) M^{-1/2}\|_{\infty} \|\partial_t u\|_W. \end{aligned}$$

Proof. For all $(w_h, v_h) \in W_h \times V_h$ we have

$$\|(u - u_h, \widehat{\mathrm{tr}}_h u - \widehat{u}_h)\|_{W_h \times Z'_h} \leq \|(u - w_h, \widehat{\mathrm{tr}}_h u - \widehat{\mathrm{tr}}_h v_h)\|_{W_h \times Z'_h} + \|(w_h - u_h, \widehat{\mathrm{tr}}_h v_h - \widehat{u}_h)\|_{W_h \times Z'_h},$$

and inserting (21), (23), and (19) gives

$$\begin{aligned} \|(w_h - u_h, \widehat{\operatorname{tr}}_h v_h - \widehat{u}_h)\|_{W_h \times Z'_h} &= \|(w_h - u_h, \operatorname{tr}_h v_h - \widetilde{u}_h)\|_{W_h \times Z'_h} \\ &\leq \beta_0^{-1} \sup_{z_h \in Z_h} \frac{b_h (w_h - u_h, \operatorname{tr}_h v_h - \widetilde{u}_h; z_h)}{\|z_h\|_{Z_h}} \,. \end{aligned}$$

Using $b_h(u_h, \tilde{u}_h; z_h) = (f, z_h)_{0,\Omega} = (Lu, z_h)_{0,\Omega} = b(u, \hat{u}; z_h)$ for $z_h \in \tilde{T}_h(W_h \times \tilde{V}_h)$ and

$$\sup_{z_h \in Z_h} \frac{b_h(w_h, \tilde{v}_h; z_h)}{\|z_h\|_{Z_h}} = \sup_{z_h \in \tilde{T}_h(W_h \times \tilde{V}_h)} \frac{b_h(w_h, \tilde{v}_h; z_h)}{\|z_h\|_{Z_h}} , \qquad (w_h, \tilde{v}_h) \in W_h \times \tilde{V}_h$$

yields first estimate. The second estimate follows from $b_h(w_h, \operatorname{tr}_h v_h; z_h) = (w_h, L_h^* z_h)_{0,\Omega_h} + \langle \widehat{\operatorname{tr}}_h v_h, z_h \rangle$ and

$$b_{h}(w_{h} - u_{h}, \operatorname{tr}_{h} v_{h} - \tilde{u}_{h}; z_{h}) = b_{h}(w_{h}, \operatorname{tr}_{h} v_{h}; z_{h}) - (Lu, z_{h})_{0,\Omega}$$

$$= (w_{h}, L_{h}^{*} z_{h})_{0,\Omega_{h}} + \langle \widehat{\operatorname{tr}}_{h} v_{h}, z_{h} \rangle + \left((M_{h} - M) \partial_{t} u, z_{h} \right)_{0,\Omega} - (L_{h} u, z_{h})_{0,\Omega}$$

$$= (w_{h} - u, L_{h}^{*} z_{h})_{0,\Omega_{h}} + \langle \widehat{\operatorname{tr}}_{h} v_{h} - \widehat{\operatorname{tr}}_{h} u, z_{h} \rangle + \left((M_{h} - M) \partial_{t} u, z_{h} \right)_{0,\Omega}.$$

Remark 11. In the case that the solution $u \in V$ has the additional regularity $u \in \mathrm{H}^{1+s}(\Omega; \mathbb{R}^m)$ with $s > k \ge 0$, a trace function $\tilde{u} \in \mathrm{L}_2(\partial \Omega_h; \mathbb{R}^m)$ with

$$(\tilde{u}, \operatorname{tr}_{h}^{*} z)_{0,\partial\Omega_{h}} = (Lu, z)_{0,\Omega_{h}} - (u, L^{*}z)_{0,\Omega_{h}}, \qquad z \in \operatorname{C}^{1}(\Omega_{h}; \mathbb{R}^{m})$$

exists. In this case, the proof of Thm. 10 only relies on the traces in L₂, and no conforming reconstruction space V_h is required for the estimates. For $W_h \supset \mathbb{P}_k(\Omega_h)$ we obtain

$$\inf_{w_h \in W_h} \|u - w_h\|_{W_h} \le \|M_h\|_{\infty}^{1/2} \inf_{w_h \in W_h} \|u - w_h\|_{0,\Omega} \le C_1 \|M_h\|_{\infty}^{1/2} h^s \|u\|_{1+s,\Omega}$$

and for $\widetilde{V}_h \supset \mathbb{P}_{k+1}(\Gamma_h)$ with $\Gamma_h = \bigcup F \subset \partial \Omega_h$, we obtain, provided that $M, M_h \in \mathcal{L}_{\infty}(\partial \Omega_h; \mathbb{R}^{m \times m}_{sym})$, the estimate

$$\inf_{v_h \in V_h} \|\widehat{\mathrm{tr}}_h u - \widehat{\mathrm{tr}}_h v_h\|_{Z'_h} = \inf_{\tilde{v}_h \in \widetilde{V}_h} \|\widetilde{u} - \widetilde{v}_h\|_{Z'_h} \le C_2 h^{-1/2} \inf_{\tilde{v}_h \in \widetilde{V}_h} \|\widetilde{u} - \widetilde{v}_h\|_{0,\partial\Omega_h} \le C_3 h^s \|u\|_{1+s,\Omega_h}$$

with constants $C_1, C_2, C_3 > 0$ depending on the shape regularity of Ω_h and on M_h . Together, this results in the convergence estimate

$$\|(u-u_h,\widehat{\mathrm{tr}}_hu-\hat{u}_h)\|_{W_h\times Z'_h} \le (1+\beta_0^{-1})\max\{C_1,C_3\}h^s\|u\|_{1+s,\Omega} + \beta_0^{-1}\|M_h^{1/2}(M_h-M)M^{-1/2}\|_{\infty}\|\partial_t u\|_W.$$

Remark 12. Choosing an extended space $Z_h^{\text{ext}} \supset Z_h$ allows for a conforming reconstruction $V_h \subset Z_h^{\text{ext}} \cap V$ satisfying (20), see [15, Sect. 6] for the local construction and for examples in case of acoustic waves.

In the nonconforming case $\tilde{V} \not\subset \operatorname{tr}_h(V_h)$, the construction of the reconstruction space V_h depends on Z_h , and the asymptotic arguments in Lem. 5 do not apply. In particular, the reconstruction space may have strong oscillations at the corners and edges of ∂K . The numerical experiments in [14] indicate that the optimal choice of Z_h has to be well balanced to ensure discrete inf-sup stability on the one hand and to limit the nonconformity on the other hand. Nevertheless, for a given choice of W_h , \tilde{V}_h , and Z_h the well-posedness of the discrete system can be guaranteed by Lem. 6, and explicitly computing a reconstruction V_h a lower bound for the inf-sup constant can be provided by Lem. 8.

6. A numerical example for an application in geophysics

Many applications rely on accurate numerical simulations of waves through complex material structures. For instance, geophysical structures like the earth's crust below the sea bed feature complex varying material properties. A typical example is the problem of full waveform inversion (FWI), where the material distribution is reconstructed from measurements of the wave field close to the surface [23]. Here, in a field survey a wave is excited at some point $S_0 \in \Omega_0$, and the scattered wave field is measured by receiver devices located at $R_0, \ldots, R_N \in \Omega_0$. During the experiment, each receiver records a time series of approximate point measurements $u(t, R_n), t \in [0, T]$. The collection of all these measurements is called a seismogram, see Fig. 5 for examples.

The recorded seismogram contains information about the material structure the wave has traveled through. Full waveform inversion techniques attempt to reconstruct from this information by applying iterative schemes of Newton-type (see, e.g., [19]). During the iteration, a large number of wave equations has to be solved numerically for different right-hand sides and varying material parameters.

To demonstrate the flexibility and the accuracy of the space-time DPG method for heterogeneous media, we consider a numerical example corresponding to the forward problem within FWI. We use the acoustic wave equation for the Marmousi benchmark, a synthetic model problem for geophysical structures in two space dimensions featuring a material distribution that is similar to what is located inside the earth crust (see, e.g., [5]). Fig. 3 shows rescaled variants of the spatial varying mass density and bulk modulus that have similar structure to the materials in this benchmark. We select the spatial domain $\Omega_0 \subset \mathbb{R}^2$, the receiver positions R_0, \ldots, R_N , and the source position $S = (t_0, S_0) \in (0, T) \times \Omega_0$ as shown in Fig. 2.



FIGURE 2. The left figure shows the spatial domain $\Omega_0 = (0, 9) \times (0, 3)$, the source position $S_0 = (2.602, 0.802)$, and the first and last receiver $R_0 = (2.002, 0.303)$ and $R_N = (4.002, 0.303)$. In space-time, the source is located at $S = (0.4, S_0)$ and the signal is measured in $\mathcal{R} =$ conv $(\{(t, R_0), \ldots, (t, R_N): t \in (0, T)\}), T = 8$, which results in the domain of interest intersecting of the forward and backward cone $\mathcal{C}_+(\{S\}) \cap \mathcal{C}_-(\mathcal{R})$. This is illustrated on the right.

For the inversion, we only require the wave field in the space-time region that contributes to the transport of information from the source to the receiver array. Thus, using Lem. 2 for the maximal wave speed $c_{\text{max}} = 1.37$, we find a superset of the relevant region of the full space-time domain $(0, T) \times \Omega_0$, see Fig. 2. At the source position S, the wave is excited by the right space for $f(x, t) = 100 \phi_t(t_0 - t)\phi_x(|S_0 - x|), (t, x) \in (0, T) \times \Omega_0$ with a Ricker wavelet for $f_0 = 100$ in time

$$\phi_t(\tau) = (1 - 2a\tau^2) \exp(-a\tau^2), \quad a = \pi^2 f_0^2 \tau^2,$$

and in space $\phi_x(r) = \chi_{[0,r_0)}(r) \cos\left(\frac{\pi}{2} \frac{r}{r_0}\right)^6$ for $r_0 = 0.05$.

The numerical simulations were performed using the DPG approximations with local finite element spaces

$$W_{K,h} = \mathbb{Q}_3(K)^3, \quad \tilde{V}_h|_{(t_{n-1},t_n)\times\partial K} = \mathbb{Q}_4(F)^3, \quad \tilde{V}_h|_{\{t_n\}\times K_0} = \mathbb{Q}_4(F)^2, \qquad Z_{K,h} = \mathbb{Q}_6(K)^3$$

on a quadrilateral mesh, yielding a scheme that converges with order 4 in $L_2((0,T) \times \Omega_0)$ for smooth solutions, see [14, Sec. 4.7]. The material parameters κ and ρ are cell-wise constant as shown in Fig. 3.

The method is realized in the parallel Finite Element system M++ [24], where the linear systems are solved with a preconditioned cg method using the reduction to the symmetric positive definite Schur complement system for the skeleton approximation \tilde{V}_h , cf. [25]. The implementation of this space-time DPG method is evaluated systematically in [14, Sec. 5] for configurations where the analytic solution is known. Moreover, basic applications of space-time DPG to FWI are discussed in [14, Sec. 6].



FIGURE 3. Approximations of rescaled spatial material distributions for Marmousi, κ on the left and ρ on the right. The first row corresponds to the discrete material parameters on a coarse quadrilateral mesh with $N_x \cdot N_y = 144 \cdot 48 = 6\,912$ cells, the second row shows the material on a uniformly refined mesh with $N_x \cdot N_y = 288 \cdot 96 = 27\,648$ cells.

The pressure component of the full numerical space-time solution is shown in Fig. 4, also see Fig. 6. This is compared with the solution on a mesh that has been truncated to the space-time region of interest. The seismograms corresponding to the materials are shown in Fig. 5.

The numerical experiments with the truncated space-time cylinder correspond to our results in Lem. 2. Up to a small relative $L_2(0, T; \mathbb{R}^3)$ difference of 0.00318, the seismograms on the full mesh and the truncated mesh coincide. The linear system of all space-time unknowns is solved approximately using an iterative solver, which can explain the small difference. On the other hand, only explicit time stepping methods guarantee finite speed of the discrete wave propagation. Since DPG is an implicit scheme, this also may explain a small difference of the results for the full space-time cylinder and the truncated domain.

The comparison of the solutions on different refinement levels of the mesh shows that both seismograms have a similar structure at the beginning but differ strongly at later times. Here, we give two possible reasons for this difference: on the one hand, the approximation quality for the wave field increases on mesh refinement, and on the other hand, on the finer mesh we have a better resolution of the material parameters M_h , i.e., a different wave equation is solved, see Thm. 10.



FIGURE 4. The left picture shows the pressure component of the numerical solution on a regular grid with $N_t \cdot N_x \cdot N_y = 128 \cdot 144 \cdot 48 = 884736$ space-time cells resulting in 156576000 space-time DoFs for all components in the skeleton space \tilde{V}_h . On the right, a truncated version of the space-time cylinder is depicted that respects the position of the source and the receivers. The truncation reduced the amount of cells by approx. 43% to 503024 cells and 89394900 DoFs.

Conclusion and Outlook. In this example we show that the method also yields accurate results for application driven simulations. We demonstrate that for wave-type equations the finite wave speed can be exploited to reduce the size of the linear system considerably while yielding the same results. Further numerical experiments are necessary to compare the roles of the discretization error for fixed material M and $h \rightarrow 0$ to the model error resulting from the approximate material M_h . For that reason, we intend to evaluate the performance of an established discretization like discontinuous Galerkin with time stepping in comparison to the space-time DPG method.

Acknowledgment. We gratefully acknowledge financial support by the Deutsche Forschungsgemeinschaft (DFG) through CRC 1173. This work was performed on the supercomputer ForHLR funded by the Ministry of Science, Research and the Arts Baden-Württemberg and by the Federal Ministry of Education and Research.

References

- K. D. Blazek, C. Stolk, and W. W. Symes. A mathematical framework for inverse wave problems in heterogeneous media. *Inverse Problems*, 29(6):065001, 2013.
- [2] D. Braess. Finite Elements. Theory, fast solvers, and applications in solid mechaics. 3th ed. Cambridge University Press, 2007.
- [3] F. Brezzi and M. Fortin. Mixed and Hybrid Finite Element Methods. Springer, 1991.
- [4] D. Broersen, W. Dahmen, and R. Stevenson. On the stability of DPG formulations of transport equations. Mathematics of Computation, 87(311):1051–1082, 2018.



FIGURE 5. All plots show measurements of the form $t \mapsto \int \phi(s-t, x-R_n) \cdot u(s, x) d(s, x)$ of the pressure component at the corresponding receivers. Here ϕ is a measurement kernel with small support in space and time. On the left-hand side, a comparison of the seismograms on the full mesh (solid) and the truncated mesh (dotted) is depicted. At the bottom, the recording at R_0 is shown for illustration purposes. Since the difference is very small (relative $L_2(0, T; \mathbb{R}^3)$ difference is 0.00318), both seismograms look very similar. On the right, we compare the seismogram corresponding to the truncated mesh (solid) to the seismogram obtained on a mesh that has been uniformly refined (dotted). While being similar to the measurements on the coarser grid, we see a different fine structure due to the different material approximation M_h .



FIGURE 6. Pressure distribution for three different numerical simulations at selected time steps t = 0.4, 1.49, 2.57, 3.66, 4.74, 5.83, 6.91, 8. The first column shows the results for the space-time DPG method using the full space-time cylinder on level 4. The second and third column correspond to space-time DPG in the truncated space-time cylinder on level 4 and level 5 respectively. To solve the linear systems, we use a parallel cg iteration with a block Jacobi preconditioner. The full problem on the left with $1.57 \cdot 10^8$ skeleton degrees of freedom is solved on 1000 cores in about 40 minutes. The truncated problems in the middle and on the right with $8.9 \cdot 10^7$ and $7.1 \cdot 10^8$ skeleton degrees of freedom are solved on 5000 cores in about 6 min and 75 min, respectively. On level 4, less than 6000 iterations and on level 5, less than 12000 iterations are sufficient to reduce the residual by a factor of $2 \cdot 10^{-8}$.

- [5] A. Brougois, M. Bourget, P. Lailly, M. Poulet, P. Ricarte, and R. Versteeg. Marmousi, model and data. In EAEG Workshop-Practical Aspects of Seismic Data Inversion, 1990.
- [6] C. Carstensen, L. Demkowicz, and J. Gopalakrishnan. Breaking spaces and forms for the DPG method and applications including Maxwell equations. *Computers & Mathematics with Applications*, 2016.
- [7] C. Carstensen and F. Hellwig. Low-order discontinuous Petrov–Galerkin finite element methods for linear elasticity. SIAM Journal on Numerical Analysis, 54(6):3388–3410, 2016.
- [8] R. Dautray and J.-L. Lions. Mathematical analysis and numerical methods for science and technology Volume 5: Evolution problems I. Springer Science & Business Media, 2012.
- [9] L. F. Demkowicz and J. Gopalakrishnan. An overview of the discontinuous Petrov–Galerkin method. In Recent Developments in Discontinuous Galerkin Finite Element Methods for Partial Differential Equations, pages 149–180. Springer, 2014.
- [10] L. F. Demkowicz, J. Gopalakrishnan, S. Nagaraj, and P. Sepulveda. A spacetime DPG method for the Schrödinger equation. SIAM J. Numer. Anal., 55(4):1740–1759, 2017.
- [11] A. Ern and J.-L. Guermond. *Theory and practice of finite elements*, volume 159. Springer Science & Business Media, 2013.
- [12] A. Ern and J.-L. Guermond. A converse to Fortin's Lemma in Banach spaces. Comptes Rendus Mathematique, 354(11):1092–1095, 2016.
- [13] A. Ern, J.-L. Guermond, and G. Caplain. An intrinsic criterion for the bijectivity of Hilbert operators related to Friedrichs systems. *Communications in partial differential equations*, 32(2):317–341, 2007.
- [14] J. Ernesti. Space-Time Methods for Acoustic Waves with Applications to Full Waveform Inversion. PhD thesis, Karlsruher Institut f
 ür Technologie (KIT), 2018.
- [15] J. Ernesti and C. Wieners. A space-time DPG method for acoustic waves. In U. Langer and O. Steinbach, editors, Space-Time Methods. Applications to Partial Differential Equations, volume 21 of Radon Series on Computational and Applied Mathematics. Walter de Gruyter, 2018. submitted, preprint http://www.math.kit.edu/user/~wieners/SpaceTimeDPG_revised.pdf.
- [16] L. C. Evans. Partial differential equations, 2. ed. American Mathematical Society, Providence, RI, 2010.
- [17] J. Gopalakrishnan and P. Sepulveda. A spacetime DPG method for acoustic waves. arXiv:1709.08268, 2017.
- [18] M. Hochbruck, T. Pazur, A. Schulz, E. Thawinan, and C. Wieners. Efficient time integration for discontinuous Galerkin approximations of linear wave equations. ZAMM, 95(3):237–259, 2015.
- [19] L. Métivier, R. Brossier, S. Operto, and J. Virieux. Acoustic multi-parameter FWI for the reconstruction of p-wave velocity, density and attenuation: preconditioned truncated Newton approach. In SEG Technical Program Expanded Abstracts 2015, pages 1198–1203. Society of Exploration Geophysicists, 2015.
- [20] S. Nagaraj, S. Petrides, and L. F. Demkowicz. Construction of DPG Fortin operators for second order problems. Computers & Mathematics with Applications, 2017.
- [21] M. Renardy and R. C. Rogers. An introduction to partial differential equations, volume 13. Springer Science & Business Media, 2006.
- [22] W. Rudin. Functional Analysis. International Series in Pure and Applied Mathematics. McGraw-Hill, Inc., New York, 1991.
- [23] W. Symes. The seismic reflection inverse problem. Inverse problems, 25(12):123008, 2009.
- [24] C. Wieners. A geometric data structure for parallel finite elements and the application to multigrid methods with block smoothing. *Computing and visualization in science*, 13(4):161–175, 2010.
- [25] C. Wieners. The skeleton reduction for finite element substructuring methods. In Numerical Mathematics and Advanced Applications ENUMATH 2015, pages 133–141. Springer, 2016.
- [26] C. Wieners and B. Wohlmuth. Robust operator estimates and the application to substructuring methods for first-order systems. ESAIM: M²AN, 48:161–175, 2014.