

Biharmonic wave maps: local wellposedness in high regularity

Sebastian Herr, Tobias Lamm,
Tobias Schmid, Roland Schnaubelt

CRC Preprint 2019/2 (revised), March 2019

KARLSRUHE INSTITUTE OF TECHNOLOGY

CRC 1173



Participating universities



Universität Stuttgart

EBERHARD KARLS
UNIVERSITÄT
TÜBINGEN



Funded by

DFG

ISSN 2365-662X

BIHARMONIC WAVE MAPS: LOCAL WELLPOSEDNESS IN HIGH REGULARITY

SEBASTIAN HERR, TOBIAS LAMM, TOBIAS SCHMID AND ROLAND SCHNAUBELT

ABSTRACT. We show the local wellposedness of biharmonic wave maps with initial data of sufficiently high Sobolev regularity and a blow-up criterion in the sup-norm of the gradient of the solutions. In contrast to the wave maps equation we use a vanishing viscosity argument and an appropriate parabolic regularization in order to obtain the existence result. The geometric nature of the equation is exploited to prove convergence of approximate solutions, uniqueness of the limit, and continuous dependence on initial data.

1. INTRODUCTION

Let (N, g) be a smooth and compact Riemannian manifold which we assume to be isometrically embedded into some Euclidean space \mathbb{R}^L . Biharmonic wave maps are critical points $u : \mathbb{R}^n \times [0, T] \rightarrow N$ of the (extrinsic) action functional

$$(1.1) \quad \Phi(u) = \frac{1}{2} \int_0^T \int_{\mathbb{R}^n} |\partial_t u|^2 - |\Delta u|^2 \, dx \, ds.$$

These maps model the movement of a thin, stiff, elastic object within the target manifold N .

The Euler-Lagrange equation of Φ has been calculated in [4] (in the case $N = S^l \subset \mathbb{R}^{l+1}$) and in [9] (for arbitrary N) and it is given by

$$(1.2) \quad \partial_t^2 u + \Delta^2 u \perp T_u N \quad \text{on } \mathbb{R}^n \times [0, T].$$

In order to obtain a more explicit form of this equation we use the fact that there exists some $\delta_0 > 0$ and a smooth family of linear maps $P_p : \mathbb{R}^L \rightarrow \mathbb{R}^L$ for $\text{dist}(p, N) < \delta_0$ such that

$$P_p : \mathbb{R}^L \rightarrow T_p N, \quad p \in N,$$

is an orthogonal projection onto the tangent space $T_p N$. The Euler-Lagrange equation (1.2) can thus be written as

$$\partial_t^2 u + \Delta^2 u = (I - P_u)(\partial_t^2 u + \Delta^2 u).$$

Exploiting that u takes values in N , we calculate

$$(1.3) \quad \begin{aligned} \partial_t^2 u + \Delta^2 u &= dP_u(u_t, u_t) + dP_u(\Delta u, \Delta u) + 4dP_u(\nabla u, \nabla \Delta u) + 2dP_u(\nabla^2 u, \nabla^2 u) \\ &\quad + 2d^2 P_u(\nabla u, \nabla u, \Delta u) + 4d^2 P_u(\nabla u, \nabla u, \nabla^2 u) \\ &\quad + d^3 P_u(\nabla u, \nabla u, \nabla u, \nabla u) \\ &=: \mathcal{N}(u), \end{aligned}$$

where the tensors $d^j P$ are described in the next section more explicitly. The main goal of this paper is to show the following local wellposedness result for the Cauchy problem corresponding to (1.2) in Sobolev spaces with sufficiently high regularity.

Date: March 5, 2019.

TL, TS and RS gratefully acknowledge financial support by the Deutsche Forschungsgemeinschaft (DFG) through CRC 1173.

Theorem 1.1. *Let $u_0, u_1 : \mathbb{R}^n \rightarrow \mathbb{R}^L$ satisfy $u_0(x) \in N$ and $u_1(x) \in T_{u_0(x)}N$ for a.e. $x \in \mathbb{R}^n$ as well as*

$$(\nabla u_0, u_1) \in H^{k-1}(\mathbb{R}^n) \times H^{k-2}(\mathbb{R}^n)$$

for some $k \in \mathbb{N}$ with $k > \lfloor \frac{n}{2} \rfloor + 2$. Then the following assertions hold:

(a) *There exists a maximal existence time*

$$T_m = T_m(u_0, u_1) > T = T(\|\nabla u_0\|_{H^{k-1}}, \|u_1\|_{H^{k-2}}) > 0$$

and a unique solution $u : \mathbb{R}^n \times [0, T_m) \rightarrow N$ of (1.2) with $u(0) = u_0$, $\partial_t u(0) = u_1$, and

$$u - u_0 \in C^0([0, T_m), H^k(\mathbb{R}^n)) \cap C^1([0, T_m), H^{k-2}(\mathbb{R}^n)).$$

(b) *For $T_0 \in (0, T_m)$ there exists a (sufficiently small) radius $R_0 > 0$ such that for all initial data (v_0, v_1) as above that satisfy*

$$\|(u_0, u_1) - (v_0, v_1)\|_{H^k(\mathbb{R}^n) \times H^{k-2}(\mathbb{R}^n)} \leq R_0,$$

the unique solution $v : \mathbb{R}^n \times [0, T_m(v_0, v_1)) \rightarrow N$ exists on $\mathbb{R}^n \times [0, T_0]$. Further, for such initial data the map $(v_0, v_1) \rightarrow (v(t), \partial_t v(t))$ is continuous in $H^k(\mathbb{R}^n) \times H^{k-2}(\mathbb{R}^n)$ for $t \in [0, T_0]$.

(c) *If $T_m < \infty$, then*

$$(1.4) \quad \int_0^{T_m} \|\nabla u(s)\|_{L^\infty}^{2k} + \|u_t(s)\|_{L^\infty}^{2k} ds = \infty.$$

In particular, for smooth initial data $u_0, u_1 : \mathbb{R}^n \rightarrow \mathbb{R}^L$ with $u_0(x) \in N$ and $u_1(x) \in T_{u_0(x)}N$ for $x \in \mathbb{R}^n$ having compact $\text{supp}(\nabla u_0) \subset \mathbb{R}^n$ and $\text{supp}(u_1) \subset \mathbb{R}^n$, there exist $T_m > 0$ and a smooth solution $u : \mathbb{R}^n \times [0, T_m) \rightarrow N$ of (1.2).

It is worthwhile to remark that both u_0 and $u(t)$ do not necessarily belong to $L^2(\mathbb{R}^n)$ and it is only the difference of these two functions which is contained in this space.

The first, second and fourth author have recently shown in [4] that there exists a global weak solution of (1.2) for initial data in the energy space $H^2 \times L^2$ in the case $N = S^l \subset \mathbb{R}^{l+1}$. In [4] a crucial ingredient is a conservation law which allows to construct the desired solution as a weak limit of a sequence of solutions of suitably regularized problems. The derivation of this conservation law relies on the fact that the action functional Φ is invariant under rotations in the highly symmetric setting $N = S^l$, and this argument does not apply to arbitrary target manifolds N .

Moreover, the third author has shown energy estimates for biharmonic wave maps in low dimensions $n = 1, 2$ in [9]. When combining this result with the above blow-up criterion (1.4), he then obtained the existence of a unique global smooth solution of (1.2) for smooth and compactly supported initial data. This result extends earlier work of Fan and Ozawa [3] for spherical target manifolds.

A local well-posedness result as in Theorem 1.1 is standard for second-order wave equations with derivative nonlinearities such as wave maps. It can be found for example in the books of Shatah and Struwe [11] and Sogge [10]. In contrast to this case, our nonlinearity $\mathcal{N}(u)$ depends on the third spatial derivative of u which cannot directly be controlled by the energy of (1.3) that only contains second order spatial derivatives. In our proof we use the geometric nature of the equation in several crucial steps in order to be able to rewrite this expression in terms of derivatives of lower order. For this reason many of our arguments are fairly delicate.

Concerning the continuous dependence of the solution on the initial data, as the nonlinearity $\mathcal{N}(u)$ depends on the third spatial derivative, there seems to be no Lipschitz estimate available in the norm $H^k \times H^{k-2}$ at least from the energy method (as we observe e.g. from the a priori estimates in Section 6). This makes the wellposedness issue more involved compared to semi-linear wave equations with derivative nonlinearities such as wave maps.

In the following we briefly outline the structure of the paper. Since the nonlinearity $\mathcal{N}(u)$ in equation (1.3) contains derivatives of up to third order we cannot directly apply

the energy estimates for the operator $\partial_t^2 + \Delta^2$ and construct the desired solution by means of a fixed-point argument. Instead, in Section 3, we use a vanishing viscosity approximation and solve the corresponding Cauchy problem for the damped problem

$$\partial_t^2 u + \Delta^2 u - \varepsilon \Delta \partial_t u \perp T_u N, \quad \varepsilon \in (0, 1].$$

In order to obtain a limiting solution for (1.2) as $\varepsilon \searrow 0$, we prove a priori energy estimates which are uniform in ε in Section 4. As a byproduct we obtain the blow up criterion in Theorem 1.1. The existence part in Theorem 1.1 is then shown in Section 5, and in Section 6 we prove that the solutions are unique. Finally we establish the continuity of the flow map in Section 7. Before we proceed, we comment on a related action functional and its Euler-Lagrange equation.

The intrinsic version of a biharmonic wave map.

The functional Φ has an intrinsic analogue Ψ defined by

$$\Psi(u) = \frac{1}{2} \int_0^T \int_{\mathbb{R}^n} |\partial_t u|^2 - |(\Delta u)^T|^2 dx ds,$$

where $(\Delta u)^T = P_u(\Delta u)$ is the tension field of a smooth function $u : \mathbb{R}^n \times [0, T] \rightarrow N$. In contrast to Φ , the functional Ψ is independent of the embedding of $N \hookrightarrow \mathbb{R}^L$. Compared to the right hand side of (1.3) the Euler-Lagrange equation differs by

$$(1.5) \quad P_u(dP_u(\nabla u, \nabla u) \cdot d^2 P_u(\nabla u, \nabla u, \cdot)) + P_u(\operatorname{div}(dP_u(\nabla u, \nabla u) \cdot dP_u(\nabla u, \cdot))).$$

Since these terms are of lower order in the analysis of the following sections, as in Theorem 1.1 one can derive the existence and uniqueness of solutions of the modified version of (1.3). We hence deduce the existence of local unique intrinsic biharmonic wave maps with initial data as in Theorem 1.1. However, we do not have a result for initial data with (only) covariant derivatives in L^2 .

2. NOTATION AND PRELIMINARIES

We write C for a generic constant only depending on N , n and k , and often also $\lesssim \dots$ instead of $\leq C(\dots)$. We note that the projectors P_p defined in the introduction are derivatives of the metric distance (with respect to N) in \mathbb{R}^L , i.e.,

$$(2.1) \quad p = \pi(p) + \frac{1}{2} \nabla_p(\operatorname{dist}^2(p, N)), \quad P_p = d_p \pi(p), \quad \operatorname{dist}(p, N) < \delta_0.$$

Moreover, if $p \in \mathbb{R}^L$ is sufficiently close to N , then π has the nearest point property, i.e., $|\pi(p) - p| = \inf_{q \in N} |q - p|$, and hence

$$d\pi|_p = d\pi(p) = d(\pi^2(p)) = d\pi|_{\pi(p)} d\pi|_p.$$

Therefore $P_p : \mathbb{R}^L \rightarrow T_{\pi(p)} N$ is well-defined. Using cut-off functions we extend the identity (2.1), and thus also the equation $P_p = d_p \pi(p)$, to all of \mathbb{R}^L . Moreover, all derivatives of P_p are bounded on \mathbb{R}^n . In this way one can investigate (1.3) without restricting the coefficients a priori. Further, for $l \in \mathbb{N}_0$ we denote by $d^l P_p$ the derivative of order l of the map P_p , which is a $(l+1)$ -linear form on \mathbb{R}^L . For the coefficients in the standard coordinates in \mathbb{R}^L we write

$$(d^j P_u)_{l_0, \dots, l_j}^k = \frac{\partial}{\partial p_{l_1}} \dots \frac{\partial}{\partial p_{l_j}} (P_p)_{l_0}^k(u).$$

We now derive (1.3) from the condition (1.2) for smooth solutions $u : \mathbb{R}^m \times [0, T] \rightarrow N$. Note that we use the sum convention, i.e. the same indices in super-/subscript means summation.

Since $\partial_t u \in T_u N$, we infer the identity

$$\begin{aligned} [(I - P_u)(\partial_t^2 u)]^k &= (\delta_l^k - (P_u)_l^k)(\partial_t^2 u^l) = \partial_t(\delta_l^k - (P_u)_l^k)(\partial_t u^l) + (\partial_m P_u)_l^k \partial_t u^l \partial_t u^m \\ &= (dP_u)_{m,l}^k \partial_t u^l \partial_t u^m \end{aligned}$$

for $k = 1, \dots, L$. Because of $\nabla u \in T_u N$, we also obtain

$$\begin{aligned} [(I - P_u)(\Delta u)]^k &= \partial^{x^\alpha} (\delta_l^k - (P_u)_l^k) (\partial_{x_\alpha} u^l) + (\partial_m P_u)_l^k \partial^{x^\alpha} u^l \partial_{x_\alpha} u^m \\ &= (dP_u)_{m,l}^k \partial^{x^\alpha} u^l \partial_{x_\alpha} u^m, \end{aligned}$$

and hence

$$\begin{aligned} [(I - P_u)(\Delta^2 u)]^k &= \Delta((dP_u)_{m,l}^k \partial^{x^\alpha} u^l \partial_{x_\alpha} u^m) + \partial^{x^\alpha} ((dP_u)_{m,l}^k \Delta u^l \partial_{x_\alpha} u^m) \\ &\quad + (dP_u)_{m,l}^k (\partial^{x^\alpha} \Delta u^l) \partial_{x_\alpha} u^m. \end{aligned}$$

The symmetry of the indices then implies

$$\begin{aligned} [(I - P_u)(\Delta^2 u)]^k &= (d^3 P_u)_{l_0, l_1, l_2, l_3}^k \partial^{x^\alpha} u^{l_0} \partial_{x_\alpha} u^{l_1} \partial^{x^\beta} u^{l_2} \partial_{x_\beta} u^{l_3} \\ &\quad + 2(dP_u)_{l_0, l_1}^k \partial_{x_\alpha} \partial^{x^\beta} u^{l_0} \partial^{x^\alpha} \partial_{x_\beta} u^{l_1} + (dP_u)_{l_0, l_1}^k \Delta u^{l_0} \Delta u^{l_1} \\ &\quad + 2(d^2 P_u)_{l_0, l_1, l_2}^k \partial^{x^\alpha} u^{l_0} \partial_{x_\alpha} u^{l_1} \Delta u^{l_2} + 4(dP_u)_{l_0, l_1}^k \partial^{x^\alpha} \Delta u^{l_0} \partial_{x_\alpha} u^{l_1} \\ &\quad + 4(d^2 P_u)_{l_0, l_1, l_2}^k \partial^{x^\alpha} u^{l_0} \partial_{x_\alpha} \partial^{x^\beta} u^{l_1} \partial_{x_\beta} u^{l_2}. \end{aligned}$$

We briefly state the expressions from (1.5) in coordinates, i.e.,

$$\begin{aligned} [P_u(dP_u(\nabla u, \nabla u) \cdot d^2 P_u(\nabla u, \nabla u, \cdot))]^l &= \sum_j (P_u)_j^l dP_u(\nabla u, \nabla u) \cdot (d^2 P_u)_{k,m,j} \partial_{x_\alpha} u^k \partial^{x^\alpha} u^m, \\ [P_u(\operatorname{div}(dP_u(\nabla u, \nabla u) \cdot dP_u(\nabla u, \cdot)))]^l &= \sum_j (P_u)_j^l \partial^{x^\alpha} (dP_u(\nabla u, \nabla u) \cdot (dP_u)_{kj} \partial_{x_\alpha} u^k) \end{aligned}$$

for $l = 1, \dots, L$. In the following we use the shorthand $\nabla^{k_1} u \star \nabla^{k_2} u$ for (linear combinations of) products of partial derivatives of the components u^l of u for $l = 1, \dots, L$. Here the partial derivatives are of order $k_1 \in \mathbb{N}$ and $k_2 \in \mathbb{N}$, respectively. With this notation we can rewrite equation (1.3) as

$$\begin{aligned} \partial_t^2 u + \Delta^2 u &= dP_u(u_t, u_t) + dP_u(\nabla^2 u \star \nabla^2 u) + dP_u(\nabla^3 u \star \nabla u) \\ &\quad + d^2 P_u(\nabla u \star \nabla u \star \nabla^2 u) + d^3 P_u(\nabla u \star \nabla u \star \nabla u \star \nabla u). \end{aligned}$$

The Leibniz formula implies the following identity.

Lemma 2.1. *For $m \in \mathbb{N}$ and $l \in \mathbb{N}_0$ we have*

$$(2.2) \quad \nabla^m (d^l P_u) = \sum_{j=1}^m \sum_{\sum_{k=1}^j m_k = m-j} d^{j+l} P_u(\nabla^{m_1+1} u \star \dots \star \nabla^{m_j+1} u).$$

In order to include the case $m = 0$ in the lemma, we will use $\sum_{j=\min\{1, m\}}^m$ for the sum in (2.2) or similar formulas. The calculation of derivatives $\nabla^m(\mathcal{N}(u))$ and $\nabla^m(\mathcal{N}(u) - \mathcal{N}(v))$ for sufficiently regular $u, v : \mathbb{R}^n \times [0, T] \rightarrow \mathbb{R}^L$ and $m \in \mathbb{N}_0$ has been included in Appendix A, employing the \star -convention. The results from Appendix A will be used frequently throughout the paper. In the following sections, we also need a version of the classical Moser estimate, see e.g. [12, Chapter 13].

Lemma 2.2. *Let $l, k \in \mathbb{N}$ and $\alpha_1, \dots, \alpha_l \in \mathbb{N}_0^n$ satisfy $\sum_{i=1}^l |\alpha_i| = k$. There exists $C > 0$ such that for all $f_1, \dots, f_l \in C_0(\mathbb{R}^n) \cap H^k(\mathbb{R}^n)$ we have*

$$(2.3) \quad \|D^{\alpha_1} f_1 \cdots D^{\alpha_l} f_l\|_{L^2} \leq C \prod_{i=1}^l \|f_i\|_{L^\infty}^{1 - \frac{|\alpha_i|}{k}} \|f_i\|_{H^k}^{\frac{|\alpha_i|}{k}}.$$

In particular,

$$(2.4) \quad \|D^{\alpha_1} f_1 \cdots D^{\alpha_l} f_l\|_{L^2} \leq C \sum_{j=1}^l \prod_{i \neq j} \|f_i\|_{L^\infty} (\|f_j\|_{H^k} + \dots + \|f_l\|_{H^k}).$$

3. EXISTENCE FOR THE PARABOLIC APPROXIMATION

Since $\mathcal{N}(u) = \mathcal{N}(u, u_t, \nabla u, \nabla^2 u, \nabla^3 u)$, energy estimates for the operator $\partial_t^2 + \Delta^2$ are not sufficient to show the existence of a solution of (1.3). Instead, we use the damped plate operator

$$\partial_t^2 + \Delta^2 - \varepsilon \Delta \partial_t,$$

with $\varepsilon \in (0, 1]$ fixed, as a regularization. More precisely, we prove the existence of a solution $u^\varepsilon : \mathbb{R}^n \times [0, T_\varepsilon] \rightarrow N$ of the Cauchy problem

$$(3.1) \quad \begin{cases} \partial_t^2 u^\varepsilon(x, t) + \Delta^2 u^\varepsilon(x, t) - \varepsilon \Delta \partial_t u^\varepsilon(x, t) \perp T_{u^\varepsilon(x, t)} N, & (x, t) \in \mathbb{R}^n \times [0, T_\varepsilon], \\ u^\varepsilon(x, 0) = u_0(x), \quad u_t^\varepsilon(0, x) = u_1(x), & x \in \mathbb{R}^n, \end{cases}$$

where $u_0, u_1 : \mathbb{R}^n \rightarrow \mathbb{R}^L$ satisfy $u_0(x) \in N$ and $u_1(x) \in T_{u_0(x)} N$ for a.e. $x \in \mathbb{R}^n$ as well as

$$(\nabla u_0, u_1) \in H^{k-1}(\mathbb{R}^n) \times H^{k-2}(\mathbb{R}^n)$$

for some $k \in \mathbb{N}$ with $k > \lfloor \frac{n}{2} \rfloor + 2$. In the following we mostly drop the super-/subscript ε and write (u, T) instead of $(u^\varepsilon, T_\varepsilon)$. We note that the condition in (3.1) reads as

$$(3.2) \quad \partial_t^2 u + \Delta^2 u - \varepsilon \Delta \partial_t u = \mathcal{N}(u) - \varepsilon(I - P_u)(\Delta \partial_t u).$$

Using $u(t, x) \in N$, we can expand

$$(3.3) \quad \varepsilon(I - P_u)(\Delta \partial_t u) = \varepsilon d^2 P_u(u_t, \nabla u, \nabla u) + \varepsilon 2dP_u(\nabla u_t, \nabla u) + \varepsilon dP_u(u_t, \Delta u).$$

We thus study the regularized problem

$$(3.4) \quad \begin{aligned} \partial_t^2 u + \Delta^2 u - \varepsilon \Delta \partial_t u &= \mathcal{N}(u) - \varepsilon d^2 P_u(u_t, \nabla u, \nabla u) - \varepsilon 2dP_u(\nabla u_t, \nabla u) - \varepsilon dP_u(u_t, \Delta u) \\ &=: \mathcal{N}_\varepsilon(u). \end{aligned}$$

We next solve (3.4) without the geometric constraint, recalling that only $u(t) - u_0 \in L^2(\mathbb{R}^n)$.

Lemma 3.1. *Let $\varepsilon \in (0, 1)$ and take $u_0, u_1 : \mathbb{R}^n \rightarrow \mathbb{R}^L$ with $u_0(x) \in N$ and $u_1(x) \in T_{u_0(x)} N$ for a.e. $x \in \mathbb{R}^n$ such that*

$$(\nabla u_0, u_1) \in H^{k-1}(\mathbb{R}^n) \times H^{k-2}(\mathbb{R}^n)$$

for some $k \in \mathbb{N}$ with $k > \lfloor \frac{n}{2} \rfloor + 2$. Then (3.4) has a unique local solution $u : \mathbb{R}^n \times [0, T_\varepsilon] \rightarrow \mathbb{R}^L$ satisfying $u(0) = u_0$, $u_t(0) = u_1$, and

$$(3.5) \quad u - u_0 \in C^0([0, T_\varepsilon], H^k(\mathbb{R}^n)) \cap C^1([0, T_\varepsilon], H^{k-2}(\mathbb{R}^n)) \cap H^1(0, T_\varepsilon; H^{k-1}(\mathbb{R}^n)).$$

In addition,

$$(3.6) \quad \nabla u \in L^2(0, T_\varepsilon; H^k(\mathbb{R}^n))$$

and there exists a constant $C < \infty$ such that for $0 \leq t \leq T_\varepsilon$

$$(3.7) \quad \begin{aligned} &\|\nabla^{k-2} u_t(t)\|_{L^2}^2 + \|\nabla^k u(t)\|_{L^2}^2 + \varepsilon \int_0^t \|\nabla^{k-1} u_t(s)\|_{L^2}^2 ds + \varepsilon \int_0^t \|\nabla^{k+1} u(s)\|_{L^2}^2 ds \\ &\leq C \left(\int_0^t \int_{\mathbb{R}^n} \nabla^{k-2}(\mathcal{N}_\varepsilon(u)) \cdot \nabla^{k-2} u_t dx ds + \|\nabla u_0\|_{H^{k-1}}^2 + \|u_1\|_{H^{k-2}}^2 \right). \end{aligned}$$

Before we prove Lemma 3.1, we reduce the problem to functions in L^2 by setting $v(x, t) = u(x, t) - u_0(x)$. We thus rewrite (3.4) as

$$(3.8) \quad \partial_t U + \mathcal{A}_k U = \begin{pmatrix} 0 \\ f_\varepsilon(U) \end{pmatrix}, \quad U(0) = \begin{pmatrix} 0 \\ u_1 \end{pmatrix},$$

where $U = \begin{pmatrix} v \\ v_t \end{pmatrix}$ and $f_\varepsilon(U)$ is defined through

$$(3.9) \quad \begin{aligned} f_\varepsilon(U) &:= \mathcal{N}(v + u_0) - \varepsilon d^2 P_{v+u_0}(v_t, \nabla(v + u_0), \nabla(v + u_0)) \\ &\quad - \varepsilon 2dP_{v+u_0}(\nabla v_t, \nabla(v + u_0)) - \varepsilon dP_{v+u_0}(v_t, \Delta(v + u_0)) - \Delta^2 u_0. \end{aligned}$$

Further the operator $\mathcal{A}_k : H^k(\mathbb{R}^n) \times H^{k-2}(\mathbb{R}^n) \supseteq \mathcal{D}(\mathcal{A}) \rightarrow H^k(\mathbb{R}^n) \times H^{k-2}(\mathbb{R}^n)$ is given by

$$(3.10) \quad \mathcal{A}_k = \begin{pmatrix} 0 & -I \\ \Delta^2 & -\varepsilon\Delta \end{pmatrix}, \quad \mathcal{D}(\mathcal{A}) = H^{k+2}(\mathbb{R}^n) \times H^k(\mathbb{R}^n).$$

Since the operators \mathcal{A}_k extend each other we drop the subscript k . It is well known that $-\mathcal{A}$ generates an analytic C^0 -semigroup $\{T_\varepsilon(t)\}_{t \geq 0}$, see e.g. [1, Prop. 2.3] for the case $k = 2$. Using also standard parabolic theory, see e.g. [6, Prop. 0.1] and [8, Prop. 1.13], we obtain a first linear existence result with some extra regularity.

Lemma 3.2. *Let $r \in \mathbb{N}_0$, $u_1 \in H^{r+1}(\mathbb{R}^n)$, and $g \in C^0([0, T], H^r(\mathbb{R}^n))$. Then there exists a unique solution U of the linear equation*

$$(3.11) \quad \partial_t U + \mathcal{A}U = \begin{pmatrix} 0 \\ g \end{pmatrix}, \quad U(0) = \begin{pmatrix} 0 \\ u_1 \end{pmatrix},$$

satisfying

$$U \in L^2(0, T; H^{r+4} \times H^{r+2}(\mathbb{R}^n)) \cap C^0(0, T; H^{r+3} \times H^{r+1}(\mathbb{R}^n)) \cap H^1(0, T; H^{r+2} \times H^r(\mathbb{R}^n)).$$

We remark that the solution of (3.11) is given by

$$(3.12) \quad U(t) = T_\varepsilon(t) \begin{pmatrix} 0 \\ u_1 \end{pmatrix} + \int_0^t T_\varepsilon(t-s) \begin{pmatrix} 0 \\ g(s) \end{pmatrix} ds.$$

We quantify the above result by the following higher-order energy estimates.

Lemma 3.3. *Let $r \in \mathbb{N}_0$, $g \in C^0([0, T], H^r(\mathbb{R}^n))$, $u_1 \in H^{r+1}(\mathbb{R}^n)$, and $u_0 : \mathbb{R}^n \rightarrow \mathbb{R}^L$ with $\nabla u_0 \in H^{r+3}(\mathbb{R}^n)$. Then v from Lemma 3.2 satisfies*

$$(3.13) \quad \begin{aligned} & \|v_t(t)\|_{H^{r+1}}^2 + \|v(t)\|_{H^{r+3}}^2 + \varepsilon \int_0^t \|\nabla v_t(s)\|_{H^{r+1}}^2 ds + \varepsilon \int_0^t \|\nabla(v+u_0)(s)\|_{H^{r+3}}^2 ds \\ & \leq C(1+T) \left(\frac{1}{\varepsilon} \int_0^t \|g(s) + \Delta^2 u_0\|_{H^r}^2 ds + \|u_1\|_{H^{r+1}}^2 + \|\nabla u_0\|_{H^{r+2}}^2 \right) \end{aligned}$$

for $0 \leq t \leq T$, and

$$(3.14) \quad \begin{aligned} & \|\nabla^{r+1} v_t(t)\|_{L^2}^2 + \|\nabla^{r+3} v(t)\|_{L^2}^2 + \varepsilon \int_0^t \|\nabla^{r+2} v_t(s)\|_{L^2}^2 ds \\ & \leq C \left(- \int_0^t \int_{\mathbb{R}^n} \nabla^r (g(s) + \Delta^2 u_0) \cdot \nabla^r \Delta v_t dx ds + \|u_1\|_{H^{r+1}}^2 + \|\nabla u_0\|_{H^{r+2}}^2 \right). \end{aligned}$$

Proof. Writing $U = (v, v_t)$ in Lemma 3.2, the function $u = v + u_0$ fulfills

$$(3.15) \quad \partial_t^2 u + \Delta^2 u - \varepsilon \Delta \partial_t u = g + \Delta^2 u_0$$

in $L^2(0, T; H^r(\mathbb{R}^n))$. We first differentiate (3.15) of order ∇^l with $l \in \{0, \dots, r\}$. Testing with $-\nabla^l \Delta u_t \in L^2_{t,x}$ and integrating by parts in x , we derive

$$(3.16) \quad \begin{aligned} & \frac{d}{dt} \|\nabla^{l+1} u_t(t)\|_{L^2}^2 + \frac{d}{dt} \|\nabla^{l+3} u(t)\|_{L^2}^2 + \varepsilon \|\nabla^{l+2} u_t(t)\|_{L^2}^2 \\ & \leq \frac{C}{\varepsilon} \|\nabla^l (g + \Delta^2 u_0)\|_{L^2}^2 + \frac{\varepsilon}{2} \|\nabla^{l+2} u_t(t)\|_{L^2}^2, \end{aligned}$$

which makes sense for a.e. t . (Here and below we use the duality (H^1, H^{-1}) in intermediate steps.) We then absorb the last term by the left-hand side and integrate the inequality in t .

To control the second summand with ε in (3.13), we test the differentiated version of (3.15) by $\varepsilon \nabla^l \Delta^2 u$. Here we proceed similarly as before, where we integrate the term

$$\varepsilon \int_0^t \int_{\mathbb{R}^n} \nabla^l \partial_t^2 u \cdot \nabla^l \Delta^2 u dx ds$$

by parts in t and x before absorbing it.

It remains to estimate the L^2 -norm of $v_t(t)$ and the H^2 -norm of $v(t)$. These inequalities follow by testing the equation with u_t and using the fact that

$$\|u - u_0\|_{L_t^\infty L^2} \leq T \|u_t\|_{L_t^\infty L^2}. \quad \square$$

Proof of Lemma 3.1. We aim at constructing a solution $U \in C^0([0, T], H^k \times H^{k-2})$, but due to $\Delta^2 u_0 \in H^{k-4}$ we have $f_\varepsilon(U) \in C^0([0, T], H^{k-4})$, which is insufficient for an application of Lemmas 3.2 and 3.3 in a fixed point argument for v .

We thus approximate u_0 by $u_0^\delta \in C^\infty(\mathbb{R}^n, \mathbb{R}^L)$ for $\delta > 0$ such that $\text{supp}(\nabla u_0^\delta) \subset \mathbb{R}^n$ is compact with

$$(3.17) \quad u_0^\delta \rightarrow u_0 \text{ a.e. and } \nabla u_0^\delta \rightarrow \nabla u_0 \text{ in } H^{k-1}(\mathbb{R}^n) \text{ as } \delta \rightarrow 0^+.$$

Defining $f_{\varepsilon, \delta}$ as above with u_0^δ instead of u_0 , we obtain $f_{\varepsilon, \delta}(U) \in C^0([0, T], H^{k-3}(\mathbb{R}^n))$. For the data (u_0^δ, u_1) we now prove the existence of a fixed point for the operator $v \mapsto \mathcal{S}(v)$ defined through

$$(3.18) \quad \begin{pmatrix} \mathcal{S}(v) \\ \partial_t \mathcal{S}(v) \end{pmatrix} = T_\varepsilon(t) \begin{pmatrix} 0 \\ u_1 \end{pmatrix} + \int_0^t T_\varepsilon(t-s) \begin{pmatrix} 0 \\ f_{\varepsilon, \delta}(v) \end{pmatrix} ds,$$

which acts on the space

$$\begin{aligned} \mathcal{B}_R(T) &:= \{v \in C^0([0, T], H^k) \cap C^1([0, T], H^{k-2}) \mid v(0) = 0, v_t(0) = u_1, \\ &\quad \|v\|_{\mathcal{B}} := \|v_t\|_{L^\infty H^{k-2}} + \|v\|_{L^\infty L^2} + \|\nabla(v + u_0^\delta)\|_{L^\infty H^{k-1}} \leq R\}, \end{aligned}$$

for parameters $R > 0$ and $T \in (0, 1)$ fixed below and the metric given by

$$\|v_1 - v_2\|_{\mathcal{B}(T)} = \|v_1 - v_2\|_{L^\infty H^k} + \|\partial_t v_1 - \partial_t v_2\|_{L^\infty H^{k-2}}, \quad v_1, v_2 \in \mathcal{B}_R(T).$$

Let $\varepsilon \in (0, 1)$ be fixed. We will show that the map

$$\mathcal{S} : \mathcal{B}_R(T) \rightarrow \mathcal{B}_R(T)$$

is strictly contractive with respect to $\|\cdot\|_{\mathcal{B}(T)}$ if we choose $R = R_\delta$ and $T = T_\delta$ with

$$(3.19) \quad \begin{aligned} R_\delta^k &= 3(\|\nabla u_0^\delta\|_{H^{k-1}} + \|u_1\|_{H^{k-2}})^k =: 3R_{0, \delta}^k, \\ T_\delta &= \frac{1}{2} \min \left\{ \left(\frac{\sqrt[k]{3} - 1}{\sqrt[k]{3}} \right)^2 \frac{\varepsilon}{\hat{C}^2(1 + 3R_{0, \delta}^k)^2}, \frac{\varepsilon}{\hat{C}^2(1 + 6R_{0, \delta}^k)^2} \right\} \end{aligned}$$

for a constant \hat{C} depending only on N, n , and k . To show this statement, we have to prove the estimates

$$(3.20) \quad \|\mathcal{S}(v)\|_{\mathcal{B}} \leq \frac{\hat{C}}{\varepsilon^{\frac{1}{2}}} T^{\frac{1}{2}} (1 + \|v\|_{\mathcal{B}}^k) \|v\|_{\mathcal{B}} + \|\nabla u_0^\delta\|_{H^{k-1}} + \|u_1\|_{H^{k-2}},$$

$$(3.21) \quad \|\mathcal{S}(v) - \mathcal{S}(\tilde{v})\|_{\mathcal{B}(T)} \leq \frac{\hat{C}}{\varepsilon^{\frac{1}{2}}} T^{\frac{1}{2}} (1 + \|v\|_{\mathcal{B}}^k + \|\tilde{v}\|_{\mathcal{B}}^k) \|v - \tilde{v}\|_{\mathcal{B}(T)}$$

for $v, \tilde{v} \in \mathcal{B}_R(T)$. To employ the inequality (3.13) for $r = k - 3$, we need to bound the norms

$$\|\mathcal{N}_\varepsilon(v(t) + u_0^\delta)\|_{H^{k-3}}^2 \quad \text{and} \quad \|\mathcal{N}_\varepsilon(v(t) + u_0^\delta) - \mathcal{N}_\varepsilon(\tilde{v}(t) + u_0^\delta)\|_{H^{k-3}}^2$$

by $C(1 + \|v\|_{\mathcal{B}}^k) \|v\|_{\mathcal{B}}$ and $C(1 + \|v\|_{\mathcal{B}}^k + \|\tilde{v}\|_{\mathcal{B}}^k) \|v - \tilde{v}\|_{\mathcal{B}(T)}$, respectively. This is done by means of Lemma A.1 and Corollary A.4 combined with a careful application of the Moser estimate in Lemma 2.2. We give the relevant details below in Section 4 in the proof of the a priori estimate and in Section 6 for the uniqueness since these parts require more thought. In this way we obtain in the fixed point $v^\delta = \mathcal{S}(v^\delta)$ satisfying

$$(3.22) \quad \|v_t^\delta\|_{L^\infty H^{k-2}}^2 + \|v^\delta\|_{L^\infty H^k}^2 + \varepsilon \int_0^{T_\delta} \|v_t^\delta(s)\|_{H^{k-1}}^2 ds + \varepsilon \int_0^{T_\delta} \|\nabla(v^\delta + u_0^\delta)\|_{H^k}^2 ds \lesssim R_\delta^2.$$

In particular, $v^\delta \in L^2(0, T_\delta; H^{k+1}) \cap H^1(0, T_\delta; H^{k-1})$.

We next define R_0 , R and $\tilde{T} > 0$ in the same way as $R_{0,\delta}$, R_δ and T_δ using u_0 instead of u_0^δ and the R_0 instead of R_0^δ . Thus,

$$R_{0,\delta} \rightarrow R_0, \quad R_\delta \rightarrow R, \quad T_\delta \rightarrow \tilde{T} \quad \text{as } \delta \rightarrow 0^+.$$

For sufficiently small $\delta > 0$ we have $T_\delta > \frac{1}{2}\tilde{T} =: T$ and $|R_{0,\delta} - R_0| \leq R_0$. Hence $v^\delta : \mathbb{R}^n \times [0, T] \rightarrow \mathbb{R}^L$ is well defined and $\|v^\delta\|_{\mathcal{B}(T)} \leq CR$ for a constant $C > 0$. Observe that for sufficiently small $\delta, \delta' > 0$, the differences $v^\delta - v^{\delta'}$ and $\partial_t v^\delta - \partial_t v^{\delta'}$ solve (3.11) with the nonlinearity

$$\mathcal{N}_\varepsilon(v^\delta + u_0^\delta) - \mathcal{N}_\varepsilon(v^{\delta'} + u_0^{\delta'}) + \Delta^2(u_0^\delta - u_0^{\delta'}) \in C^0([0, T], H^{k-3}).$$

Similar to the proof of the Lipschitz estimate (3.21), Lemma 3.3 then yields the bound

$$\begin{aligned} & \|v^\delta - v^{\delta'}\|_{\mathcal{B}(T)}^2 + \varepsilon \int_0^{T_\delta} \|v_t^\delta(s) - v_t^{\delta'}(s)\|_{H^{k-1}}^2 ds + \varepsilon \int_0^{T_\delta} \|\nabla(v^\delta - v^{\delta'}) + \nabla(u_0^\delta - u_0^{\delta'})\|_{H^k}^2 ds \\ & \leq C \frac{T}{\varepsilon} (1 + R^{2k}) \|v^\delta - v^{\delta'}\|_{\mathcal{B}(T)}^2 + \tilde{C}_{\varepsilon, R} \|\nabla u_0^\delta - \nabla u_0^{\delta'}\|_{H^{k-1}}^2. \end{aligned}$$

Hence, if $T = T(\varepsilon)$ is sufficiently small, as $\delta \rightarrow 0$ the functions v^δ tend to a function

$$v \in C^0([0, T], H^k) \cap C^1([0, T], H^{k-2}) \cap H^1(0, T; H^{k-1})$$

with $\nabla(v + u_0) \in L^2(0, T; H^k)$, where the limits exist in these spaces. In particular, (v, v_t) is a solution of (3.8) and $u = v + u_0$ solves (3.4). Moreover, by (3.13) the function $u^\delta = v^\delta + u_0^\delta$ satisfies inequality (3.7), and therefore this estimate also holds for u since $u_t^\delta \rightarrow u_t$ strongly in $C^0([0, T], H^{k-2})$ and $\mathcal{N}_\varepsilon(u^\delta) \rightarrow \mathcal{N}_\varepsilon(u)$ strongly in $L^2(0, T; H^{k-2})$ because of Corollary A.4 and Lemma 2.2.

For the uniqueness of v , we note that, for a second solution \tilde{v} , the functions $w = v - \tilde{v}$ and $w_t = v_t - \tilde{v}_t$ solve (3.11) with the nonlinearity $\mathcal{N}_\varepsilon(v + u_0) - \mathcal{N}_\varepsilon(\tilde{v} + u_0) \in C^0([0, T], H^{k-3})$. Lemma 3.3 then yields the estimate

$$(3.23) \quad \|v - \tilde{v}\|_{\mathcal{B}(T)}^2 \leq C \frac{T}{\varepsilon} (1 + R^{2k}) \|v - \tilde{v}\|_{\mathcal{B}(T)}^2.$$

(Note that u_0 from the Lemma is different, namely $u_0 = 0$.) Hence, if T is sufficiently small, we obtain $v = \tilde{v}$ and thus $u = v + u_0$ is unique. \square

We next show that the above solution actually takes values in the target manifold.

Proposition 3.4. *Let $\varepsilon \in (0, 1)$ and take $u_0, u_1 : \mathbb{R}^n \rightarrow \mathbb{R}^L$ with $u_0(x) \in N$ and $u_1(x) \in T_{u_0(x)}N$ for a.e. $x \in \mathbb{R}^n$ satisfying*

$$(\nabla u_0, u_1) \in H^{k-1}(\mathbb{R}^n) \times H^{k-2}(\mathbb{R}^n)$$

for some $k \in \mathbb{N}$ with $k > \lfloor \frac{n}{2} \rfloor + 2$. Then there exists a maximal existence time $T_{\varepsilon, m} \in (0, \infty]$ and a unique solution $u \in \mathbb{R}^n \times [0, T_{\varepsilon, m}] \rightarrow N$ of (3.1) with $u(0) = u_0$, $\partial_t u(0) = u_1$,

$$u - u_0 \in C^0([0, T_{\varepsilon, m}], H^k) \cap C^1([0, T_{\varepsilon, m}], H^{k-2}) \cap H_{loc}^1([0, T_{\varepsilon, m}], H^{k-1}(\mathbb{R}^n))$$

and $\nabla u \in L_{loc}^2([0, T_{\varepsilon, m}], H^k(\mathbb{R}^n))$ which satisfies (3.7) for $t \in [0, T_{\varepsilon, m}]$.

Proof. Fix $\varepsilon \in (0, 1)$. Let $u : \mathbb{R}^n \times [0, T] \rightarrow \mathbb{R}^L$ be the solution of (3.1) constructed in Lemma 3.1. We first show that $u(x, t) \in N$ for $x \in \mathbb{R}^n$ and $t > 0$ small enough. Since

$$C^0([0, T], H^k) \hookrightarrow C^0(\mathbb{R}^n \times [0, T])$$

and $u_0 \in N$ a.e. on \mathbb{R}^n , there exists a time $\tilde{T} \in (0, T]$ such that for $t \in [0, \tilde{T}]$ the distance

$$\|\text{dist}(u(t), N)\|_{L^\infty} \leq \sup_{x \in \mathbb{R}^n} |u(x, t) - u_0(x)| \lesssim \|u(t) - u_0\|_{H^k}$$

is so small that $\bar{u} = \pi(u)$ is well-defined. We then let $w = \bar{u} - u$ and we note that $w(0) = \partial_t w(0) = 0$. Calculating

$$\partial_t^2 \bar{u} = d\pi_u \partial_t^2 u + d^2 \pi_u(u_t, u_t),$$

$$\begin{aligned}
\Delta \bar{u}_t &= d\pi_u \Delta u_t + d^2 \pi_u (\Delta u, u_t) + 2d^2 \pi_u (\nabla u_t, \nabla u) + d^3 \pi_u (\nabla u, \nabla u, u_t), \\
\Delta^2 \bar{u} &= d\pi_u \Delta^2 u + d^2 \pi_u (\Delta u, \Delta u) + 4d^2 \pi_u (\nabla u, \nabla \Delta u) + 2d^2 \pi_u (\nabla^2 u, \nabla^2 u) \\
&\quad + 2d^3 \pi_u (\nabla u, \nabla u, \Delta u) + 4d^3 \pi_u (\nabla u, \nabla u, \nabla^2 u) \\
&\quad + d^4 \pi_u (\nabla u, \nabla u, \nabla u, \nabla u),
\end{aligned}$$

we conclude that

$$\begin{aligned}
(\partial_t^2 + \Delta^2 - \varepsilon \Delta \partial_t)w &= d\pi_u \left((\partial_t^2 + \Delta^2 - \varepsilon \Delta \partial_t)u \right) + \mathcal{N}_\varepsilon(u) - \mathcal{N}_\varepsilon(u) \\
&= d\pi_u(\mathcal{N}_\varepsilon(u)) \in T_{\bar{u}}N.
\end{aligned}$$

Next, we note that

$$w_t = \left((\pi - I)(u) \right)_t = (d\pi_{\bar{u}} - I)u_t \perp T_{\bar{u}}N.$$

By testing the above equation for w by w_t , it follows

$$\partial_t \frac{1}{2} \int_{\mathbb{R}^n} |w_t|^2 dx + \partial_t \frac{1}{2} \int_{\mathbb{R}^n} |\Delta w|^2 dx + \varepsilon \int_{\mathbb{R}^n} |\nabla w_t|^2 dx = 0.$$

This fact implies that $w_t = 0$ and hence $w = 0$, which means that $u \in N$.

The claimed uniqueness follows similarly to the end of the proof of Lemma 3.1. Finally, we let $T_{\varepsilon, m} \geq \tilde{T}$ be the supremum of times $T' > 0$ such that we have a solution $u : [0, T'] \times \mathbb{R}^n \rightarrow N$ of (3.1) with $u(0) = u_0$, $\partial_t u(0) = u_1$,

$$u - u_0 \in C^0([0, T'], H^k) \cap C^1([0, T'], H^{k-2}) \cap H^1(0, T'; H^{k-1}(\mathbb{R}^n))$$

and $\nabla u \in L^2(0, T'; H^k(\mathbb{R}^n))$ which satisfies (3.7) on $[0, T']$. \square

Remark 3.5. We remark that up to now we fixed $\varepsilon \in (0, 1)$. Since the constants in the upper bound in estimates such as (3.22) are of order $O(\varepsilon^{-1})$, we have to prove ε independent estimates in the next section.

4. THE A PRIORI ESTIMATE

We now prove an a priori estimate for the solution $u^\varepsilon : \mathbb{R}^n \times [0, T_{\varepsilon, m}] \rightarrow N$ of the equation

$$(4.1) \quad \partial_t^2 u^\varepsilon + \Delta^2 u^\varepsilon - \varepsilon \Delta \partial_t u^\varepsilon \perp T_{u^\varepsilon} N \quad \text{on } \mathbb{R}^n \times [0, T_{\varepsilon, m}]$$

given by Proposition 3.4 with $\varepsilon \in (0, 1)$ and initial data $u_0, u_1 : \mathbb{R}^n \rightarrow \mathbb{R}^L$ such that $u_0(x) \in N$ and $u_1(x) \in T_{u_0(x)}N$ for a.e. $x \in \mathbb{R}^n$ as well as

$$(\nabla u_0, u_1) \in H^{k-1}(\mathbb{R}^n) \times H^{k-2}(\mathbb{R}^n)$$

for some $k \in \mathbb{N}$ with $k > \lfloor \frac{n}{2} \rfloor + 2$. As before we write u instead of u^ε , and we fix a number $T < T_{\varepsilon, m}$. Moreover, (3.7) says that

$$\begin{aligned}
&\|\nabla^{k-2} u_t(t)\|_{L^2}^2 + \|\nabla^k u(t)\|_{L^2}^2 + \varepsilon \int_0^t \|\nabla^{k-1} u_t(s)\|_{L^2}^2 ds \\
(4.2) \quad &\lesssim \int_0^t \int_{\mathbb{R}^n} \nabla^{k-2} [\mathcal{N}(u) - \varepsilon(I - P_u)(\Delta u_t)] \cdot \nabla^{k-2} u_t dx ds + \|\nabla^{k-2} u_1\|_{L^2}^2 + \|\nabla^k u_0\|_{L^2}^2
\end{aligned}$$

for $t \in [0, T]$. We recall that the summand with ε on the right-hand side is well defined because of (3.3).

In the following, we often make use of the relations $\mathcal{N}(u) \perp T_u N$ and $u_t \in T_u N$ which hold since $u(x, t) \in N$ for a.e. $(x, t) \in \mathbb{R}^n \times [0, T]$. In particular, $\mathcal{N}(u) = (I - P_u)\mathcal{N}(u)$. Using this fact, we first write

$$\begin{aligned}
(4.3) \quad \nabla^{k-2}(\mathcal{N}(u))\nabla^{k-2}u_t &= \sum_{\substack{m_1+m_2=k-2 \\ m_1>0}} \nabla^{m_1}(I - P_u) \star \nabla^{m_2}(\mathcal{N}(u))\nabla^{k-2}u_t \\
&\quad + \nabla^{k-2}(\mathcal{N}(u))(I - P_u)\nabla^{k-2}u_t
\end{aligned}$$

$$\begin{aligned}
&= \sum_{\substack{m_1+m_2=k-2 \\ m_1>0}} \nabla^{m_1}(I - P_u) \star \nabla^{m_2}(\mathcal{N}(u)) \nabla^{k-2} u_t \\
&\quad - \sum_{\substack{l_1+l_2=k-2 \\ l_1>0}} \nabla^{k-2}(\mathcal{N}(u)) \star \nabla^{l_1}[(I - P_u)] \nabla^{l_2} u_t \\
&=: I_1 + I_2,
\end{aligned}$$

where the second equality follows from the Leibniz formula

$$(4.4) \quad 0 = \nabla^{k-2} [(I - P_u)u_t] = \sum_{\substack{l_1+l_2=k-2 \\ l_1>0}} \nabla^{l_1}[(I - P_u)] \star \nabla^{l_2} u_t + (I - P_u) \nabla^{k-2} u_t.$$

In (4.2) we thus split

$$\begin{aligned}
&\int_{\mathbb{R}^n} \nabla^{k-2}(\mathcal{N}(u) - \varepsilon(I - P_u)(\Delta u_t)) \cdot \nabla^{k-2} u_t \, dx \\
&= \int_{\mathbb{R}^n} \nabla^{k-2}(\mathcal{N}(u)) \cdot \nabla^{k-2} u_t \, dx - \varepsilon \int_{\mathbb{R}^n} \nabla^{k-2}((I - P_u)(\Delta u_t)) \cdot \nabla^{k-2} u_t \, dx \\
(4.5) \quad &= \int_{\mathbb{R}^n} I_1 \, dx + \int_{\mathbb{R}^n} I_2 \, dx - \varepsilon \int_{\mathbb{R}^n} \nabla^{k-2}((I - P_u)(\Delta u_t)) \cdot \nabla^{k-2} u_t \, dx,
\end{aligned}$$

We start by estimating

$$\int_{\mathbb{R}^n} I_1 \, dx \leq \sum_{\substack{m_1+m_2=k-2 \\ m_1>0}} \|\nabla^{m_1}(I - P_u) \star \nabla^{m_2}(\mathcal{N}(u))\|_{L^2} \|\nabla^{k-2} u_t\|_{L^2}.$$

Lemma 2.1 yields the identity

$$(4.6) \quad \nabla^{m_1}(I - P_u) = - \sum_{j=1}^{m_1} \sum_{\sum_{i=1}^j \tilde{k}_i = m_1 - j} d^j P_u(\nabla^{\tilde{k}_1+1} u \star \dots \star \nabla^{\tilde{k}_j+1} u),$$

which implies the pointwise inequality

$$(4.7) \quad |\nabla^{m_1}(I - P_u)| \lesssim \sum_{j=1}^{m_1} \sum_{\sum_{i=1}^j \tilde{k}_i = m_1 - j} |\nabla^{\tilde{k}_1+1} u| \dots |\nabla^{\tilde{k}_j+1} u|.$$

On the other hand, Lemma A.1 allows us to bound $|\nabla^{m_2}(\mathcal{N}(u))|$ pointwise (up to a constant) by terms of the form

$$(4.8) \quad |\nabla^{\tilde{m}_1+1} u| \dots |\nabla^{\tilde{m}_i+1} u| [|\nabla^{k_1} u_t| |\nabla^{k_2} u_t| + |\nabla^{k_1+2} u| |\nabla^{k_2+2} u| + |\nabla^{k_1+3} u| |\nabla^{k_2+1} u|],$$

$$(4.9) \quad |\nabla^{\tilde{m}_1+1} u| \dots |\nabla^{\tilde{m}_i+1} u| [|\nabla^{k_1+1} u| |\nabla^{k_2+1} u| |\nabla^{k_3+2} u|],$$

$$(4.10) \quad |\nabla^{\tilde{m}_1+1} u| \dots |\nabla^{\tilde{m}_i+1} u| [|\nabla^{k_1+1} u| |\nabla^{k_2+1} u| |\nabla^{k_3+1} u| |\nabla^{k_4+1} u|],$$

where $i = 1, \dots, m_2$ and $\tilde{m}_1 + \dots + \tilde{m}_i + k_1 + \dots = m_2 - i$ are as in Lemma A.1. Moreover, in the case $i = 0$ (where no derivatives fall on the coefficients) the terms are of the form

$$\begin{aligned}
&|\nabla^{k_1} u_t| |\nabla^{k_2} u_t| + |\nabla^{k_1+2} u| |\nabla^{k_2+2} u| + |\nabla^{k_1+3} u| |\nabla^{k_2+1} u|, \\
&|\nabla^{k_1+1} u| |\nabla^{k_2+1} u| |\nabla^{k_3+2} u|, \\
&|\nabla^{k_1+1} u| |\nabla^{k_2+1} u| |\nabla^{k_3+1} u| |\nabla^{k_4+1} u|,
\end{aligned}$$

where $k_j \in \mathbb{N}_0$ and $k_1 + k_2 + \dots = m_2$. Note that $m_2 \leq k - 3$ since $m_1 > 0$. In the following we use the notation (4.8) - (4.10) for all five cases, setting $i = 0$ for the latter three.

Combining the above considerations with Lemma 2.2, we can now estimate the norm

$$\|\nabla^{m_1}(I - P_u) \nabla^{m_2}(\mathcal{N}(u))\|_{L^2},$$

where we distinguish five cases according to the terms in the brackets in (4.8) - (4.10).

Case 1: $\nabla^{k_1} u_t \star \nabla^{k_2} u_t$

We use Lemma 2.2 with

$$f_1 = \nabla u, \dots, f_j = \nabla u, f_{j+1} = \nabla u, \dots, f_{j+i} = \nabla u, f_{j+i+1} = u_t, f_{j+i+2} = u_t,$$

and derivatives of order

$$\tilde{k}_1 + \dots + \tilde{k}_j + \tilde{m}_1 + \dots + \tilde{m}_i + k_1 + k_2 = m_1 + m_2 - i - j = k - 2 - (i + j).$$

Employing also Young's inequality, it follows

$$\begin{aligned} & \left\| |\nabla^{\tilde{k}_1+1} u| \dots |\nabla^{\tilde{k}_j+1} u| |\nabla^{\tilde{m}_1+1} u| \dots |\nabla^{\tilde{m}_i+1} u| |\nabla^{k_1} u_t| |\nabla^{k_2} u_t| \right\|_{L^2} \\ & \lesssim \left((1 + \|\nabla u\|_{L^\infty}^{k-3}) \|u_t\|_{L^\infty}^2 + (1 + \|\nabla u\|_{L^\infty}^{k-2}) \|u_t\|_{L^\infty} \right) (\|\nabla u\|_{H^{k-2-i-j}} + \|u_t\|_{H^{k-2-i-j}}) \\ & \lesssim (1 + \|\nabla u\|_{L^\infty}^{k-1} + \|u_t\|_{L^\infty}^{k-1}) (\|\nabla u\|_{H^{k-1}} + \|u_t\|_{H^{k-2}}). \end{aligned}$$

The other cases will be treated similarly.

Case 2: $\nabla^{k_1+2} u \star \nabla^{k_2+2} u$

Here it is exploited that $m_1 > 0$ in I_1 due to the cancellation from (4.4). This time Lemma 2.2 is applied with $f_1 = \dots = f_{j+i+2} = \nabla u$ and derivatives of order

$$\tilde{k}_1 + \dots + \tilde{k}_j + \tilde{m}_1 + \dots + \tilde{m}_i + k_1 + k_2 + 2 = m_1 + m_2 + 2 - i - j = k - (i + j) \leq k - 1$$

since $j > 0$ by (4.6). We estimate

$$\begin{aligned} & \left\| |\nabla^{\tilde{k}_1+1} u| \dots |\nabla^{\tilde{k}_j+1} u| |\nabla^{\tilde{m}_1+1} u| \dots |\nabla^{\tilde{m}_i+1} u| |\nabla^{k_1+2} u| |\nabla^{k_2+2} u| \right\|_{L^2} \\ & \lesssim \sum_{i,j} \|\nabla u\|_{L^\infty}^{i+j+1} \|\nabla u\|_{H^{k-i-j}} \lesssim (1 + \|\nabla u\|_{L^\infty}^{k-1}) \|\nabla u\|_{H^{k-1}}. \end{aligned}$$

Case 3: $\nabla^{k_1+3} u \star \nabla^{k_2+1} u$

As in the previous case, $C(1 + \|\nabla u\|_{L^\infty}^{k-1}) \|\nabla u\|_{H^{k-1}}$ dominates

$$\left\| |\nabla^{\tilde{k}_1+1} u| \dots |\nabla^{\tilde{k}_j+1} u| |\nabla^{\tilde{m}_1+1} u| \dots |\nabla^{\tilde{m}_i+1} u| |\nabla^{k_1+3} u| |\nabla^{k_2+1} u| \right\|_{L^2}.$$

Case 4: $\nabla^{k_1+1} u \star \nabla^{k_2+1} u \star \nabla^{k_3+2} u$

We apply Lemma 2.2 to the functions $f_1 = \dots = f_{j+i+3} = \nabla u$ with derivatives of order $\tilde{k}_1 + \dots + \tilde{k}_j + \tilde{m}_1 + \dots + \tilde{m}_i + k_1 + k_2 + k_3 + 1 = m_1 + m_2 + 1 - i - j = k - 1 - (i + j)$, leading to the bound

$$\begin{aligned} & \left\| |\nabla^{\tilde{k}_1+1} u| \dots |\nabla^{\tilde{k}_j+1} u| |\nabla^{\tilde{m}_1+1} u| \dots |\nabla^{\tilde{m}_i+1} u| |\nabla^{k_1+1} u| |\nabla^{k_2+1} u| |\nabla^{k_3+2} u| \right\|_{L^2} \\ & \lesssim \sum_{i,j} \|\nabla u\|_{L^\infty}^{i+j+2} \|\nabla u\|_{H^{k-2-i-j}} \lesssim (1 + \|\nabla u\|_{L^\infty}^k) \|\nabla u\|_{H^{k-1}}. \end{aligned}$$

Case 5: $\nabla^{k_1+1} u \star \nabla^{k_2+1} u \star \nabla^{k_3+1} u \star \nabla^{k_4+1} u$

We now use Lemma 2.2 with $f_1 = \dots = f_{j+i+4} = \nabla u$ and derivatives of order

$$\tilde{k}_1 + \dots + \tilde{k}_j + \tilde{m}_1 + \dots + \tilde{m}_i + k_1 + k_2 + k_3 + k_4 = m_1 + m_2 - i - j = k - 2 - (i + j).$$

Hence, we have

$$\begin{aligned} & \left\| |\nabla^{\tilde{k}_1+1} u| \dots |\nabla^{\tilde{k}_j+1} u| |\nabla^{\tilde{m}_1+1} u| \dots |\nabla^{\tilde{m}_i+1} u| |\nabla^{k_1+1} u| |\nabla^{k_2+1} u| |\nabla^{k_3+1} u| |\nabla^{k_4+1} u| \right\|_{L^2} \\ & \lesssim \sum_{i,j} \|\nabla u\|_{L^\infty}^{i+j+3} \|\nabla u\|_{H^{k-2-i-j}} \lesssim (1 + \|\nabla u\|_{L^\infty}^{k+1}) \|\nabla u\|_{H^{k-1}}. \end{aligned}$$

Summing up the five cases, we infer

$$(4.11) \quad \|I_1\|_{L^1} \lesssim (1 + \|\nabla u\|_{L^\infty}^{k-1} + \|u_t\|_{L^\infty}^{k-1}) (\|\nabla u\|_{H^{k-1}} + \|u_t\|_{H^{k-2}}).$$

Next, in I_2 from (4.5) we integrate by parts in order to conclude

$$\int_{\mathbb{R}^n} I_2 dx = \sum_{\substack{l_1+l_2=k-2 \\ l_1>0}} \int_{\mathbb{R}^n} \nabla^{k-3} (\mathcal{N}(u)) \star [\nabla^{l_1+1} (I - P_u) \nabla^{l_2} u_t] dx$$

$$\begin{aligned}
& + \sum_{\substack{l_1+l_2=k-2 \\ l_1>0}} \int_{\mathbb{R}^n} \nabla^{k-3}(\mathcal{N}(u)) \star [\nabla^{l_1}(I - P_u)\nabla^{l_2+1}u_t] dx \\
& =: I_2^1 + I_2^2.
\end{aligned}$$

These terms are estimated by

$$(4.12) \quad |I_2^1| \lesssim \sum_{\substack{l_1+l_2=k-2 \\ l_1>0}} \|\nabla^{k-3}(\mathcal{N}(u))\|_{L^2} \|\nabla^{l_1+1}(I - P_u)\nabla^{l_2+1}u_t\|_{L^2},$$

$$(4.13) \quad |I_2^2| \lesssim \sum_{\substack{l_1+l_2=k-2 \\ l_1>0}} \|\nabla^{k-3}(\mathcal{N}(u))\|_{L^2} \|\nabla^{l_1}(I - P_u)\nabla^{l_2+1}u_t\|_{L^2}.$$

We control $\|\nabla^{k-3}(\mathcal{N}(u))\|_{L^2}$ by terms of the form (4.8) - (4.10) in the L^2 norm, obtaining as above

$$\|\nabla^{k-3}(\mathcal{N}(u))\|_{L^2} \lesssim (1 + \|\nabla u\|_{L^\infty}^k + \|u_t\|_{L^\infty}^{k-2})(\|\nabla u\|_{H^{k-1}} + \|u_t\|_{H^{k-2}}).$$

Equation (4.6) and Lemma 2.2 further imply

$$\begin{aligned}
\|\nabla^{l_1+1}(I - P_u)\nabla^{l_2+1}u_t\|_{L^2} & \lesssim \sum_{j=1}^{l_1+1} \sum_{\sum_{i=1}^j \tilde{m}_i = l_1+1-j} \|\nabla^{\tilde{m}_1+1}u \dots \nabla^{\tilde{m}_j+1}u\|_{L^2} \|\nabla^{l_2+1}u_t\|_{L^2} \\
& \lesssim (1 + \|\nabla u\|_{L^\infty}^{k-1} + \|u_t\|_{L^\infty}^{k-1})(\|\nabla u\|_{H^{k-1}} + \|u_t\|_{H^{k-2}})
\end{aligned}$$

where $\tilde{m}_1 + \dots + \tilde{m}_j + l_2 = k - 1 - i \leq k - 2$. Similarly, we have

$$\begin{aligned}
\|\nabla^{l_1}(I - P_u)\nabla^{l_2+1}u_t\|_{L^2} & \lesssim \sum_{j=1}^{l_1} \sum_{\sum_{i=1}^j \tilde{m}_i = l_1-j} \|\nabla^{\tilde{m}_1+1}u \dots \nabla^{\tilde{m}_j+1}u\|_{L^2} \|\nabla^{l_2+1}u_t\|_{L^2} \\
& \lesssim (1 + \|\nabla u\|_{L^\infty}^{k-2} + \|u_t\|_{L^\infty}^{k-2})(\|\nabla u\|_{H^{k-1}} + \|u_t\|_{H^{k-2}})
\end{aligned}$$

by Lemma 2.2 with $\tilde{m}_1 + \dots + \tilde{m}_j + l_2 + 1 = k - 1 - i \leq k - 2$, since $l_1 > 0$. The above three inequalities yield

$$(4.14) \quad \|I_2\|_{L^1} \lesssim (1 + \|\nabla u\|_{L^\infty}^{2k-1} + \|u_t\|_{L^\infty}^{2k-1})(\|\nabla u\|_{H^{k-1}}^2 + \|u_t\|_{H^{k-2}}^2).$$

Finally, for the regularization term, we observe

$$\begin{aligned}
-\varepsilon \int_{\mathbb{R}^n} \nabla^{k-2}[(I - P_u)(\Delta u_t)] \nabla^{k-2}u_t dx & = \varepsilon \int_{\mathbb{R}^n} \nabla^{k-3}[(I - P_u)(\Delta u_t)] \nabla^{k-1}u_t dx \\
& \leq C \|\nabla^{k-3}[(I - P_u)(\Delta u_t)]\|_{L^2}^2 + \frac{\varepsilon}{2} \|\nabla^{k-1}u_t\|_{L^2}^2.
\end{aligned}$$

In view of (3.3), to bound $\|\nabla^{k-3}[(I - P_u)(\Delta u_t)]\|_{L^2}^2$ it suffices to estimate

$$(4.15) \quad \|\nabla^{\tilde{m}_1+1}u \dots \nabla^{\tilde{m}_i+1}u [|\nabla^{k_1+1}u_t| |\nabla^{k_2+1}u| + |\nabla^{k_1}u_t| |\nabla^{k_2+2}u|]\|_{L^2}^2,$$

$$(4.16) \quad \|\nabla^{\tilde{m}_1+1}u \dots \nabla^{\tilde{m}_i+1}u [|\nabla^{k_1}u_t| |\nabla^{k_2+1}u| |\nabla^{k_3+1}u|]\|_{L^2}^2,$$

where $\tilde{m}_1 + \dots + \tilde{m}_i + k_1 + k_2 + 1 = k - 2 - i$ and $\tilde{m}_1 + \dots + \tilde{m}_i + k_1 + k_2 + k_3 = k - 3 - i$, respectively. As before, Lemma 2.2 implies the inequalities

$$(4.17) \quad \begin{aligned} & \|\nabla^{\tilde{m}_1+1}u \dots \nabla^{\tilde{m}_i+1}u [|\nabla^{k_1+1}u_t| |\nabla^{k_2+1}u| + |\nabla^{k_1}u_t| |\nabla^{k_2+2}u|]\|_{L^2}^2 \\ & \lesssim (1 + \|\nabla u\|_{L^\infty}^{2(k-2)} + \|u_t\|_{L^\infty}^{2(k-2)})(\|u_t\|_{H^{k-2}}^2 + \|\nabla u\|_{H^{k-2}}^2), \end{aligned}$$

$$(4.18) \quad \begin{aligned} & \|\nabla^{\tilde{m}_1+1}u \dots \nabla^{\tilde{m}_i+1}u [|\nabla^{k_1}u_t| |\nabla^{k_2+1}u| |\nabla^{k_3+1}u|]\|_{L^2}^2 \\ & \lesssim (1 + \|\nabla u\|_{L^\infty}^{2(k-1)} + \|u_t\|_{L^\infty}^{2(k-1)})(\|u_t\|_{H^{k-2}}^2 + \|\nabla u\|_{H^{k-2}}^2). \end{aligned}$$

Putting together (4.11), (4.14), (4.17) and (4.18), we arrive at the inequality

$$\left| \int_{\mathbb{R}^n} \nabla^{k-2}(\mathcal{N}(u) - \varepsilon(I - P_u)(\Delta u_t)) \cdot \nabla^{k-2}u_t dx \right|$$

$$\lesssim (1 + \|\nabla u\|_{L^\infty}^{2k} + \|u_t\|_{L^\infty}^{2k})(\|\nabla u\|_{H^{k-1}}^2 + \|u_t\|_{H^{k-2}}^2) + \frac{\varepsilon}{2} \|\nabla^{k-1} u_t\|_{L^2}^2.$$

Subtracting the last term on both sides of (4.2), for $t \in [0, T]$ we conclude

$$(4.19) \quad \begin{aligned} & \|\nabla^{k-2} u_t(t)\|_{L^2}^2 + \|\nabla^k u(t)\|_{L^2}^2 + \frac{\varepsilon}{2} \int_0^t \|\nabla^{k-1} u_t(s)\|_{L^2}^2 ds \\ & \lesssim \int_0^t \left[(1 + \|\nabla u\|_{L^\infty}^{2k} + \|u_t\|_{L^\infty}^{2k})(\|\nabla u\|_{H^{k-1}}^2 + \|u_t\|_{H^{k-2}}^2) \right] ds + \|\nabla^{k-2} u_1\|_{L^2}^2 + \|\nabla^k u_0\|_{L^2}^2. \end{aligned}$$

It remains to bound the lower order terms. Testing (4.1) by $u_t \in T_u N$, we infer

$$(4.20) \quad \|u_t(t)\|_{L^2}^2 + \|\Delta u(t)\|_{L^2}^2 + \varepsilon \int_0^t \|\nabla u_t(s)\|_{L^2}^2 ds = \|u_1\|_{L^2}^2 + \|\Delta u_0\|_{L^2}^2.$$

Since also

$$\frac{d}{dt} \int_{\mathbb{R}^n} |\nabla u|^2 dx \leq \int_{\mathbb{R}^n} |u_t|^2 dx + \int_{\mathbb{R}^n} |\Delta u|^2 dx,$$

it follows

$$(4.21) \quad \begin{aligned} \|\nabla u(t)\|_{L^2}^2 & \leq \|\nabla u_0\|_{L^2}^2 + \int_0^t \|\Delta u(s)\|_{L^2}^2 + \|u_t(s)\|_{L^2}^2 ds \\ & = \|\nabla u_0\|_{L^2}^2 + t(\|u_1\|_{L^2}^2 + \|\Delta u_0\|_{L^2}^2) \end{aligned}$$

for $t \in [0, T]$. The other derivatives are treated via interpolation, more precisely

$$\begin{aligned} \|\nabla^l u_t\|_{L^2}^2 & \lesssim \|\nabla^{k-1} u_t\|_{L^2}^{\frac{2(l-1)}{k-2}} \|\nabla u_t\|_{L^2}^{\frac{2(k-1-l)}{k-2}}, \quad l = 2, \dots, k-2, \\ \|\nabla^l u_t\|_{L^2}^2 & \lesssim \|\nabla^{k-2} u_t\|_{L^2}^{\frac{2l}{k-2}} \|u_t\|_{L^2}^{\frac{2(k-2-l)}{k-2}}, \quad l = 1, \dots, k-3, \\ \|\nabla^l u\|_{L^2}^2 & \lesssim \|\nabla^k u\|_{L^2}^{\frac{2(l-2)}{k-2}} \|\Delta u\|_{L^2}^{\frac{2(k-l)}{k-2}}, \quad l = 3, \dots, k-1. \end{aligned}$$

Estimate (4.19) and the above inequalities lead to the core estimate

$$(4.22) \quad \begin{aligned} & \|u_t(t)\|_{H^{k-2}}^2 + \|\nabla u(t)\|_{H^{k-1}}^2 + \frac{\varepsilon}{2} \int_0^t \|\nabla u_t(s)\|_{H^{k-2}}^2 ds \\ & \lesssim \int_0^t \left[(1 + \|\nabla u\|_{L^\infty}^{2k} + \|u_t\|_{L^\infty}^{2k})(\|\nabla u\|_{H^{k-1}}^2 + \|u_t\|_{H^{k-2}}^2) \right] ds \\ & \quad + (1+T)(\|u_1\|_{H^{k-2}}^2 + \|\nabla u_0\|_{H^{k-1}}^2), \quad t \in [0, T]. \end{aligned}$$

for solutions of (3.1) and $T < T_{\varepsilon, m}$. Using Gronwall's lemma we also obtain

$$(4.23) \quad \begin{aligned} & \sup_{t \in [0, T]} \left(\|u_t(t)\|_{H^{k-2}}^2 + \|\nabla u(t)\|_{H^{k-1}}^2 \right) \\ & \leq C(1+T) \left(\|u_1\|_{H^{k-2}}^2 + \|\nabla u_0\|_{H^{k-1}}^2 \right) \exp \left(\int_0^T (1 + \|\nabla u\|_{L^\infty}^{2k} + \|u_t\|_{L^\infty}^{2k}) ds \right). \end{aligned}$$

At least for small times we want to remove the dependence on u on the right-hand side of (4.22) and thus we introduce the quantity

$$\alpha(t) = \|\nabla u(t)\|_{L^2}^2 + \|\Delta u(t)\|_{L^2}^2 + \|\nabla^k u(t)\|_{L^2}^2 + \|u_t(t)\|_{L^2}^2 + \|\nabla^{k-2} u_t(t)\|_{L^2}^2$$

for $t \in [0, T_{\varepsilon, m})$. We observe that $\alpha(t)$ is equivalent to the square of the Sobolev norms appearing in (4.22). Since the solutions to (3.1) are (locally) unique, our reasoning is also valid for any initial time $t_0 \in (0, T_{\varepsilon, m})$. The estimates (4.19), (4.20) and (4.21) thus imply

$$\alpha(t) - \alpha(t_0) \leq C \int_{t_0}^t (1 + \alpha(s)^k) \alpha(s) ds.$$

By the above arguments, the function α is differentiable a.e. so that

$$(4.24) \quad \frac{d}{dt} \alpha(t) \leq C(1 + \alpha(t)^k) \alpha(t)$$

for a.e. $0 \leq t_0 \leq t < T_{\varepsilon, m}$. We now proceed similarly to [5], where regularization by the (intrinsic) biharmonic energy has been applied in order to obtain the existence of local Schrödinger maps.

Lemma 4.1. *Let $\varepsilon \in (0, 1)$ and take data $u_0, u_1 : \mathbb{R}^n \rightarrow \mathbb{R}^L$ with $u_0(x) \in N$ and $u_1(x) \in T_{u_0(x)}N$ for a.e. $x \in \mathbb{R}^n$ satisfying*

$$(\nabla u_0, u_1) \in H^{k-1}(\mathbb{R}^n) \times H^{k-2}(\mathbb{R}^n) \quad \text{for some } k \in \mathbb{N} \text{ with } k > \lfloor \frac{n}{2} \rfloor + 2.$$

Let $T_{\varepsilon, m} > 0$ be the maximal existence time of the solution $u^\varepsilon : \mathbb{R}^n \times [0, T_{\varepsilon, m}) \rightarrow N$ of (3.1) with $u^\varepsilon(0) = u_0$ and $\partial_t u^\varepsilon(0) = u_1$ from Proposition 3.4. Then there is a time $T_0 = T_0(\|\nabla u_0\|_{H^{k-1}}, \|u_1\|_{H^{k-2}}) > 0$ such that $T_{\varepsilon, m} > T_0$ for all $\varepsilon \in (0, 1)$.

Proof. Let $\varepsilon \in (0, 1)$ and $t \in [0, T_{\varepsilon, m})$. We write $u = u^\varepsilon$. From (4.24) we infer

$$(4.25) \quad \frac{d}{dt} \log \left(\frac{\alpha}{(1 + \alpha^k)^{\frac{1}{k}}} \right) = \frac{\alpha'}{(1 + \alpha^k) \alpha} \leq C,$$

With $\alpha_0 = \alpha(0)$ it follows

$$\begin{aligned} \frac{\alpha(t)^k}{(1 + \alpha(t)^k)} &\leq e^{Ctk} \frac{\alpha_0^k}{(1 + \alpha_0^k)} \leq (1 + 4Ctk) \frac{\alpha_0^k}{(1 + \alpha_0^k)}, \\ \alpha(t)^k &\leq (1 + 4Ctk) \alpha_0^k + 4Ctk \alpha_0^k \alpha^k \end{aligned}$$

for $0 \leq t \leq \frac{1}{8Ck}$, and hence

$$\alpha(t)^k \leq 2(1 + 4Ctk) \alpha_0^k \leq 3 \alpha_0^k$$

for $0 \leq t \leq \frac{1}{8Ck} \min\{1, \frac{1}{\alpha_0^k}\} =: T_0$. Since α and the Sobolev norms are equivalent, we infer

$$(4.26) \quad \|u_t(t)\|_{H^{k-2}}^2 + \|\nabla u(t)\|_{H^{k-1}}^2 \leq c_0(\|u_1\|_{H^{k-2}}^2 + \|\nabla u_0\|_{H^{k-1}}^2)$$

for $t \in [0, \min\{T_{\varepsilon, m}, T_0\})$ and some constant $c_0 = c_0(k, n) > 0$.

We now assume by contradiction that $T_{\varepsilon, m} \leq T_0$ for some (fixed) $\varepsilon \in (0, 1)$. We apply the contraction argument in the proof of Lemma 3.1 for the initial time $t_0 \in [0, T_{\varepsilon, m})$ and data $(u(t_0), u_t(t_0))$ in the fixed-point space $\mathcal{B}_r(T)$ with radius

$$r^k = 3r(t_0)^k := 3 \left(\|\nabla u(t_0)\|_{H^{k-1}} + \|u_t(t_0)\|_{H^{k-2}} \right)^k.$$

Since $t_0 < T_0$, estimate (4.26) yields the uniform bound

$$r(t_0) \leq \sqrt{2c_0}(\|u_1\|_{H^{k-2}}^2 + \|\nabla u_0\|_{H^{k-1}}^2)^{1/2} =: \hat{c}_0.$$

As a result, the time

$$T := \frac{1}{4} \min \left\{ \left(\frac{\sqrt[k]{3} - 1}{\sqrt[k]{3}} \right)^2 \frac{\varepsilon}{\hat{C}^2(1 + 3\hat{c}_0^k)^2}, \frac{\varepsilon}{\hat{C}^2(1 + 6\hat{c}_0^k)^2} \right\}.$$

is less or equal than the time T_δ for $\mathcal{B}_r(T)$ in (3.19). Therefore, the solution can be uniquely extended to $[0, t_0 + T]$ in the regularity class of Proposition 3.4. For $t_0 > T_{\varepsilon, m} - T$ this fact contradicts the maximality of $T_{\varepsilon, m}$, showing the result. \square

5. PROOF OF THE MAIN THEOREM

We now combine the existence result from Proposition 3.4 with Lemma 4.1. Thus, there exists a solution $u^\varepsilon : \mathbb{R}^n \times [0, T_0] \rightarrow N$ of (3.1) for each $\varepsilon \in (0, 1)$, where $T_0 > 0$ only depends on $\|\nabla u_0\|_{H^{k-1}}$ and $\|u_1\|_{H^{k-2}}$. From (4.26) and the inequality

$$\|u^\varepsilon - u_0\|_{L_t^\infty L_x^2} \leq T_0 \|u_t^\varepsilon\|_{L_t^\infty L_x^2},$$

we extract a limit $u : \mathbb{R}^n \times [0, T_0] \rightarrow \mathbb{R}^L$ as $\varepsilon \rightarrow 0^+$ of the solutions $u^\varepsilon|_{[0, T_0]}$ in the sense

$$\nabla^{l_1} u^\varepsilon \xrightarrow{*} \nabla^{l_1} u, \quad u^\varepsilon - u_0 \xrightarrow{*} u - u_0, \quad \text{and} \quad \nabla^{l_2-2} u_t^\varepsilon \xrightarrow{*} \nabla^{l_2-2} u_t \quad \text{in} \quad L^\infty(0, T_0; L^2),$$

where $1 \leq l_1 \leq k$ and $0 \leq l_2 \leq k$. (Here and below we do not indicate that we pass to subsequences.) In particular,

$$u - u_0 \in L^\infty(0, T_0; H^k) \cap W^{1, \infty}(0, T_0; H^{k-2})$$

and $(\nabla u, \partial_t u)$ is weakly continuous in $H^{k-1} \times H^{k-2}$. We first assume $k \geq 4$ (which is no restriction if $n \geq 2$). Estimating the nonlinearity similarly to Section 4, we also deduce from (3.3) and (4.26) that $\partial_t^2 u^\varepsilon \in C^0([0, T_0], H^{k-4})$ is uniformly bounded as $\varepsilon \rightarrow 0^+$. Compactness and Sobolev's embedding further yield

$$\nabla^3 u^\varepsilon \rightarrow \nabla^3 u \quad \text{in} \quad C^0([0, T_0], L_{loc}^2(\mathbb{R}^n)),$$

$$(5.1) \quad \partial_t u^\varepsilon \rightarrow \partial_t u, \quad u^\varepsilon \rightarrow u, \quad \nabla u^\varepsilon \rightarrow \nabla u, \quad \nabla^2 u^\varepsilon \rightarrow \nabla^2 u \quad \text{locally uniformly on} \quad \mathbb{R}^n \times [0, T_0].$$

More precisely for $\alpha \in (0, 1)$ and $v^\varepsilon = u^\varepsilon - u_0$, our a priori estimates and [7, Prop. 1.1.4] imply uniform bounds (in ε) in the spaces

$$(5.2) \quad v^\varepsilon \in C^\alpha H^{k-2\alpha}, \quad \nabla v^\varepsilon \in C^\alpha H^{k-1-2\alpha}, \quad \nabla^2 v^\varepsilon \in C^\alpha H^{k-2-2\alpha}, \quad \partial_t v^\varepsilon \in C^\alpha H^{k-2-2\alpha}.$$

As a result, u takes values in N . Moreover, since (4.22) and (4.26) give

$$(5.3) \quad \int_0^T \|\sqrt{\varepsilon} \nabla u_t^\varepsilon(s)\|_{H^{k-2}}^2 ds \\ \lesssim (T_0(1 + \|u_1\|_{H^{k-2}}^{2k} + \|\nabla u_0\|_{H^{k-1}}^{2k}) + 1)(\|u_1\|_{H^{k-2}}^2 + \|\nabla u_0\|_{H^{k-1}}^2)$$

and $k \geq 3$, we infer that $\varepsilon \Delta \partial_t u^\varepsilon \rightarrow 0$ in $L_{t,x}^2$. Combining this fact with (5.1) and recalling (3.4), we conclude

$$\mathcal{N}_\varepsilon(u^\varepsilon) \rightarrow \mathcal{N}(u) \quad \text{in} \quad L_{loc}^2(\mathbb{R}^n \times [0, T_0]).$$

In the case $n = 1$ and $k = 3$ we obtain the convergence $\mathcal{N}_\varepsilon(u^\varepsilon) \rightarrow \mathcal{N}(u)$ in the sense of the duality (H^1, H^{-1}) because we still have

$$\nabla u^\varepsilon \rightarrow \nabla u, \quad \nabla^2 u^\varepsilon \rightarrow \nabla^2 u, \quad \partial_t u^\varepsilon \rightarrow \partial_t u$$

locally uniformly, as well as $\nabla^3 u^\varepsilon \rightarrow \nabla^3 u$ and $\nabla \partial_t u^\varepsilon \rightarrow \nabla \partial_t u$ in $C^0([0, T_0], H_{loc}^{-1})$ as $\varepsilon \rightarrow 0^+$.

Summing up, we have constructed a local solution $u : [0, T_0] \times \mathbb{R}^n \rightarrow N$ of (1.3) with $u(0) = u_0$ and $\partial_t u(0) = u_1$ such that $(\nabla u, \partial_t u)$ is bounded and weakly continuous in $H^{k-1} \times H^{k-2}$.

In Lemma 6.1 it will be shown that such a solution is locally unique. We recall from the proof of Proposition 4.1 that the solution $u : \mathbb{R}^n \times [0, T] \rightarrow N$ for some $T > 0$ can be extended if $\limsup_{t \rightarrow T^-} (\|\nabla u(t)\|_{H^{k-1}} + \|u_t(t)\|_{H^{k-2}}) < \infty$. There thus exists a maximal time of existence $T_m \in (T_0, \infty]$ of u with

$$\limsup_{t \rightarrow T_m^-} (\|\nabla u(t)\|_{H^{k-1}} + \|u_t(t)\|_{H^{k-2}}) = \infty \quad \text{if} \quad T_m < \infty.$$

Arguing as in Section 4, we establish the energy equality

$$(5.4) \quad \|\nabla^k u\|_{L^2}^2 + \|\nabla^{k-2} u_t\|_{L^2}^2 = 2 \int_0^t \int_{\mathbb{R}^n} \nabla^{k-2}(\mathcal{N}(u)) \cdot \nabla^{k-2} u_t \, dx \, ds \\ + \|\nabla^k u_0\|_{L^2}^2 + \|\nabla^{k-2} u_1\|_{L^2}^2$$

for $t \in [0, T_m)$. (The integral is well-defined in view of the cancellation of one derivative in (4.3).) However, in contrast to the approximations u^ε , the solution u has only k weak spatial derivatives (and $\partial_t u$ has $k-2$). For this reason, when deriving (5.4) we have to replace one spatial derivative by a difference quotient. The details are outlined in Appendix C.

We conclude that the highest derivatives $\nabla^{k-2}u_t, \nabla^k u : [0, T_m) \rightarrow L^2$ are continuous, employing their weak continuity and that the right-hand side of (5.4) is continuous in t . The continuity of the lower order derivatives can be shown as in the next section, so that

$$u - u_0 \in C^0([0, T_m), H^k) \cap C^1([0, T_m), H^{k-2})$$

as asserted. Finally, following the proof of the a priori estimate in Section 4 we can derive the blow-up criterion (1.4), cf. Appendix C.

To show Theorem 1.1 it thus remains to establish the uniqueness statement and the continuous dependence on the initial data, which is done in the next Sections 6 and 7.

6. UNIQUENESS

Lemma 6.1. *Let $u, v : \mathbb{R}^n \times [0, T] \rightarrow N$ be two solutions of (1.2) with initial data $u_0 : \mathbb{R}^n \rightarrow N$ and $u_1 : \mathbb{R}^n \rightarrow \mathbb{R}^L$ such that $u_1 \in T_{u_0}N$ on \mathbb{R}^n and*

$$(\nabla u_0, u_1) \in H^{k-1}(\mathbb{R}^n) \times H^{k-2}(\mathbb{R}^n)$$

for some $k \in \mathbb{N}$ with $k > \lfloor \frac{n}{2} \rfloor + 2$. Also let

$$u - u_0, v - u_0 \in L^\infty(0, T; H^k(\mathbb{R}^n)) \cap W^{1, \infty}(0, T; H^{k-2}(\mathbb{R}^n)).$$

Then $u|_{[0, T]} = v|_{[0, T]}$.

Proof of Lemma 6.1. We derive the uniqueness statement from a Gronwall argument based on the equality

$$(6.1) \quad \frac{d}{dt} \frac{1}{2} \int_{\mathbb{R}^n} |\nabla^l w_t|^2 + |\nabla^{l+2} w|^2 dx = \int_{\mathbb{R}^n} \nabla^l (\mathcal{N}(u) - \mathcal{N}(v)) \cdot \nabla^l w_t dx,$$

for $w = u - v$, $l \in \{0, \dots, k-3\}$ and $t \in [0, T]$, which is a consequence of (1.3). Setting

$$\mathcal{E}(t) = \|w(t)\|_{H^{k-1}}^2 + \|w_t(t)\|_{H^{k-3}}^2,$$

we want to prove

$$(6.2) \quad \frac{d}{dt} \mathcal{E}(t) \leq C(1 + \|\nabla u\|_{H^{k-1}}^{2k} + \|u_t\|_{H^{k-2}}^{2k} + \|\nabla v\|_{H^{k-1}}^{2k} + \|v_t\|_{H^{k-2}}^{2k}) \mathcal{E}(t)$$

for $t \in [0, T]$. We first estimate (6.1) in the case $l = k-3$. Since u and v map into N , we have $\mathcal{N}(u) = (I - P_u)\mathcal{N}(u)$ and analogously for v . It follows

$$\begin{aligned} \mathcal{N}(u) - \mathcal{N}(v) &= (I - P_u)\mathcal{N}(u) - (I - P_v)\mathcal{N}(v) \\ &= (P_v - P_u)\mathcal{N}(u) + (I - P_v)(\mathcal{N}(u) - \mathcal{N}(v)), \end{aligned}$$

and hence

$$\begin{aligned} \nabla^{k-3}(\mathcal{N}(u) - \mathcal{N}(v)) \cdot \nabla^{k-3} w_t &= \nabla^{k-3}[(P_v - P_u)\mathcal{N}(u)] \cdot \nabla^{k-3} w_t \\ &\quad + \nabla^{k-3}[(I - P_v)(\mathcal{N}(u) - \mathcal{N}(v))] \cdot \nabla^{k-3} w_t. \end{aligned}$$

In this way, we can avoid that all derivatives fall on $\nabla^3 w$. We next write

$$\begin{aligned} \nabla^{k-3}[(P_v - P_u)\mathcal{N}(u)] \cdot \nabla^{k-3} w_t &= (P_v - P_u) \nabla^{k-3}[\mathcal{N}(u)] \cdot \nabla^{k-3} w_t \\ &\quad + \sum_{\substack{l_1 + l_2 = k-3 \\ l_1 > 0}} \nabla^{l_1}[(P_v - P_u)] \star \nabla^{l_2}[\mathcal{N}(u)] \cdot \nabla^{k-3} w_t =: I_1 + I_2. \end{aligned}$$

Observe that

$$\int_{\mathbb{R}^n} I_1 dx \lesssim \|w\|_{L^\infty} \|\nabla^{k-3} \mathcal{N}(u)\|_{L^2} \|\nabla^{k-3} w_t\|_{L^2}.$$

We then control $\|\nabla^{k-3}\mathcal{N}(u)\|_{L^2}$ using Lemma 2.2 as above for the a priori estimate (4.22). Further, Lemma A.2 implies that $\int_{\mathbb{R}^n} I_2 dx$ is bounded by terms of the form

$$(6.3) \quad \|w\|_{L^\infty} \|\nabla^{m_1+1}u\| \cdots \|\nabla^{m_j+1}u\| \|\nabla^{l_2}\mathcal{N}(u)\|_{L^2} \|\nabla^{k-3}w_t\|_{L^2},$$

$$(6.4) \quad \|\nabla^{k-3}w_t\|_{L^2} \|\nabla^{m_1+1}w\| \|\nabla^{m_2+1}h_1\| \cdots \|\nabla^{m_j+1}h_{j-1}\| \|\nabla^{l_2}\mathcal{N}(u)\|_{L^2},$$

where m_1, \dots, m_j and h_1, \dots, h_{j-1} are as in Lemma A.2. In (6.3) we then estimate as above in the a priori estimate. For (6.4), it suffices to control terms of the form

$$(6.5) \quad |\nabla^{m_1+1}w| |\nabla^{m_2+1}h_1| \cdots |\nabla^{m_j+1}h_{j-1}| |\nabla^{\tilde{m}_1+1}u| \cdots |\nabla^{\tilde{m}_i+1}u| [|\nabla^{k_1}u_t| |\nabla^{k_2}u_t| \cdots],$$

where $[|\nabla^{k_1}u_t| |\nabla^{k_2}u_t| \cdots]$ is given as in the nonlinearity $\mathcal{N}(u)$ and the orders $m_1, \dots, m_j, \tilde{m}_1, \dots, \tilde{m}_i$, and k_1, k_2, \dots are as used before. To apply Lemma 2.2, as above we choose

$$f_1 = w, f_2 = \nabla h_1, \dots, f_j = \nabla h_{j-1}, f_{j+1} = \nabla u, \dots, f_{i+j} = \nabla u,$$

and $f_{i+j+1}, f_{i+j+2}, \dots$, according to the respective terms in $\mathcal{N}(u)$. We can thus estimate (6.5) in L^2 by

$$\begin{aligned} & \left(\|w\|_{L^\infty}^{1-\frac{m_1}{k-2-i-j}} \|w\|_{H^{k-2-i-j}}^{\frac{m_1}{k-2-i-j}} + \|w\|_{L^\infty}^{1-\frac{m_1}{k-1-i-j}} \|w\|_{H^{k-1-i-j}}^{\frac{m_1}{k-1-i-j}} \right) \\ & \quad \cdot (1 + \|\nabla u\|_{H^{k-1}}^{2k} + \|u_t\|_{H^{k-2}}^{2k} + \|\nabla v\|_{H^{k-1}}^{2k} + \|v_t\|_{H^{k-2}}^{2k}) \\ & \lesssim \|w\|_{H^{k-1}} (1 + \|\nabla u\|_{H^{k-1}}^{2k} + \|u_t\|_{H^{k-2}}^{2k} + \|\nabla v\|_{H^{k-1}}^{2k} + \|v_t\|_{H^{k-2}}^{2k}), \end{aligned}$$

noting that $l_1 > 0$, $j \geq 1$ and $i + j < k - 2$. We continue by computing

$$\begin{aligned} & \nabla^{k-3}[(I - P_v)(\mathcal{N}(u) - \mathcal{N}(v))] \cdot \nabla^{k-3}w_t \\ &= \nabla^{k-3}(\mathcal{N}(u) - \mathcal{N}(v))(I - P_v)\nabla^{k-3}w_t + \sum_{\substack{l_1+l_2=k-3 \\ l_1>0}} \nabla^{l_1}(I - P_v) \star \nabla^{l_2}(\mathcal{N}(u) - \mathcal{N}(v)) \cdot \nabla^{k-3}w_t \\ &= \nabla^{k-3}(\mathcal{N}(u) - \mathcal{N}(v))\nabla^{k-3}[(P_u - P_v)u_t] - \sum_{\substack{l_1+l_2=k-3 \\ l_1>0}} \nabla^{k-3}(\mathcal{N}(u) - \mathcal{N}(v)) \cdot \nabla^{l_1}[(I - P_v)] \star \nabla^{l_2}w_t \\ & \quad + \sum_{\substack{l_1+l_2=k-3 \\ l_1>0}} \nabla^{l_1}(I - P_v) \star \nabla^{l_2}(\mathcal{N}(u) - \mathcal{N}(v)) \cdot \nabla^{k-3}w_t =: J_1 + J_2 + J_3. \end{aligned}$$

where the second equality is a consequence of

$$(I - P_v)w_t = (I - P_v)u_t = [(I - P_v) - (I - P_u)]u_t = (P_u - P_v)u_t.$$

We use integration by parts to treat $\int J_1 dx$ and $\int J_2 dx$. Here we assume that $k \geq 4$. (If $k = 3$ the estimate becomes easier and we only employ integration by parts for $dP_v(\nabla^3 w \star \nabla u)$ in the difference $\mathcal{N}(u) - \mathcal{N}(v)$.) It follows

$$\begin{aligned} \int_{\mathbb{R}^n} J_1 dx &= - \int_{\mathbb{R}^n} \nabla^{k-4}[\mathcal{N}(u) - \mathcal{N}(v)] \cdot \nabla^{k-2}[(P_u - P_v)u_t] dx, \\ \int_{\mathbb{R}^n} J_2 dx &= \sum_{\substack{l_1+l_2=k-3 \\ l_1>0}} \int_{\mathbb{R}^n} \nabla^{k-4}[\mathcal{N}(u) - \mathcal{N}(v)] \cdot [\nabla^{l_1+1}(I - P_v) \star \nabla^{l_2}w_t \\ & \quad + \nabla^{l_1}(I - P_v) \star \nabla^{l_2+1}w_t] dx. \end{aligned}$$

We first bound

$$\int_{\mathbb{R}^n} J_1 dx \lesssim \|\nabla^{k-4}[\mathcal{N}(u) - \mathcal{N}(v)]\|_{L^2} \|\nabla^{k-2}[(P_u - P_v)u_t]\|_{L^2}.$$

Corollary A.3, Lemma A.2 and Lemma 2.2 yield

$$\begin{aligned} \|\nabla^{k-4}[\mathcal{N}(u) - \mathcal{N}(v)]\|_{L^2} &\lesssim (\|w\|_{H^{k-1}} + \|w_t\|_{H^{k-3}}) \\ & \quad \cdot (1 + \|\nabla u\|_{H^{k-1}}^{2k} + \|u_t\|_{H^{k-2}}^{2k} + \|\nabla v\|_{H^{k-1}}^{2k} + \|v_t\|_{H^{k-2}}^{2k}), \end{aligned}$$

$$\|\nabla^{k-2}[(P_u - P_v)u_t]\|_{L^2} \lesssim \|w\|_{H^{k-1}} (1 + \|\nabla u\|_{H^{k-1}}^{2k} + \|u_t\|_{H^{k-2}}^{2k} + \|\nabla v\|_{H^{k-1}}^{2k} + \|v_t\|_{H^{k-2}}^{2k}).$$

The integrals of J_2 and J_3 are treated similarly. Summing up, we obtain

$$\frac{d}{dt} \int_{\mathbb{R}^n} |\nabla^{k-3} w_t|^2 + |\nabla^{k-1} w|^2 dx \lesssim \mathcal{E}(t) (1 + \|\nabla u\|_{H^{k-1}}^{2k} + \|u_t\|_{H^{k-2}}^{2k} + \|\nabla v\|_{H^{k-1}}^{2k} + \|v_t\|_{H^{k-2}}^{2k}).$$

We can similarly derive the estimate (integrating $dP_v(\nabla^3 w \star \nabla u)$ by parts)

$$\frac{d}{dt} \int_{\mathbb{R}^n} |w_t|^2 + |\Delta w|^2 dx \lesssim \mathcal{E}(t) (1 + \|\nabla u\|_{H^{k-1}}^{2k} + \|u_t\|_{H^{k-2}}^{2k} + \|\nabla v\|_{H^{k-1}}^{2k} + \|v_t\|_{H^{k-2}}^{2k}).$$

Interpolation on the left-hand side then yields

$$\frac{d}{dt} \mathcal{E}(t) \lesssim \mathcal{E}(t) (1 + \|\nabla u(t)\|_{H^{k-1}}^{2k} + \|u_t(t)\|_{H^{k-2}}^{2k} + \|\nabla v(t)\|_{H^{k-1}}^{2k} + \|v_t(t)\|_{H^{k-2}}^{2k}).$$

By assumption, we have $\mathcal{E}(0) = 0$ and

$$\sup_{t \in [0, T]} (\|\nabla u(t)\|_{H^{k-1}}^{2k} + \|u_t(t)\|_{H^{k-2}}^{2k} + \|\nabla v(t)\|_{H^{k-1}}^{2k} + \|v_t(t)\|_{H^{k-2}}^{2k}) < \infty,$$

so that \mathcal{E} on $[0, T]$ as asserted. \square

7. CONTINUITY OF THE FLOW MAP

We now prove that the solutions of the Cauchy problem for (1.2) depend continuously on the initial data. As seen in the previous section, the difference $u - v$ of two solutions u and v satisfies estimates in which one loses a derivative compared the a priori bounds such as (4.22) for the solutions u and v themselves. To deal with this problem, we apply the Bona-Smith argument, which is outlined e.g. in [13] (for the Burgers equation) and in [2] (for the KdV equation); see also the references therein.

Let T_m be the maximal existence time of the solution u with initial data (u_0, u_1) from Theorem 1.1. Fix $T_0 \in (0, T_m)$. Take data (v_0, v_1) as in the theorem satisfying

$$(7.1) \quad \|(u_0, u_1) - (v_0, v_1)\|_{H^k \times H^{k-2}} \leq R$$

for some $R > 0$. (We note that we have to assume $u_0 - v_0 \in L^2$ in order to establish the a priori estimate for the difference of the solutions as in the Section 6.) We use regularized data (u_0^δ, u_1^δ) and (v_0^δ, v_1^δ) in the sense of Lemma B.1 from Appendix B, where $\delta \in (0, \delta^*]$ for some $\delta^* > 0$ depending on N . The corresponding solutions are denoted by u^δ and v^δ . They satisfy the regularity assertions of part a) of Theorem 1.1 for *all* $k > \lfloor \frac{n}{2} \rfloor + 2$. It is crucial that the a priori estimates for u^δ and v^δ are uniform in δ . We split $u - v$ into

$$u - v = u - u^\delta + u^\delta - v^\delta + v^\delta - v$$

and bound each of the differences in $H^k \times H^{k-2}$.

In order to estimate $u^\delta - u$ and $v^\delta - v$, we use the geometric structure (as before in Section 6). It allows us to fix a (small) parameter $\delta > 0$ for which the differences are small in $H^k \times H^{k-2}$. This can be done uniformly for (v_0, v_1) in a certain ball around (u_0, u_1) . For fixed δ , one can then estimate $u^\delta - v^\delta$ employing their extra regularity, but paying the price of a large constant (arising from the small parameter δ). We can control this constant, however, by choosing a small radius $R > 0$ in (7.1).

We start with some preparations concerning the cancellations caused by the geometric constraints. As in Section 6, we have

$$(7.2) \quad \begin{aligned} \mathcal{N}(u^\delta) - \mathcal{N}(u) &= (P_u - P_{u^\delta})(\mathcal{N}(u^\delta)) + (I - P_u)(\mathcal{N}(u^\delta) - \mathcal{N}(u)), \\ (I - P_u)(u^\delta - u)_t &= (P_{u^\delta} - P_u)u_t^\delta. \end{aligned}$$

We then calculate (again similar to Section 6)

$$(7.3) \quad \begin{aligned} &\int_{\mathbb{R}^n} \nabla^{k-2}(\mathcal{N}(u^\delta) - \mathcal{N}(u)) \cdot \nabla^{k-2}(u^\delta - u)_t dx \\ &= \int_{\mathbb{R}^n} (P_u - P_{u^\delta}) \nabla^{k-2}[\mathcal{N}(u^\delta)] \cdot \nabla^{k-2}(u^\delta - u)_t dx \end{aligned}$$

$$\begin{aligned}
& + \sum_{\substack{l_1+l_2=k-2 \\ l_1>0}} \int_{\mathbb{R}^n} \nabla^{l_1} [(P_{u^\delta} - P_u)] \star \nabla^{l_2} \mathcal{N}(u^\delta) \cdot \nabla^{k-2}(u^\delta - u)_t dx \\
& + \sum_{\substack{l_1+l_2=k-2 \\ l_1>0}} \int_{\mathbb{R}^n} \nabla^{l_1} (I - P_u) \star \nabla^{l_2} [\mathcal{N}(u^\delta) - \mathcal{N}(u)] \cdot \nabla^{k-2}(u^\delta - u)_t dx \\
& + \int_{\mathbb{R}^n} \nabla^{k-2} [\mathcal{N}(u^\delta) - \mathcal{N}(u)] \cdot (I - P_u) \nabla^{k-2}(u^\delta - u)_t dx.
\end{aligned}$$

Using integration by parts and (7.2), the last term is rewritten as

$$\begin{aligned}
(7.4) \quad & \int_{\mathbb{R}^n} \nabla^{k-2} [\mathcal{N}(u^\delta) - \mathcal{N}(u)] \cdot (I - P_u) \nabla^{k-2}(u^\delta - u)_t dx \\
& = \sum_{\substack{l_1+l_2=k-2 \\ l_1>0}} \int_{\mathbb{R}^n} \nabla^{k-3} [\mathcal{N}(u^\delta) - \mathcal{N}(u)] \cdot \nabla(\nabla^{l_1} (I - P_u) \star \nabla^{l_2}(u^\delta - u)_t) dx \\
& - \sum_{\substack{l_1+l_2=k-1 \\ l_1>0}} \int_{\mathbb{R}^n} \nabla^{k-3} [\mathcal{N}(u^\delta) - \mathcal{N}(u)] \cdot \nabla^{l_1} [(P_{u^\delta} - P_u)] \star \nabla^{l_2} u_t^\delta dx \\
& - \int_{\mathbb{R}^n} \nabla^{k-3} [\mathcal{N}(u^\delta) - \mathcal{N}(u)] \cdot (P_{u^\delta} - P_u) \nabla^{k-1} u_t^\delta dx,
\end{aligned}$$

which is well defined by the higher regularity of u^δ . Technically this has to be established by difference quotients as in Appendix C, however we omit the details here. The advantage of estimating $u^\delta - u$ is that the *bad terms* (with respect to the regularity of u)

$$(7.5) \quad \|\nabla^{k-2} \mathcal{N}(u^\delta)\|_{L^2} \quad \text{and} \quad \|\nabla^{k-1} u_t^\delta\|_{L^2}$$

will be bounded by the regularized initial data from Lemma B.1. Their norm will grow as $\delta \rightarrow 0^+$ in a controlled way. Moreover, when estimating (7.3) and (7.4), these bad terms only appear in the products

$$\begin{aligned}
& \|u^\delta - u\|_{L^\infty} \|\nabla^{k-2} \mathcal{N}(u^\delta)\|_{L^2} \|\nabla^{k-2}(u^\delta - u)_t\|_{L^2}, \\
& \|u^\delta - u\|_{L^\infty} \|\nabla^{k-3} (\mathcal{N}(u^\delta) - \mathcal{N}(u))\|_{L^2} \|\nabla^{k-1} u_t^\delta\|_{L^2}.
\end{aligned}$$

Here the decay of $\|u^\delta - u\|_{L^\infty}$ as $\delta \rightarrow 0^+$ will compensate the growth in (7.5). We now carry out the details in several steps.

Step 1. Since $T_0 < T_m$, we have the bound

$$\sup_{t \in [0, T_0]} (\|\nabla u(t)\|_{H^{k-1}} + \|u_t(t)\|_{H^{k-2}}) =: \bar{C} < \infty.$$

Lemma B.1 allows us to fix a parameter $\delta'_1 \in (0, \delta^*]$ depending on (u_0, u_1) such that

$$(7.6) \quad \|(\nabla u_0^\delta, u_1^\delta)\|_{H^{k-1} \times H^{k-2}} \leq 3\bar{C}/2$$

for all $\delta \in (0, \delta'_1]$. We let $\delta \in (0, \delta'_1]$ and also $R \leq \bar{C}/2$ in (7.1). Hence

$$(7.7) \quad \|(\nabla v_0, v_1)\|_{H^{k-1} \times H^{k-2}} \leq \|(\nabla u_0, u_1)\|_{H^{k-1} \times H^{k-2}} + R \leq 3\bar{C}/2,$$

$$(7.8) \quad \|(\nabla v_0^\delta, v_1^\delta)\|_{H^{k-1} \times H^{k-2}} \leq \|(\nabla u_0^\delta, u_1^\delta)\|_{H^{k-1} \times H^{k-2}} + R \leq 2\bar{C}.$$

We define a time $\tilde{T}_0 > 0$ as in Lemma 4.1, replacing $\alpha(0)$ there by a multiple of \bar{C} . We then combine the uniform a priori bound (4.26) for the approximate solution to the ϵ -problem for v on $[0, \tilde{T}_0]$ with (7.7). Likewise one treats u^δ and v^δ using (7.6) and (7.8), respectively. Following the existence proof in Section 5, we then see that the solutions $u_{[0, \tilde{T}_0]}$, $v_{[0, \tilde{T}_0]}$, $u_{[0, \tilde{T}_0]}^\delta$, and $v_{[0, \tilde{T}_0]}^\delta$ exist on $[0, \tilde{T}_0]$. Proceeding as in Sections 4 and 6, we further obtain a constant $\tilde{C} = \tilde{C}(N, k, \tilde{T}_0) > 0$ such that

$$(7.9) \quad \|\nabla u\|_{H^m}^2 + \|u_t\|_{H^{m-1}}^2 \leq \tilde{C} (\|\nabla u_0\|_{H^m}^2 + \|u_1\|_{H^{m-1}}^2),$$

$$(7.10) \quad \|\nabla v\|_{H^m}^2 + \|v_t\|_{H^{m-1}}^2 \leq \tilde{C}(\|\nabla v_0\|_{H^m}^2 + \|v_1\|_{H^{m-1}}^2),$$

$$(7.11) \quad \|u - v\|_{H^m}^2 + \|u_t - v_t\|_{H^{m-2}}^2 \leq \tilde{C}(\|u_0 - v_0\|_{H^m}^2 + \|u_1 - v_1\|_{H^{m-2}}^2).$$

on $[0, \tilde{T}_0]$ and for orders $m \in \{2, \dots, k-1\}$. Analogously, u^δ and v^δ satisfy the estimates (7.9) respectively (7.10), and the differences $u - u^\delta$, $v - v^\delta$ and $u^\delta - v^\delta$ fulfill (7.11) with the same constant \tilde{C} independent of $\delta \in (0, \delta^*]$. For the regularized data we can replace here k by $k+1$, deriving

$$(7.12) \quad \begin{aligned} \|\nabla u^\delta\|_{H^k}^2 + \|u_t^\delta\|_{H^{k-1}}^2 &\leq \tilde{C}(\|\nabla u_0^\delta\|_{H^k}^2 + \|u_1^\delta\|_{H^{k-1}}^2), \\ \|\nabla v^\delta\|_{H^k}^2 + \|v_t^\delta\|_{H^{k-1}}^2 &\leq \tilde{C}(\|\nabla v_0^\delta\|_{H^k}^2 + \|v_1^\delta\|_{H^{k-1}}^2). \end{aligned}$$

Step 2. Estimating (7.3) and (7.4) as in Section 6, we derive

$$\begin{aligned} \frac{d}{dt} (\|u - u^\delta\|_{H^k}^2 + \|u_t - u_t^\delta\|_{H^{k-2}}^2) &\leq C \|u - u^\delta\|_{L^\infty} \|\nabla^{k-2} \mathcal{N}(u^\delta)\|_{L^2} \|\nabla^{k-2}(u_t - u_t^\delta)\|_{L^2} \\ &\quad + C \|u - u^\delta\|_{L^\infty} \|\nabla^{k-3}(\mathcal{N}(u^\delta) - \mathcal{N}(u))\|_{L^2} \|\nabla^{k-1} u_t^\delta\|_{L^2} \\ &\quad + C(\|u - u^\delta\|_{H^k}^2 + \|u_t - u_t^\delta\|_{H^{k-2}}^2) \end{aligned}$$

for some $C = C(N, \bar{C}, \tilde{C}) > 0$. The nonlinearities are treated as in Sections 4 and 6. Using also (7.9), (7.11) and (7.12), we then conclude

$$\begin{aligned} \frac{d}{dt} (\|u - u^\delta\|_{H^k}^2 + \|u_t - u_t^\delta\|_{H^{k-2}}^2) &\leq C \|u - u^\delta\|_{H^{k-1}} (1 + \|\nabla u^\delta\|_{H^k} + \|u_t^\delta\|_{H^{k-2}}) (\|u_t\|_{H^{k-2}} + \|u_t^\delta\|_{H^{k-2}}) \\ &\quad + C \|u - u^\delta\|_{H^{k-1}} (1 + \|\nabla u\|_{H^{k-1}} + \|\nabla u^\delta\|_{H^{k-1}} + \|u_t\|_{H^{k-3}} + \|u_t^\delta\|_{H^{k-3}}) \|u_t^\delta\|_{H^{k-1}} \\ &\quad + C(\|u - u^\delta\|_{H^k}^2 + \|u_t - u_t^\delta\|_{H^{k-2}}^2) \\ &\leq C(\|u_0 - u_0^\delta\|_{H^{k-1}} + \|u_t - u_t^\delta\|_{H^{k-3}}) (1 + \|\nabla u_0^\delta\|_{H^k} + \|u_1^\delta\|_{H^{k-1}}) \\ &\quad + C(\|u - u^\delta\|_{H^k}^2 + \|u_t - u_t^\delta\|_{H^{k-2}}^2) \end{aligned}$$

on $[0, \tilde{T}_0]$. Gronwall's inequality and Lemma B.1 thus yield

$$\begin{aligned} \sup_{t \in [0, \tilde{T}_0]} (\|u - u^\delta\|_{H^k}^2 + \|u_t - u_t^\delta\|_{H^{k-2}}^2) &\leq \frac{C\tilde{T}_0}{\sqrt{\delta}} (\|u_0 - u_0^\delta\|_{H^{k-1}} + \|u_1 - u_1^\delta\|_{H^{k-3}}) + C(\|u_0 - u_0^\delta\|_{H^k}^2 + \|u_1 - u_1^\delta\|_{H^{k-2}}^2) = o(1) \end{aligned}$$

as $\delta \rightarrow 0^+$. In view of our a priori bounds, we can estimate $v - v^\delta$ in the same way. Here we have to split the initial values, obtaining

$$\begin{aligned} \sup_{t \in [0, \tilde{T}_0]} (\|v - v^\delta\|_{H^k}^2 + \|v_t - v_t^\delta\|_{H^{k-2}}^2) &\leq \frac{C\tilde{T}_0}{\sqrt{\delta}} (\|v_0 - v_0^\delta\|_{H^{k-1}} + \|v_1 - v_1^\delta\|_{H^{k-3}}) + C(\|v_0 - v_0^\delta\|_{H^k}^2 + \|v_1 - v_1^\delta\|_{H^{k-2}}^2) \\ &\leq \frac{C\tilde{T}_0}{\sqrt{\delta}} (\|u_0 - u_0^\delta\|_{H^{k-1}} + \|u_1 - u_1^\delta\|_{H^{k-3}}) + C(\|u_0 - u_0^\delta\|_{H^k}^2 + \|u_1 - u_1^\delta\|_{H^{k-2}}^2) \\ &\quad + \frac{C\tilde{T}_0}{\sqrt{\delta}} (\|u_0 - v_0\|_{H^{k-1}} + \|u_1 - v_1\|_{H^{k-3}} + \|u_0^\delta - v_0^\delta\|_{H^{k-1}} + \|u_1^\delta - v_1^\delta\|_{H^{k-3}}) \\ &\quad + C(\|u_0 - v_0\|_{H^k}^2 + \|u_1 - v_1\|_{H^{k-2}}^2 + \|u_0^\delta - v_0^\delta\|_{H^k}^2 + \|u_1^\delta - v_1^\delta\|_{H^{k-2}}^2). \end{aligned}$$

Lemma B.1 now implies that

$$\sup_{t \in [0, \tilde{T}_0]} (\|v - v^\delta\|_{H^k}^2 + \|v_t - v_t^\delta\|_{H^{k-2}}^2)$$

$$\begin{aligned} &\leq \frac{C\tilde{T}_0}{\sqrt{\delta}} (\|u_0 - u_0^\delta\|_{H^{k-1}} + \|u_1 - u_1^\delta\|_{H^{k-3}}) + C(\|u_0 - u_0^\delta\|_{H^k}^2 + \|u_1 - u_1^\delta\|_{H^{k-2}}^2) \\ &\quad + \frac{C\tilde{T}_0}{\sqrt{\delta}} R + CR^2. \end{aligned}$$

On the regularized level, we use the coarse estimate

$$\begin{aligned} \sup_{t \in [0, \tilde{T}_0]} (\|u^\delta - v^\delta\|_{H^k}^2 + \|u_t^\delta - v_t^\delta\|_{H^{k-2}}^2) &\leq \frac{C}{\sqrt{\delta}} \tilde{T}_0 (\|v_0^\delta - u_0^\delta\|_{H^k} + \|v_1^\delta - u_1^\delta\|_{H^{k-2}}) \\ &\quad + C(\|u_0^\delta - v_0^\delta\|_{H^k}^2 + \|u_1^\delta - v_1^\delta\|_{H^{k-2}}^2) \\ &\leq \frac{C\tilde{T}_0}{\sqrt{\delta}} R + CR^2. \end{aligned}$$

Since $u - v = u - u^\delta + u^\delta - v^\delta + v^\delta - v$, it follows

$$(7.13) \quad \sup_{t \in [0, \tilde{T}_0]} (\|u - v\|_{H^k}^2 + \|u_t - v_t\|_{H^{k-2}}^2) \leq \frac{C\tilde{T}_0}{\sqrt{\delta}} (\|u_0 - u_0^\delta\|_{H^{k-1}} + \|u_1 - u_1^\delta\|_{H^{k-3}}) \\ + C(\|u_0 - u_0^\delta\|_{H^k}^2 + \|u_1 - u_1^\delta\|_{H^{k-2}}^2) \\ + \frac{C\tilde{T}_0}{\sqrt{\delta}} R + CR^2.$$

Now take $\eta \in (0, \bar{C}/2]$ and $r_1 \in (0, \eta]$. We first fix $\delta = \delta_1 = \delta_1(r_1) \in (0, \delta'_1]$ and then choose $R_1 = R_1(\delta_1) \in (0, \bar{C}/2]$ such that for all $R \in (0, R_1]$ we have

$$(7.14) \quad \sup_{t \in [0, \tilde{T}_0]} (\|u - v\|_{H^k}^2 + \|u_t - v_t\|_{H^{k-2}}^2) \leq r_1 \leq \eta.$$

In the above reasoning we now replace (u_0, u_1) with corresponding solution u by data (\hat{u}_0, \hat{u}_1) with solution \hat{u} that satisfy the same assumptions as (v_0, v_1) . The function \hat{u} thus fulfills the same a priori estimates as v and also (7.14). Moreover, we assume that

$$(7.15) \quad \|(\hat{u}_0, \hat{u}_1) - (v_0, v_1)\|_{H^k \times H^{k-2}} \leq \hat{R}$$

for some radius $\hat{R} > 0$. We can then repeat the above arguments replacing u by \hat{u} . The resulting regularization parameter $\hat{\delta}_1$ depends on \hat{u} , and thus also the upper bound $\hat{R}_1 = \hat{R}_1(\hat{\delta}_1)$ for the radii in (7.15). For given $0 \leq \hat{r}_1 \leq \hat{\eta}$, we infer

$$(7.16) \quad \sup_{t \in [0, \tilde{T}_0]} (\|\hat{u} - v\|_{H^k}^2 + \|\hat{u}_t - v_t\|_{H^{k-2}}^2) \leq \hat{r}_1 \leq \hat{\eta}$$

provided that $0 < \hat{R} \leq \hat{R}_1$ in (7.15).

Step 3. In the case $\tilde{T}_0 \geq T_0$ the proof is complete. Otherwise we repeat the same argument starting from

$$(u_0^{(1)}, u_1^{(1)}) = (u(\tilde{T}_0), u_t(\tilde{T}_0)) \quad \text{and} \quad (v_0^{(1)}, v_1^{(1)}) = (v(\tilde{T}_0), v_t(\tilde{T}_0)).$$

Observe that (7.14) yields

$$\|(\nabla v_0^{(1)}, v_1^{(1)})\|_{H^{k-1} \times H^{k-2}} \leq \eta + \|(\nabla u_0^{(1)}, u_1^{(1)})\|_{H^{k-1} \times H^{k-2}} \leq 3\bar{C}/2.$$

For a sufficiently small $\delta'_2 \in (0, \delta^*]$ and all $\delta \in (0, \delta'_2]$, we derive

$$\|(\nabla (u_0^{(1)})^\delta, (u_1^{(1)})^\delta)\|_{H^{k-1} \times H^{k-2}}, \|(\nabla (v_0^{(1)})^\delta, (v_1^{(1)})^\delta)\|_{H^{k-1} \times H^{k-2}} \leq 2\bar{C}$$

as in (7.6) and (7.8). Based on these bounds we can repeat the arguments of Steps 1 and 2 on the interval $[\tilde{T}_0, \min\{2\tilde{T}_0, T_0\}] =: J_1$. However we have to replace the bound (7.1) involving R by (7.14) which yields

$$\|(u_0^{(1)}, u_1^{(1)}) - (v_0^{(1)}, v_1^{(1)})\|_{H^k \times H^{k-2}} \leq r_1.$$

Let $r_2 \in (0, \eta]$. Lemma B.1 allows us to fix a parameter $\delta = \delta_2 = \delta_2(r_2) \in (0, \delta_2']$ such that

$$\begin{aligned} & \frac{C\tilde{T}_0}{\sqrt{\delta}} (\|u_0^{(1)} - (u_0^{(1)})^\delta\|_{H^{k-1}} + \|u_0^{(1)} - (u_0^{(1)})^\delta\|_{H^{k-3}}) \\ & + C(\|u_0^{(1)} - (u_0^{(1)})^\delta\|_{H^k} + \|u_0^{(1)} - (u_0^{(1)})^\delta\|_{H^{k-2}}) \leq r_2/4. \end{aligned}$$

As in (7.13) we then obtain

$$\sup_{t \in J_1} (\|u - v\|_{H^k}^2 + \|u_t - v_t\|_{H^{k-2}}^2) \leq r_2/4 + r_2/4 + \frac{C\tilde{T}_0}{\sqrt{\delta_2}} r_1 + Cr_1^2 \leq r_2 \leq \eta$$

if we choose r_1 , and hence R , small enough.

Again we can argue in the same way for \hat{u} instead of u , replacing r_i , δ_i and R by \hat{r}_i , $\hat{\delta}_i$ and \hat{R} . For given $0 < \hat{r}_2 \leq \hat{\eta}$, we thus obtain

$$\sup_{t \in J_1} (\|\hat{u} - v\|_{H^k}^2 + \|\hat{u}_t - v_t\|_{H^{k-2}}^2) \leq \hat{r}_2/4 + \hat{r}_2/4 + \frac{C\tilde{T}_0}{\sqrt{\hat{\delta}_2}} \hat{r}_1 + C\hat{r}_1^2 \leq \hat{r}_2 \leq \hat{\eta}$$

if \hat{r}_1 and \hat{R} are small enough.

Step 4.

The previous step can be repeated m times until $m\tilde{T}_0 \geq T_0$. We set $R_0 = R(\bar{C}/2)$ (with $\eta = \bar{C}/2$) and use the resulting radius $\hat{R} = \hat{R}(\hat{\eta})$ for the continuity at \hat{u} , concluding the proof of the continuous dependence and thus of Theorem 1.1.

APPENDIX A. DERIVATIVES OF THE NONLINEARITY

In this section we assume $u, v : \mathbb{R}^n \times [0, T] \rightarrow \mathbb{R}^L$ are smooth maps. The calculations hold if u and v are sufficiently regular to apply the Leibniz formula (e.g. with weak derivatives in L^2). Lemma 2.1 and the Leibniz formula imply the following substitution rule.

Lemma A.1. *Let $l \in \mathbb{N}$. Then we have*

$$\nabla^l(\mathcal{N}(u)) = J_1 + J_2 + J_3,$$

where the terms J_1 , J_2 , and J_3 are of the form (with $k_i, m_i \in \mathbb{N}_0$)

$$J_1 = \sum_{(*)} d^{j+1} P_u(\nabla^{m_1+1} u \star \dots \star \nabla^{m_j+1} u) [\nabla^{k_1} u_t \star \nabla^{k_2} u_t + \nabla^{k_1+2} u \star \nabla^{k_2+2} u + \nabla^{k_1+3} u \star \nabla^{k_2+1} u]$$

with $(*) : 0 \leq m \leq l$, $\sum_{i=1}^2 k_i = l - m$, $j = \min\{1, m\}, \dots, m$, $\sum_{k=1}^j m_k = m - j$;

$$J_2 = \sum_{(*)} d^{j+2} P_u(\nabla^{m_1+1} u \star \dots \star \nabla^{m_j+1} u) [\nabla^{k_1+1} u \star \nabla^{k_2+1} u \star \nabla^{k_3+2} u]$$

with $(*) : 0 \leq m \leq l$, $\sum_{i=1}^3 k_i = l - m$, $j = \min\{1, m\}, \dots, m$, $\sum_{k=1}^j m_k = m - j$;

$$J_3 = \sum_{(*)} d^{j+3} P_u(\nabla^{m_1+1} u \star \dots \star \nabla^{m_j+1} u) [\nabla^{k_1+1} u \star \nabla^{k_2+1} u \star \nabla^{k_3+1} u \star \nabla^{k_4+1} u]$$

with $(*) : 0 \leq m \leq l$, $\sum_{i=1}^4 k_i = l - m$, $j = \min\{1, m\}, \dots, m$, $\sum_{k=1}^j m_k = m - j$.

The following lemmata are used to prove the existence of a fixed point in Section 3 and the uniqueness result in Section 6.

Lemma A.2. *Let $m \in \mathbb{N}$, $k \in \mathbb{N}_0$, and $w = u - v$. For $m \geq 2$ we have*

(A.1)

$$\begin{aligned} \nabla^m(d^k P_u - d^k P_v) &= \sum_{j=1}^m \sum_{m_1+\dots+m_j=m-j} (d^{j+k} P_u - d^{j+k} P_v)(\nabla^{m_1+1} u, \dots, \nabla^{m_j+1} u) \\ &+ \sum_{j=2}^m \sum_{m_1+\dots+m_j=m-j} d^{j+k} P_v(\nabla^{m_1+1} w, \nabla^{m_2+1} u, \dots, \nabla^{m_j+1} u) \end{aligned}$$

$$\begin{aligned}
& + \sum_{j=2}^m \sum_{m_1+\dots+m_j=m-j} d^{j+k} P_v(\nabla^{m_1+1}v, \nabla^{m_2+1}w, \nabla^{m_3+1}u, \dots, \nabla^{m_j+1}u) \\
& : \\
& + \sum_{j=2}^m \sum_{m_1+\dots+m_j=m-j} d^{j+k} P_v(\nabla^{m_1+1}v, \dots, \nabla^{m_{j-1}+1}v, \nabla^{m_j+1}w),
\end{aligned}$$

and for $m = 1$

$$(A.2) \quad \nabla(d^k P_u - d^k P_v) = (d^k P_u - d^k P_v)(\nabla u) + d^k P_v(\nabla w).$$

Proof. The result follows from subtracting the expansion in Lemma 2.1 for $d^k P_v$

$$\nabla^m(d^k P_v) = \sum_{j=1}^m \sum_{m_1+\dots+m_j=m-j} d^{j+k} P_v(\nabla^{m_1+1}v \star \dots \star \nabla^{m_j+1}v),$$

from the same expansion of $\nabla^m(d^k P_u)$. Then subsequently adding and subtracting the intermediate terms in the formula above gives the result. \square

Corollary A.3. Let $m \in \mathbb{N}$, $k \in \mathbb{N}_0$, and $w = u - v$. Then we have

$$\begin{aligned}
& \nabla^m [(dP_u - dP_v)(u_t \cdot u_t + \nabla^2 u \star \nabla^2 u + \nabla^3 u \star \nabla u)] \\
& = \sum_{(*)} (d^{j+1} P_u - d^{j+1} P_v)(\nabla^{m_1+1}u, \dots, \nabla^{m_j+1}u)(\nabla^{k_1}u_t \star \nabla^{k_2}u_t \\
& \quad + \nabla^{k_1+2}u \star \nabla^{k_2+2}u + \nabla^{k_1+3}u \star \nabla^{k_2}u) \\
& + \sum_{(**)} d^{j+1} P_v(\nabla^{m_1+1}w, \nabla^{m_2+1}u, \dots, \nabla^{m_j+1}u)(\nabla^{k_1}u_t \star \nabla^{k_2}u_t \\
& \quad + \nabla^{k_1+2}u \star \nabla^{k_2+2}u + \nabla^{k_1+3}u \star \nabla^{k_2}u) \\
& : \\
& + \sum_{(**)} d^{j+1} P_v(\nabla^{m_1+1}v, \dots, \nabla^{m_{j-1}+1}v, \nabla^{m_j+1}w)(\nabla^{k_1}u_t \star \nabla^{k_2}u_t \\
& \quad + \nabla^{k_1+2}u \star \nabla^{k_2+2}u + \nabla^{k_1+3}u \star \nabla^{k_2}u),
\end{aligned}$$

where $(*) : j = 1, \dots, m$ and $m_1 + \dots + m_j + k_1 + k_2 = m - j$, and $(**) : j = 2, \dots, m$ and $m_1 + \dots + m_j + k_1 + k_2 = m - j$. Likewise we have

$$\begin{aligned}
& \nabla^m [(d^2 P_u - d^2 P_v)(\nabla u \star \nabla u \star \nabla^2 u)] \\
& = \sum_{(*)} (d^{j+2} P_u - d^{j+2} P_v)(\nabla^{m_1+1}u, \dots, \nabla^{m_j+1}u)(\nabla^{k_1+1}u \star \nabla^{k_2+1}u \star \nabla^{k_3+2}u) \\
& + \sum_{(**)} d^{j+2} P_v(\nabla^{m_1+1}w, \nabla^{m_2+1}u, \dots, \nabla^{m_j+1}u)(\nabla^{k_1+1}u \star \nabla^{k_2+1}u \star \nabla^{k_3+2}u) \\
& : \\
& + \sum_{(**)} d^{j+2} P_v(\nabla^{m_1+1}v, \dots, \nabla^{m_{j-1}+1}v, \nabla^{m_j+1}w)(\nabla^{k_1+1}u \star \nabla^{k_2+1}u \star \nabla^{k_3+2}u)
\end{aligned}$$

where $(*) : j = 1, \dots, m$ and $m_1 + \dots + m_j + k_1 + k_2 + k_3 = m - j$, and $(**) : j = 2, \dots, m$ and $m_1 + \dots + m_j + k_1 + k_2 + k_3 = m - j$. Further

$$\begin{aligned}
& \nabla^m [(d^3 P_u - d^3 P_v)(\nabla u \star \nabla u \star \nabla u \star \nabla u)] \\
& = \sum_{(*)} (d^{j+3} P_u - d^{j+3} P_v)(\nabla^{m_1+1}u, \dots, \nabla^{m_j+1}u)(\nabla^{k_1+1}u \star \nabla^{k_2+1}u \star \nabla^{k_3+1}u \star \nabla^{k_4+1}u) \\
& + \sum_{(**)} d^{j+3} P_v(\nabla^{m_1+1}w, \nabla^{m_2+1}u, \dots, \nabla^{m_j+1}u)(\nabla^{k_1+1}u \star \nabla^{k_2+1}u \star \nabla^{k_3+1}u \star \nabla^{k_4+1}u)
\end{aligned}$$

$$\begin{aligned}
& : \\
& + \sum_{(**)} d^{j+3} P_v(\nabla^{m_1+1} v, \dots, \nabla^{m_{j-1}+1} v, \nabla^{m_j+1} w)(\nabla^{k_1+1} u \star \nabla^{k_2+1} u \star \nabla^{k_3+1} u \star \nabla^{k_4+1} u)
\end{aligned}$$

where we sum over $(*) : j = 1, \dots, m$ and $m_1 + \dots + m_j + k_1 + k_2 + k_3 + k_4 = m - j$, $(**)$: $j = 2, \dots, m$ and $m_1 + \dots + m_j + k_1 + k_2 + k_3 + k_4 = m - j$.

Also, the case $m = 1$ is similar.

Proof. The assertions are consequences of the Leibniz rule and Lemma A.2. \square

Corollary A.4. We have for $m \in \mathbb{N}$, $m \geq 2$ and $w = u - v$ that

$$\nabla^m(\mathcal{N}(u) - \mathcal{N}(v))$$

is a linear combination of terms of the form

$$\begin{aligned}
& (d^{j+1} P_u - d^{j+1} P_v)(\nabla^{m_1+1} u, \dots, \nabla^{m_j+1} u)(\nabla^{k_1} u_t \star \nabla^{k_2} u_t \\
& \quad + \nabla^{k_1+2} u \star \nabla^{k_2+2} u + \nabla^{k_1+3} u \star \nabla^{k_2} u), \\
& d^{j+1} P_v(\nabla^{m_1+1} w, \nabla^{m_2+1} h_1, \dots, \nabla^{m_j+1} h_{j-1})(\nabla^{k_1} u_t \star \nabla^{k_2} u_t \\
& \quad + \nabla^{k_1+2} u \star \nabla^{k_2+2} u + \nabla^{k_1+3} u \star \nabla^{k_2} u), \\
& (d^{j+2} P_u - d^{j+2} P_v)(\nabla^{m_1+1} u, \dots, \nabla^{m_j+1} u)(\nabla^{k_1+1} u \star \nabla^{k_2+1} u \star \nabla^{k_3+2} u), \\
& d^{j+2} P_v(\nabla^{m_1+1} w, \nabla^{m_2+1} h_1, \dots, \nabla^{m_j+1} h_{j-1})(\nabla^{k_1+1} u \star \nabla^{k_2+1} u \star \nabla^{k_3+2} u), \\
& (d^{j+3} P_u - d^{j+3} P_v)(\nabla^{m_1+1} u, \dots, \nabla^{m_j+1} u)(\nabla^{k_1+1} u \star \nabla^{k_2+1} u \star \nabla^{k_3+1} u \star \nabla^{k_4+1} u), \\
& d^{j+3} P_v(\nabla^{m_1+1} w, \nabla^{m_2+1} h_1, \dots, \nabla^{m_j+1} h_{j-1})(\nabla^{k_1+1} u \star \nabla^{k_2+1} u \star \nabla^{k_3+1} u \star \nabla^{k_4+1} u), \quad \text{and} \\
& d^{j+1} P_v(\nabla^{m_1+1} v, \dots, \nabla^{m_j+1} v)(\nabla^{k_1} w_t \star \nabla^{k_2} h_t + \nabla^{k_1+2} w \star \nabla^{k_2+2} h \\
& \quad + \nabla^{k_1+3} w \star \nabla^{k_2} h + \nabla^{k_1+3} h \star \nabla^{k_2} w), \quad h \in \{u, v\}, \\
& d^{j+2} P_v(\nabla^{m_1+1} v, \dots, \nabla^{m_j+1} v)(\nabla^{k_1+1} w \star \nabla^{k_2+1} h_1 \star \nabla^{k_3+2} h_2 \\
& \quad + \nabla^{k_1+1} h_1 \star \nabla^{k_2+1} h_2 \star \nabla^{k_3+2} w), \\
& d^{j+3} P_v(\nabla^{m_1+1} v, \dots, \nabla^{m_j+1} v)(\nabla^{k_1+1} w \star \nabla^{k_2+1} h_1 \star \nabla^{k_3+1} h_2 \star \nabla^{k_4+1} h_3),
\end{aligned}$$

where $j, k_1, k_2, k_3, k_4, m_1, \dots, m_j$ and $H, h_1, \dots, h_{j-1} \in \{u, v\}$ are as above in Corollary A.3. Also, we have a similar (but simpler) statement for $m = 1$.

Proof. We write, according to the definition of $\mathcal{N}(u)$ in (1.3),

$$\begin{aligned}
\mathcal{N}(u) - \mathcal{N}(v) &= (dP_u - dP_v)(u_t \cdot u_t + \nabla^2 u \star \nabla^2 u + \nabla^3 u \star \nabla u) \\
& \quad + (d^2 P_u - d^2 P_v)(\nabla u \star \nabla u \star \nabla^2 u) + (d^3 P_u - d^3 P_v)(\nabla u \star \nabla u \star \nabla u \star \nabla u) \\
& \quad + dP_v(w_t \cdot u_t + v_t \cdot w_t + \nabla w \star \nabla u + \nabla v \star \nabla w + \nabla^3 w \star \nabla u + \nabla^3 v \star \nabla w) \\
& \quad + d^2 P_v(\nabla w \star \nabla u \star \nabla^2 u + \nabla v \star \nabla w \star \nabla^2 u + \nabla v \star \nabla v \star \nabla^2 w) \\
& \quad + d^3 P_v(\nabla w \star \nabla u \star \nabla u \star \nabla u + \nabla v \star \nabla w \star \nabla u \star \nabla u \\
& \quad \quad + \nabla v \star \nabla v \star \nabla w \star \nabla u + \nabla v \star \nabla v \star \nabla v \star \nabla w).
\end{aligned}$$

Then, we use Corollary A.3 for the first three terms in the sum above. For the latter three, we use Lemma 2.1 and the Leibniz rule. \square

Let $\varepsilon \in (0, 1)$. We recall from (3.4) the definition

$$\mathcal{N}_\varepsilon(u) = \mathcal{N}(u) - \varepsilon d^2 P_u(u_t, \nabla u, \nabla u) - \varepsilon 2dP_u(\nabla u_t, \nabla u) - \varepsilon dP_u(u_t, \Delta u).$$

Lemma A.5. *For $m \in \mathbb{N}_0$ the derivative $\nabla^m(\mathcal{N}_\varepsilon(u))$ compared to $\nabla^m(\mathcal{N}(u))$ contains the additional terms*

$$d^{j+1}P_u(\nabla^{m_1+1}u \star \dots \star \nabla^{m_j+1}u)(\nabla^{k_1}u_t \star \nabla^{k_2+2}u + \nabla^{k_1+1}u_t \star \nabla^{k_2+1}u), \quad \text{and}$$

$$d^{j+2}P_u(\nabla^{m_1+1}u \star \dots \star \nabla^{m_j+1}u)(\nabla^{k_1}u_t \star \nabla^{k_2+1}u \star \nabla^{k_3+1}u),$$

with $j, m_1, \dots, m_j, k_1, k_2, k_3$ similarly to Lemma A.1.

Further $\nabla^m(\mathcal{N}_\varepsilon(u)) - \nabla^m(\mathcal{N}_\varepsilon(v))$ compared to $\nabla^m(\mathcal{N}(u)) - \nabla^m(\mathcal{N}(v))$ contains additional terms of the form

$$(d^{j+1}P_u - d^{j+1}P_v)(\nabla^{m_1+1}u, \dots, \nabla^{m_j+1}u)(\nabla^{k_1}u_t \star \nabla^{k_2+2}u + \nabla^{k_1+1}u_t \star \nabla^{k_2+1}u),$$

$$d^{j+1}P_v(\nabla^{m_1+1}w, \nabla^{m_2+1}h_1, \dots, \nabla^{m_j+1}h_{j-1})(\nabla^{k_1}u_t \star \nabla^{k_2+2}u + \nabla^{k_1+1}u_t \star \nabla^{k_2+1}u),$$

$$(d^{j+2}P_u - d^{j+2}P_v)(\nabla^{m_1+1}u, \dots, \nabla^{m_j+1}u)(\nabla^{k_1}u_t \star \nabla^{k_2+1}u \star \nabla^{k_3+1}u),$$

$$d^{j+2}P_v(\nabla^{m_1+1}w, \nabla^{m_2+1}h_1, \dots, \nabla^{m_j+1}h_{j-1})(\nabla^{k_1}u_t \star \nabla^{k_2+1}u \star \nabla^{k_3+1}u), \quad \text{and}$$

$$d^{j+1}P_v(\nabla^{m_1+1}v, \dots, \nabla^{m_j+1}v)(\nabla^{k_1}w_t \star \nabla^{k_2+2}h + \nabla^{k_1+1}w_t \star \nabla^{k_2+1}h$$

$$+ \nabla^{k_1}h \star \nabla^{k_2+2}w + \nabla^{k_1+1}h_t \star \nabla^{k_2+1}w), \quad h \in \{u, v\},$$

$$d^{j+2}P_v(\nabla^{m_1+1}v, \dots, \nabla^{m_j+1}v)(\nabla^{k_1}w_t \star \nabla^{k_2+1}h_1 \star \nabla^{k_3+1}h_2$$

$$+ \nabla^{k_1}(h_1)_t \star \nabla^{k_2+1}h_2 \star \nabla^{k_3+1}w),$$

with $w = u - v$ and $j, m_1, \dots, m_j, k_1, k_2, k_3, h_1, \dots, h_{j-1}$ similarly to Corollary A.4.

The implicit constants may depend on ε here.

APPENDIX B. APPROXIMATION OF THE INITIAL DATA

In this section we construct certain approximations of initial data in order to conclude continuous dependence of the solution on the initial data. As in the previous sections, take functions $u_0, u_1 : \mathbb{R}^n \rightarrow \mathbb{R}^L$ with $u_0 \in N$, $u_1 \in T_{u_0}N$ a.e. on \mathbb{R}^n , and

$$(\nabla u_0, u_1) \in H^{k-1}(\mathbb{R}^n) \times H^{k-2}(\mathbb{R}^n).$$

for some $k > \lfloor \frac{n}{2} \rfloor + 2$ with $k \in \mathbb{N}$.

Lemma B.1. *Let the functions (u_0, u_1) be as above. Then there is a number $\delta^* = \delta^*(N) > 0$ such that for $\delta \in (0, \delta^*]$ there exist maps $u_0^\delta, u_1^\delta \in C^\infty(\mathbb{R}^n, \mathbb{R}^L)$ such that $u_0^\delta, u_1^\delta \in H^m$ for all $m \in \mathbb{N}$, $u_0^\delta \in N$ and $u_1^\delta \in T_{u_0^\delta}N$ on \mathbb{R}^n which satisfy*

$$(B.1) \quad u_0 - u_0^\delta \in L^2 \quad \text{and} \quad \|u_0 - u_0^\delta\|_{L^2} \leq C_0 \delta,$$

$$(B.2) \quad \|(\nabla u_0^\delta, u_1^\delta) - (\nabla u_0, u_1)\|_{H^{k-2} \times H^{k-3}} = o(\sqrt{\delta}) \quad \text{as } \delta \rightarrow 0^+,$$

$$(B.3) \quad \|(\nabla u_0^\delta, u_1^\delta) - (\nabla u_0, u_1)\|_{H^{k-1} \times H^{k-2}} = o(1) \quad \text{as } \delta \rightarrow 0^+,$$

$$(B.4) \quad \|(\nabla u_0^\delta, u_1^\delta)\|_{H^k \times H^{k-1}} \leq C_0 \frac{1}{\sqrt{\delta}}$$

for a constant $C_0 = C_0(\|P_p\|_{C_b^k}, \|\nabla u_0\|_{H^{k-1}}, \|u_1\|_{H^{k-2}}) > 0$. Further let (v_0, v_1) be as above with $u_0 - v_0 \in H^k(\mathbb{R}^n)$ and

$$\|(u_0, u_1) - (v_0, v_1)\|_{H^k \times H^{k-2}} \leq R$$

for some $R > 0$. Then for $\delta \in (0, \delta^*]$ we have

$$(B.5) \quad \|(\nabla v_0^\delta, v_1^\delta)\|_{H^k \times H^{k-1}} \leq C_0(1 + R^k) \frac{1}{\sqrt{\delta}},$$

$$(B.6) \quad \|(u_0^\delta, u_1^\delta) - (v_0^\delta, v_1^\delta)\|_{H^k \times H^{k-2}} \leq C_0(1 + R^k) \|(u_0, u_1) - (v_0, v_1)\|_{H^k \times H^{k-2}}.$$

Proof. We choose the caloric extension for regularization, i.e., we consider $\eta_\delta * u_0$ and $\eta_\delta * u_1$ where

$$\eta_\delta(x) = (4\pi\delta)^{-\frac{n}{2}} e^{-\frac{|x|^2}{4\delta}}, \quad \delta > 0, \quad x \in \mathbb{R}^n,$$

and $T(\delta)f = \eta_\delta * f$ is the heat semigroup. Since $u_1 \in C_b^0(\mathbb{R}^n)$ and $u_0 \in C_b^2(\mathbb{R}^n)$ by assumption, the convolution is well defined for u_0 and u_1 . Moreover, $\eta_\delta * u_0$ tends to u_0 and $\eta_\delta * u_1$ to u_1 uniformly as $\delta \rightarrow 0^+$, as well as

$$\nabla(\eta_\delta * u_0) \rightarrow \nabla u_0 \quad \text{in } H^{k-1}(\mathbb{R}^n), \quad \eta_\delta * u_1 \rightarrow u_1 \quad \text{in } H^{k-2}(\mathbb{R}^n) \quad \text{as } \delta \rightarrow 0^+.$$

The uniform convergence yields

$$(B.7) \quad \text{dist}(u_0 * \eta_\delta(x), N) \leq |u_0 * \eta_\delta(x) - u_0(x)| \rightarrow 0 \quad \text{as } \delta \rightarrow 0^+$$

uniformly in $x \in \mathbb{R}^n$. Hence, if $\delta > 0$ is small enough we can define

$$u_0^\delta := \pi(u_0 * \eta_\delta) \quad \text{and} \quad u_1^\delta := P_{u_0 * \eta_\delta}(u_1 * \eta_\delta).$$

Recall that π is the nearest point map and that $P_{u_0 * \eta_\delta}(u_1 * \eta_\delta) \in T_{u_0^\delta} N$ by definition of the projector P and u_0^δ . Especially we have

$$\begin{aligned} |u_0^\delta(x) - u_0 * \eta_\delta(x)| &= \text{dist}(u_0 * \eta_\delta(x), N) \leq |u_0(x) - u_0 * \eta_\delta(x)|, \\ |u_0^\delta(x) - u_0(x)| &\leq 2|u_0(x) - u_0 * \eta_\delta(x)| \end{aligned}$$

for $x \in \mathbb{R}^n$. We further note that u_0^δ and u_1^δ are smooth maps and that we have the uniform convergence

$$u_0^\delta \rightarrow u_0, \quad u_1^\delta \rightarrow u_1$$

as $\delta \rightarrow 0^+$ by construction of u_0^δ (and the mean value theorem for u_1^δ). Assertion (B.1) follows from

$$\|\delta^{-1}(u_0 * \eta_\delta - u_0)\|_{L^2} = \left\| \frac{1}{\delta} \int_0^\delta (\Delta u_0) * \eta_s ds \right\|_{L^2} \lesssim \|\Delta u_0\|_{L^2},$$

by Young's inequality for the convolution. Since $\nabla u_0^\delta = P_{u_0 * \eta_\delta}((\nabla u_0) * \eta_\delta)$, we further have to treat the terms

$$\begin{aligned} P_{u_0 * \eta_\delta}((\nabla u_0) * \eta_\delta) - \nabla u_0 &= P_{u_0 * \eta_\delta}((\nabla u_0) * \eta_\delta - \nabla u_0) + (P_{u_0 * \eta_\delta} - P_{u_0})\nabla u_0, \\ P_{u_0 * \eta_\delta}(u_1 * \eta_\delta) - u_1 &= P_{u_0 * \eta_\delta}(u_1 * \eta_\delta - u_1) + (P_{u_0 * \eta_\delta} - P_{u_0})u_1. \end{aligned}$$

We start by estimating (by means of the mean value theorem for P)

$$\begin{aligned} \|P_{u_0 * \eta_\delta}((\nabla u_0) * \eta_\delta) - \nabla u_0\|_{L^2} &\leq \|P_{u_0 * \eta_\delta}((\nabla u_0) * \eta_\delta - \nabla u_0)\|_{L^2} + \|(P_{u_0 * \eta_\delta} - P_{u_0})\nabla u_0\|_{L^2} \\ &\lesssim \delta \left(O(1) + \|\nabla u_0\|_{L^2} \left\| \frac{1}{\delta}(u_0 * \eta_\delta - u_0) \right\|_{L^\infty} \right), \end{aligned}$$

where $\frac{1}{\delta}(u_0 * \eta_\delta - u_0) \rightarrow \Delta u_0$ uniformly as $\delta \rightarrow 0^+$ since $u_0 \in C_b^2(\mathbb{R}^n)$. Similarly, employing Lemmas 2.1, 2.2 and A.2 as before, we see

$$\begin{aligned} &\|\nabla^{k-2}(P_{u_0 * \eta_\delta}((\nabla u_0) * \eta_\delta) - \nabla u_0)\|_{L^2} \\ &\lesssim \sum_{l_1+l_2=k-2} \left[\|\nabla^{l_1}(P_{u_0 * \eta_\delta}) \cdot \nabla^{l_2}((\nabla u_0) * \eta_\delta - \nabla u_0)\|_{L^2} + \|\nabla^{l_1}(P_{u_0 * \eta_\delta} - P_{u_0}) \cdot \nabla^{l_2+1}u_0\|_{L^2} \right] \\ &\lesssim (1 + \|\nabla u_0\|_{H^{k-2}}^k + \|(\nabla u_0) * \eta_\delta\|_{H^{k-2}}^k) \|(\nabla u_0) * \eta_\delta - \nabla u_0\|_{H^{k-2}} \\ &\quad + \delta \|\nabla^{k-1}u_0\|_{L^2}^k \|\delta^{-1}(u_0 * \eta_\delta - u_0)\|_{L^\infty} \\ &\lesssim o(\sqrt{\delta}) \quad \text{as } \delta \rightarrow 0^+. \end{aligned}$$

Here we also use [7, Prop. 2.2.4]. Interpolation and an analogous argument for u_1^δ in H^{k-3} then allows us to conclude (B.2). Assertion (B.3) is shown in the same way, with $o(1)$ instead of $o(\sqrt{\delta})$ in the upper bound. For (B.4), we compute

$$\begin{aligned} & \left\| \nabla^k (P_{u_0 * \eta_\delta} ((\nabla u_0) * \eta_\delta)) \right\|_{L^2} \\ & \lesssim \sum_{\substack{l_1 + l_2 = k \\ l_1 > 0}} \left\| \nabla^{l_1} (P_{u_0 * \eta_\delta}) \cdot (\nabla^{l_2 + 1} u_0 * \eta_\delta) \right\|_{L^2} + \left\| P_{u_0 * \eta_\delta} \nabla (\nabla^k u_0 * \eta_\delta) \right\|_{L^2} \\ & \lesssim (1 + \|\nabla u_0\|_{H^{k-1}}^k) \|\nabla u_0\|_{H^{k-1}} + \left\| P_{u_0 * \eta_\delta} \nabla (\nabla^k u_0 * \eta_\delta) \right\|_{L^2} \end{aligned}$$

as before. The last term is bounded via

$$\left\| P_{u_0 * \eta_\delta} \nabla (\nabla^k u_0 * \eta_\delta) \right\|_{L^2} \lesssim \left\| (\nabla^k u_0) * \nabla (\eta_\delta) \right\|_{L^2} \lesssim \frac{1}{\sqrt{\delta}} \|\nabla u_0\|_{H^{k-1}}$$

again by Young's inequality. Similarly, the term $\nabla^{k-1} u_1^\delta$ is estimated in $L^2(\mathbb{R}^n)$. The above reasoning also shows (B.5) if we choose the constant $C_0 > 0$ suitably. In order to prove (B.6), similarly as above we compute

$$\left\| u_0^\delta - v_0^\delta \right\|_{L^2} \lesssim \|\eta_\delta * (u_0 - v_0)\|_{L^2} \lesssim \|u_0 - v_0\|_{L^2}.$$

by the mean value theorem and Young's inequality. Writing

$$\begin{aligned} & P_{u_0 * \eta_\delta} ((\nabla u_0) * \eta_\delta) - P_{v_0 * \eta_\delta} ((\nabla v_0) * \eta_\delta) \\ & = P_{u_0 * \eta_\delta} ((\nabla u_0) * \eta_\delta - (\nabla v_0) * \eta_\delta) + (P_{u_0 * \eta_\delta} - P_{v_0 * \eta_\delta}) ((\nabla v_0) * \eta_\delta), \end{aligned}$$

we deduce

$$\begin{aligned} & \left\| \nabla^{k-1} (P_{u_0 * \eta_\delta} ((\nabla u_0) * \eta_\delta) - P_{v_0 * \eta_\delta} ((\nabla v_0) * \eta_\delta)) \right\|_{L^2} \\ & \lesssim \sum_{l_1 + l_2 = k-1} \left\| \nabla^{l_1} (P_{u_0 * \eta_\delta}) \cdot \nabla^{l_2} ((\nabla u_0) * \eta_\delta - (\nabla v_0) * \eta_\delta) \right\|_{L^2} \\ & \quad + \sum_{l_1 + l_2 = k-1} \left\| \nabla^{l_1} (P_{u_0 * \eta_\delta} - P_{v_0 * \eta_\delta}) \cdot (\nabla^{l_2 + 1} v_0) * \eta_\delta \right\|_{L^2} \\ & \lesssim (1 + \|\nabla u_0\|_{H^{k-1}}^k + \|\nabla v_0\|_{H^{k-1}}^k) \|\nabla u_0 - \nabla v_0\|_{H^{k-1}} + \left\| \nabla^k v_0 \right\|_{L^2}^k \|u_0 - v_0\|_{L^\infty} \\ & \lesssim (1 + \|\nabla u_0\|_{H^{k-1}}^k + R^k) \|\nabla u_0 - \nabla v_0\|_{H^{k-1}} + \left\| \nabla^k v_0 \right\|_{L^2}^k \|u_0 - v_0\|_{H^k} \\ & \lesssim (1 + \|\nabla u_0\|_{H^{k-1}}^k + R^k) \|\nabla u_0 - \nabla v_0\|_{H^{k-1}}, \end{aligned}$$

The claim (B.6) then follows by interpolation and a proper choice of $C_0 > 0$. Finally the estimate for

$$u_1^\delta - v_1^\delta = P_{u_0 * \eta_\delta} (u_1 * \eta_\delta - v_1 * \eta_\delta) + (P_{u_0 * \eta_\delta} - P_{v_0 * \eta_\delta}) (v_1 * \eta_\delta)$$

works similarly. \square

APPENDIX C. ESTABLISHING THE IDENTITY (5.4)

For $f, g \in H^1(\mathbb{R}^n)$, $h \in \mathbb{R}$ and $i \in \{1, \dots, n\}$ we set

$$D_h^i f(x) = \frac{1}{h} (f(x + e_i h) - f(x)).$$

Observe that $D_h^i (fg)(x) = (D_h^i f)(x)g(x + e_i h) + f(x)(D_h^i g)(x)$. Since we only use the product rule integrated over $x \in \mathbb{R}^n$ and $g(\cdot + he_i) \rightarrow g$ strongly in H^1 as $h \rightarrow 0$, we drop the h -dependence in $g(\cdot + e_i h)$ in the following calculation.

Let u be the solution of (1.3) obtained in Section 5. We compute

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(\left\| D_h^i \nabla^{k-3} u_t \right\|_{L^2}^2 + \left\| D_h^i \nabla^{k-1} u \right\|_{L^2}^2 \right) = \int_{\mathbb{R}^n} D_h^i \nabla^{k-3} \left((I - P_u) \mathcal{N}(u) \right) \cdot D_h^i \nabla^{k-3} u_t \, dx \\ & = \sum_{l=1}^{k-3} \int_{\mathbb{R}^n} D_h^i (\nabla^l (I - P_u) \star \nabla^{k-3-l} \mathcal{N}(u)) \cdot D_h^i \nabla^{k-3} u_t \, dx \end{aligned}$$

$$\begin{aligned}
& + \int_{\mathbb{R}^n} D_h^i (I - P_u) \nabla^{k-3} \mathcal{N}(u) \cdot D_h^i \nabla^{k-3} u_t \, dx + \int_{\mathbb{R}^n} D_h^i \nabla^{k-3} \mathcal{N}(u) \cdot (I - P_u) D_h^i \nabla^{k-3} u_t \, dx \\
& = \sum_{l=1}^{k-3} \int_{\mathbb{R}^n} D_h^i (\nabla^l (I - P_u) \star \nabla^{k-3-l} \mathcal{N}(u)) \cdot D_h^i \nabla^{k-3} u_t \, dx \\
& + \int_{\mathbb{R}^n} D_h^i (I - P_u) \nabla^{k-3} \mathcal{N}(u) \cdot D_h^i \nabla^{k-3} u_t \, dx + \int_{\mathbb{R}^n} D_h^i (\nabla^{k-3} \mathcal{N}(u) \cdot (I - P_u) D_h^i \nabla^{k-3} u_t) \, dx \\
& + \int_{\mathbb{R}^n} \nabla^{k-3} \mathcal{N}(u) \cdot D_h^i (D_h^i (I - P_u) \nabla^{k-3} u_t) \, dx \\
& + \sum_{l=1}^{k-3} \int_{\mathbb{R}^n} \nabla^{k-3} \mathcal{N}(u) \cdot (D_h^i)^2 (\nabla^l (I - P_u) \star \nabla^{k-3-l} u_t) \, dx =: \int_{\mathbb{R}^n} T_h^i(u) \, dx,
\end{aligned}$$

where the second identity follows from $(I - P_u)u_t = 0$. For a fixed time $t \in [0, T_m)$, the regularity of u yields the limit

$$\begin{aligned}
\lim_{h \rightarrow 0} \int_{\mathbb{R}^n} T_h^i(u(t)) \, dx & = \sum_{l=1}^{k-3} \int_{\mathbb{R}^n} \partial_{x_i} (\nabla^l (I - P_u) \star \nabla^{k-3-l} \mathcal{N}(u)) \cdot \nabla^{k-3} \partial_{x_i} u_t \, dx \\
& - \int_{\mathbb{R}^n} dP_u(\partial_{x_i} u, \nabla^{k-3} \mathcal{N}(u)) \cdot \nabla^{k-3} \partial_{x_i} u_t \, dx \\
& - \int_{\mathbb{R}^n} \nabla^{k-3} \mathcal{N}(u) \cdot \partial_{x_i} (dP_u(\partial_{x_i} u, \nabla^{k-3} u_t)) \, dx \\
& + \sum_{l=1}^{k-3} \int_{\mathbb{R}^n} \nabla^{k-3} \mathcal{N}(u) \cdot \partial_{x_i}^2 (\nabla^l (I - P_u) \star \nabla^{k-3-l} u_t) \, dx \\
& =: \int_{\mathbb{R}^n} T^i(u(t)) \, dx.
\end{aligned}$$

Here we also used that

$$\int_{\mathbb{R}^n} D_h^i (\nabla^{k-3} \mathcal{N}(u) \cdot (I - P_u) D_h^i \nabla^{k-3} u_t) \, dx \rightarrow 0 \quad \text{as } h \rightarrow 0$$

by Gauss' Theorem. Estimating as in Section 4, we derive

$$\left| \int_{\mathbb{R}^n} T^i(u(t)) \, dx \right| \lesssim \sup_{s \in [0, T]} (1 + \|\nabla u(s)\|_{H^{k-1}}^{2k} + \|u_t(s)\|_{H^{k-2}}^{2k}) (\|\nabla u(s)\|_{H^{k-1}}^2 + \|u_t(s)\|_{H^{k-2}}^2).$$

for $t \in [0, T]$ and $T < T_m$. In the limit $h \rightarrow 0$ it follows

$$\|\nabla^{k-3} \partial_{x_i} u_t\|_{L^2}^2 + \|\nabla^{k-1} \partial_{x_i} u\|_{L^2}^2 = 2 \int_0^t \int_{\mathbb{R}^n} T^i(u(s)) \, dx \, ds + \|\nabla^{k-3} \partial_{x_i} u_1\|_{L^2}^2 + \|\nabla^{k-1} \partial_{x_i} u_0\|_{L^2}^2$$

by dominated convergence. The right-hand side is continuous in t , and hence the highest derivatives $\nabla^k u_t, \nabla^{k-2} u : [0, T_m) \rightarrow L^2$ are continuous, since we already know their weak continuity. Finally, summing over $i = 1, \dots, n$ and estimating $T^i(u)$ as in Section 4, we conclude the blow-up criterion from (1.4) for the solution u .

REFERENCES

1. Robert Denk and Roland Schnaubelt. A structurally damped plate equation with Dirichlet-Neumann boundary conditions. *J. Differential Equations*, 259(4):1323–1353, 2015.
2. M. Burak Erdogan and Nikolaos Tzirakis. *Dispersive Partial Differential Equations: Wellposedness and Applications*. Cambridge University Press, 2016.
3. Jishan Fan and Tohru Ozawa. On regularity criterion for the 2D wave maps and the 4D biharmonic wave maps. In *Current advances in nonlinear analysis and related topics*, volume 32 of *GAKUTO Internat. Ser. Math. Sci. Appl.*, pages 69–83. Gakkotosho, Tokyo, 2010.
4. Sebastian Herr, Tobias Lamm, and Roland Schnaubelt. Biharmonic wave maps into spheres. *arXiv e-prints*, December 2018. arXiv:1812.03718.
5. Carlos Kenig, Tobias Lamm, Daniel Pollack, Gigliola Staffilani, and Tatiana Toro. The Cauchy problem for Schrödinger flows into Kähler manifolds. *Discrete Contin. Dyn. Syst.*, 27(2):389–439, 2010.

6. Irena Lasiecka and Roberto Triggiani. *Control theory for partial differential equations: continuous and approximation theories. I*. Cambridge University Press, Cambridge, 2000.
7. Alessandra Lunardi. *Analytic semigroups and optimal regularity in parabolic*. Birkhäuser/ Springer Basel AG, Basel, 1995.
8. Alessandra Lunardi. *Interpolation theory*. Edizioni della Normale, Pisa, 2018. Third edition.
9. Tobias Schmid. Energy bounds for biharmonic wave maps in low dimensions. CRC 1173-Preprint 2018/51, Karlsruhe Institute of Technology, 2018.
10. Christopher D. Sogge. *Lectures on non-linear wave equations*. International Press, Boston, MA, second edition, 2008.
11. Jalal Shatah and Michael Struwe. *Geometric wave equations*. American Mathematical Society, Providence, RI, 1998.
12. Michael E. Taylor. *Partial differential equations III. Nonlinear equations*. Springer, New York, second edition, 2011.
13. Nikolay Tzvetkov. Ill-posedness issues for nonlinear dispersive equations. In *Lectures on nonlinear dispersive equations*, volume 27 of *GAKUTO Internat. Ser. Math. Sci. Appl.*, pages 63–103. Gakkotosho, Tokyo, 2006.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, POSTFACH 10 01 31, 33501 BIELEFELD, GERMANY
E-mail address: herr@math.uni-bielefeld.de

DEPARTMENT OF MATHEMATICS, KARLSRUHE INSTITUTE OF TECHNOLOGY, ENGLERSTRASSE 2, 76131
KARLSRUHE, GERMANY
E-mail address: tobias.lamm@kit.edu, tobias.schmid@kit.edu, roland.schnaubelt@kit.edu