

On the convergence of Lawson methods for semilinear stiff problems

Marlis Hochbruck, Jan Leibold, Alexander Ostermann

CRC Preprint 2019/20, October 2019

KARLSRUHE INSTITUTE OF TECHNOLOGY

CRC 1173



Wave
phenomena

Participating universities



Universität Stuttgart

EBERHARD KARLS
UNIVERSITÄT
TÜBINGEN



Funded by

DFG

ISSN 2365-662X

On the convergence of Lawson methods for semilinear stiff problems

Marlis Hochbruck · Jan Leibold ·
Alexander Ostermann

Received: date / Accepted: date

Abstract Since their introduction in 1967, Lawson methods have achieved constant interest in the time discretization of evolution equations. The methods were originally devised for the numerical solution of stiff differential equations. Meanwhile, they constitute a well-established class of exponential integrators, which has turned out to be competitive for solving space discretizations of certain types of partial differential equations.

The popularity of Lawson methods is in some contrast to the fact that they may have a bad convergence behaviour, since they do not satisfy any of the stiff order conditions. The aim of this paper is to explain this discrepancy. It is shown that non-stiff order conditions together with appropriate regularity assumptions imply high-order convergence of Lawson methods. Note, however, that the term regularity here includes the behaviour of the solution at the boundary. For instance, Lawson methods will behave well in the case of periodic boundary conditions, but they will show a dramatic order reduction for, e.g., Dirichlet boundary conditions. The precise regularity assumptions required for high-order convergence are worked out in this paper and related to the corresponding assumptions for splitting schemes. In contrast to previous work, the analysis is based on expansions of the exact and the numerical solution along the flow of the homogeneous problem. Numerical examples for the Schrödinger equation are included.

Keywords Lawson methods · error analysis · stiff problems

M. Hochbruck and J. Leibold
Institut für Angewandte und Numerische Mathematik, Karlsruher Institut für Technologie,
D-76149 Karlsruhe, Germany
E-mail: marlis.hochbruck@kit.edu, jan.leibold@kit.edu

A. Ostermann
Institut für Mathematik, Universität Innsbruck, A-6020 Innsbruck, Austria
E-mail: alexander.ostermann@uibk.ac.at

1 Introduction

Exponential integrators are a well-established class of methods for the numerical solution of semilinear stiff differential equations. If the problem stems from a spatial semi-discretization of an evolutionary partial differential equation (PDE), the very form of the domain of the spatial differential operator enters the convergence analysis. The *stiff order conditions*, which guarantee a certain order of convergence independently of the considered problem, must be independent of the domain of this operator (which, in general, involves certain boundary conditions). This is the main reason why stiff order conditions for exponential integrators are quite involved (see [7] and [14]).

For particular problems, however, less conditions are often required for obtaining a certain order of convergence.¹ It was already observed in [8] that periodic boundary conditions do not lead to any order reduction in exponential integrators of collocation type in contrast to homogeneous Dirichlet boundary conditions, which restrict the order of convergence considerably (close to the stage order, depending on the precise situation). Full-order convergence for periodic boundary conditions was also noticed in [11] and [2].

A similar behaviour can be observed for Lawson methods which are obtained by a linear variable transformation from (explicit) Runge–Kutta methods (see [13] and Section 2 below). These methods are very attractive, since they can be easily constructed from any known Runge–Kutta method. Unfortunately, Lawson methods exhibit a strong order reduction, in general. For particular problems, however, they show full order of convergence (see [3], [1], and [16]). By construction, Lawson methods do satisfy the order conditions for non-stiff problems. Such conditions will be called *non-stiff* or *classical order conditions* henceforth. However, Lawson methods *do not* satisfy any of the stiff order conditions, as detailed in [7], [9], and [14]. This fact can result in a dramatic order reduction, even down to order one for parabolic problems with homogeneous Dirichlet boundary conditions.

So far, the derivation of (stiff) order conditions for exponential integrators was based on standard expansions of the exact and the numerical solution. There, the main assumption on the problem is that the exact solution and its composition with the nonlinearity are both sufficiently smooth in time (see [7] and [14]). Any additional regularity in space is not of immediate benefit in this analysis. This is in contrast to splitting methods, where spatial regularity usually shows up in form of commutator bounds (see, e.g., [10]).

In this paper, we study the convergence behaviour of Lawson methods for semilinear problems. One of the main contributions of this paper is a different expansion of the solution. It is still based on the variation-of-constants formula but the nonlinearity is expanded along the flow of the homogeneous problem. This expansion can be derived in a systematic way using trees as in [5] and [14]. The expansion of the exact solution is carried out in terms

¹ The same is true for ordinary differential equations (ODEs), where linear problems, e.g., require less order conditions for Runge–Kutta methods than nonlinear ones.

of elementary integrals, that of the numerical solution in terms of elementary quadrature rules. We show that classical, non-stiff order conditions together with (problem-dependent) assumptions on the exact solution give full order of convergence. This involves regularity of the solution in *space and time*. Our main result for Lawson methods is stated in Theorems 4.7 and 4.8. We prove that a Lawson method converges with order p , if the order of the underlying Runge–Kutta method is at least p and the solution satisfies appropriate regularity assumptions. These conditions are studied in detail for methods of orders one and two, respectively, and they are related to the corresponding conditions that arise in the analysis of splitting methods. In particular, this is worked out for the nonlinear Schrödinger equation. Our error analysis also reveals a different behaviour between the first-order Lawson method and the exponential Euler method, which is visible in numerical experiments.

The outline of the paper is as follows. In Section 2, we recall the construction of Lawson methods. The expansion of the numerical and the exact solution in terms of elementary integrals is given in Section 3. There, we also introduce the analytic (finite dimensional) framework which typically occurs when discretizing a semilinear parabolic or hyperbolic PDE in space. Order conditions and convergence results are given in Section 4. The resulting regularity assumptions are discussed in Section 5. These assumptions are related to the corresponding conditions for splitting methods. Numerical examples that illustrate the required regularity assumptions and the proven convergence behaviour are also presented.

2 Lawson methods

Consider a semilinear system of stiff differential equations

$$u'(t) + Au(t) = g(t, u(t)), \quad u(0) = u_0, \quad (1)$$

where the stiffness stems from the linear part of the equation, i.e., from A , which is either an unbounded linear operator or its spatial discretization, i.e., a matrix. The precise assumptions on A and g will be given in Section 3. For the numerical solution of (1), Lawson [13] considered the following local change of variables:

$$w(t) = e^{tA}u(t).$$

Note that when applied to evolution equations, this transformation has to be done in a formal way, since e^{tA} might not be a meaningful object in our general framework.

Inserting the new variables into (1) gives the transformed differential equation

$$\begin{aligned} w'(t) &= e^{tA}(u'(t) + Au(t)) \\ &= e^{tA}g(t, e^{-tA}w(t)), \quad w(0) = u_0. \end{aligned} \quad (2)$$

For the solution of this problem, an s -stage explicit Runge–Kutta method with coefficients b_i, c_i, a_{ij} satisfying the simplifying assumptions $c_1 = 0$ and

$$\sum_{j=1}^{i-1} a_{ij} = c_i, \quad i = 2, \dots, s, \quad (3)$$

is considered. Transforming the Runge–Kutta discretization of (2) back to the original variables yields the corresponding Lawson method for (1)

$$u_{n+1} = e^{-hA}u_n + h \sum_{i=1}^s b_i e^{-(1-c_i)hA} G_{ni}, \quad (4a)$$

$$G_{ni} = g(t_n + c_i h, U_{ni}), \quad (4b)$$

$$U_{ni} = e^{-c_i h A} u_n + h \sum_{j=1}^{i-1} a_{ij} e^{-(c_i - c_j)hA} G_{nj}, \quad i = 1, \dots, s. \quad (4c)$$

Here, u_n is the numerical approximation to the exact solution $u(t)$ at time $t = t_n = nh$, and h is the step size. Note that this method makes explicit use of the action of the matrix exponential function. Depending on the properties of A , the nodes c_1, \dots, c_s have to fulfill additional assumptions, see Assumption 3.1 in the next section. Because of these actions of the matrix exponential, Lawson methods form a particular class of exponential integrators. For a review on such integrators, we refer to [9].

For a non-stiff ordinary differential equation (1), it is obvious that the order of the Runge–Kutta method applied to (2) coincides with that of the corresponding Lawson method applied to (1). It is the aim of this paper to show that this is also true in the stiff situation, if appropriate regularity assumptions hold (we will explain the meaning of regularity in the context of discretized PDEs in Section 5).

3 Expansion of the exact and the numerical solution

By adding $t' = 1$ to (1), the differential equation is transformed to autonomous form. It is well known that Runge–Kutta methods of order at least one satisfying (3) are invariant under this transformation. Therefore, we restrict ourselves henceforth to the autonomous problem

$$u'(t) + Au(t) = g(u(t)), \quad u(0) = u_0. \quad (5)$$

Let X be a Hilbert space or a Banach space with norm $\|\cdot\|$. Our main assumptions on A and g are as follows.

Assumption 3.1 Let A belong to a family \mathcal{F} of linear operators on X such that $-A$ generates a group e^{-tA} satisfying

$$\|e^{-tA}\| \leq C_{\mathcal{F}}, \quad (6)$$

with a moderate constant $C_{\mathcal{F}}$, uniformly for all $t \in \mathbb{R}$ and all operators $A \in \mathcal{F}$. It is sufficient to require that $-A$ generates a bounded semigroup (i.e., (6) for $t \geq 0$), if the nodes c_i of the considered explicit Runge–Kutta method are ordered as $0 = c_1 \leq c_2 \leq \dots \leq c_s \leq 1$.

The set of infinitesimal generators of non-expansive (semi)groups in X is a possible choice for the family \mathcal{F} . In addition, the above assumption is typically satisfied in situations where (5) stems from a spatial discretization of a semilinear parabolic or hyperbolic partial differential equation. The important fact here is that the constant $C_{\mathcal{F}}$ is independent of the spatial mesh width for finite difference and finite element methods, and independent of the number of ansatz functions in spectral methods. As our error bounds derived below do not depend on A itself but only on the constant $C_{\mathcal{F}}$, they also apply to spatially discretized systems.

Assumption 3.2 For a given integer $p \geq 0$, the nonlinearity g is p times differentiable with bounded derivatives in a neighborhood of the solution of (5).

We recall that the solution of (5) can be represented in terms of the variation-of-constants formula

$$u(\theta h) = e^{-\theta h A} u_0 + h \int_0^\theta e^{-(\theta-\lambda)h A} g(u(\lambda h)) d\lambda.$$

Applying this formula recursively and expanding the nonlinearity along the flow of the homogeneous problem yields the following expansion of the exact solution

$$\begin{aligned} u(h) &= e^{-hA} u_0 + h \int_0^1 e^{-(1-\lambda)hA} g\left(e^{-\lambda hA} u_0 + h \int_0^\lambda e^{-(\lambda-\eta)hA} g(u(\eta h)) d\eta\right) d\lambda \\ &= e^{-hA} u_0 + h \int_0^1 e^{-(1-\lambda)hA} g_\lambda d\lambda \\ &\quad + h^2 \int_0^1 e^{-(1-\lambda)hA} g'_\lambda \int_0^\lambda e^{-(\lambda-\eta)hA} g_\eta d\eta d\lambda \\ &\quad + h^3 \int_0^1 e^{-(1-\lambda)hA} g'_\lambda \int_0^\lambda e^{-(\lambda-\eta)hA} g'_\eta \int_0^\eta e^{-(\eta-\xi)hA} g_\xi d\xi d\eta d\lambda \\ &\quad + \frac{1}{2} h^3 \int_0^1 e^{-(1-\lambda)hA} g''_\lambda \left(\int_0^\lambda e^{-(\lambda-\eta)hA} g_\eta d\eta, \int_0^\lambda e^{-(\lambda-\xi)hA} g_\xi d\xi \right) d\lambda \\ &\quad + \mathcal{O}(h^4), \end{aligned} \tag{7}$$

where we have used the shorthand notation

$$g_\eta = g_\eta(u_0) = g(e^{-\eta h A} u_0), \quad g_\eta^{(k)} = g_\eta^{(k)}(u_0) = g^{(k)}(e^{-\eta h A} u_0), \quad k \geq 1. \tag{8}$$

Note that here and throughout the whole section, the constant in the Landau symbol \mathcal{O} only depends on $C_{\mathcal{F}}$ and the derivatives of g , but not explicitly on A itself, i.e., not on the stiffness. Also note that this expansion differs

considerably from the previous work (see, e.g., [7, 14]) where the nonlinearity $g(u(t))$ was expanded with respect to t .

Next we perform a similar expansion of the numerical solution (4), which yields (again in the autonomous case)

$$\begin{aligned}
u_1 &= e^{-hA}u_0 + h \sum_{i=1}^s b_i e^{-(1-c_i)hA} g_{c_i} \\
&\quad + h^2 \sum_{i=1}^s b_i e^{-(1-c_i)hA} g'_{c_i} \sum_{j=1}^{i-1} a_{ij} e^{-(c_i-c_j)hA} g_{c_j} \\
&\quad + h^3 \sum_{i=1}^s b_i e^{-(1-c_i)hA} g'_{c_i} \sum_{j=1}^{i-1} a_{ij} e^{-(c_i-c_j)hA} g'_{c_j} \sum_{k=1}^{j-1} a_{jk} e^{-(c_j-c_k)hA} g_{c_k} \\
&\quad + \frac{1}{2} h^3 \sum_{i=1}^s b_i e^{-(1-c_i)hA} g''_{c_i} \left(\sum_{j=1}^{i-1} a_{ij} e^{-(c_i-c_j)hA} g_{c_j}, \sum_{k=1}^{i-1} a_{ik} e^{-(c_i-c_k)hA} g_{c_k} \right) \\
&\quad + \mathcal{O}(h^4). \tag{9}
\end{aligned}$$

As we have used the variation-of-constants formula and its discrete counterpart, respectively, the expansions of the exact and the numerical solution reflect the well-known tree structure of (explicit) Runge–Kutta methods. In the following we use the classic trees which are well-established for studying the classical order conditions for Runge–Kutta methods; see [5, Section II.2], [4, Section III.1] and references given there.

Let \mathcal{LT} denote the set of rooted labelled trees; see Def. II.2.2 in [5]. Their nodes are numbered in such a way that the number of a children's node is larger than that of the corresponding mother's node. The set \mathcal{T} of unlabelled rooted trees is defined (as in [5, Def. II.2.4]) as equivalence classes of labelled trees under admissible renumbering.

For $\mathcal{S} = \mathcal{T}$ or $\mathcal{S} = \mathcal{LT}$, respectively, and $\tau_1, \dots, \tau_k \in \mathcal{S}$ we denote by $\tau = [\tau_1, \dots, \tau_k] \in \mathcal{S}$ (a k tuple without ordering) the tree that consists of a new root which is connected to the branches τ_1, \dots, τ_k . Furthermore, by $\varrho(\tau)$ we denote the order of the tree $\tau \in \mathcal{S}$. It is defined as the number of nodes of τ . Trees of order less or equal than p are denoted by

$$\mathcal{S}_p = \{\tau \in \mathcal{S} \mid \varrho(\tau) \leq p\}.$$

For $\tau \in \mathcal{T}$ the elementary differential $D(\tau)$ of a smooth function g is defined recursively in the following way. For $\tau = \bullet$ we have $D(\bullet)(w) = g(w)$, and for $\tau = [\tau_1, \dots, \tau_k]$ we have

$$D(\tau)(w) = g^{(k)}(w)(D(\tau_1)(w), \dots, D(\tau_k)(w)).$$

Motivated by the expansion (7) of the exact solution we define elementary integrals.

Definition 3.3 For $\tau \in \mathcal{LT}$ and $0 \leq \zeta \leq 1$ we define the *elementary integral* $G_\zeta(\tau)$, its *integrand* $\Psi_\zeta(\tau)$, and the *multivariate integration operator* $I_\zeta(\tau)$ with its domain of integration $\mathcal{D}_\zeta(\tau)$, recursively in the following way.

(a) For $\tau = \bullet$ and a univariate function f , we set

$$\begin{aligned}\Psi_\zeta(\bullet)(\lambda, w) &= e^{-(\zeta-\lambda)hA} g_\lambda(w), \quad 0 \leq \lambda \leq \zeta, \\ I_\zeta(\bullet)f &= \int_0^\zeta f(\lambda) d\lambda.\end{aligned}$$

(b) For $\tau = [\tau_1, \dots, \tau_k] \in \mathcal{LT}$ and a multivariate function f in $\varrho = \varrho(\tau)$ variables, we set (for $0 \leq \lambda \leq \zeta$)

$$\begin{aligned}\Psi_\zeta(\tau)(\lambda, \lambda_2, \dots, \lambda_\varrho, w) &= e^{-(\zeta-\lambda)hA} g_\lambda^{(k)}(w) (\Psi_\lambda(\tau_1)(\cdot_1, w), \dots, \Psi_\lambda(\tau_k)(\cdot_k, w)), \\ I_\zeta(\tau)f &= \int_0^\zeta I_\lambda(\tau_1) \cdots I_\lambda(\tau_k) f(\lambda, \cdot, \dots, \cdot) d\lambda.\end{aligned}$$

Here, \cdot_j refers to the $\varrho(\tau_j)$ variables corresponding to the j th subtree τ_j , i.e., all the indices numbering the nodes in τ_j , sorted in increasing order. The integral operator $I_\lambda(\tau_j)$ acts on the variables corresponding to τ_j .

Finally, we define for all $\tau \in \mathcal{LT}$ the *elementary integrals* as

$$G_\zeta(\tau)(w) = I_\zeta(\tau) \Psi_\zeta(\tau)(\cdot, w)$$

and set $\Psi(\tau) = \Psi_1(\tau)$, $\mathcal{D}(\tau) = \mathcal{D}_1(\tau)$, $I(\tau) = I_1(\tau)$, and $G(\tau) = G_1(\tau)$.

Example 3.4 With the definition of the integrands we obtain

$$\begin{aligned}\Psi_\zeta(4 \begin{array}{c} \bullet \\ \bullet \\ \bullet \\ \bullet \end{array} 2)(\lambda, \lambda_2, \lambda_3, \lambda_4, w) &= \Psi_\zeta(2 \begin{array}{c} \bullet \\ \bullet \\ \bullet \\ \bullet \end{array} 4)(\lambda, \lambda_2, \lambda_3, \lambda_4, w) \\ &= e^{-(\zeta-\lambda)hA} g_\lambda^{(2)}(w) (\Psi_\lambda(\begin{array}{c} \bullet \\ \bullet \\ \bullet \end{array} 3)(\lambda_2, \lambda_3, w), \Psi_\lambda(\begin{array}{c} \bullet \\ \bullet \end{array} 4)(\lambda_4, w)) \\ &= e^{-(\zeta-\lambda)hA} g_\lambda^{(2)}(w) (e^{-(\lambda-\lambda_2)hA} g'_{\lambda_2}(w) \Psi_{\lambda_2}(\begin{array}{c} \bullet \\ \bullet \end{array} 3)(\lambda_3, w), e^{-(\lambda-\lambda_4)hA} g_{\lambda_4}(w)) \\ &= e^{-(\zeta-\lambda)hA} g_\lambda^{(2)}(w) (e^{-(\lambda-\lambda_2)hA} g'_{\lambda_2}(w) e^{-(\lambda_2-\lambda_3)hA} g_{\lambda_3}(w), e^{-(\lambda-\lambda_4)hA} g_{\lambda_4}(w))\end{aligned}$$

and

$$\begin{aligned}\Psi_\zeta(3 \begin{array}{c} \bullet \\ \bullet \\ \bullet \end{array} 2)(\lambda, \lambda_2, \lambda_3, \lambda_4, w) &= \Psi_\zeta(2 \begin{array}{c} \bullet \\ \bullet \\ \bullet \\ \bullet \end{array} 3)(\lambda, \lambda_2, \lambda_3, \lambda_4, w) \\ &= e^{-(\zeta-\lambda)hA} g_\lambda^{(2)}(w) (\Psi_\lambda(\begin{array}{c} \bullet \\ \bullet \\ \bullet \end{array} 4)(\lambda_2, \lambda_4, w), \Psi_\lambda(\begin{array}{c} \bullet \\ \bullet \end{array} 3)(\lambda_3, w)) \\ &= e^{-(\zeta-\lambda)hA} g_\lambda^{(2)}(w) (e^{-(\lambda-\lambda_2)hA} g'_{\lambda_2}(w) e^{-(\lambda_2-\lambda_4)hA} g_{\lambda_4}(w), e^{-(\lambda-\lambda_3)hA} g_{\lambda_3}(w)).\end{aligned}$$

It is straightforward to verify that the elementary integrals satisfy the recurrence relation

$$G_\zeta(\bullet)(w) = \int_0^\zeta e^{-(\zeta-\lambda)hA} g_\lambda(w) d\lambda,$$

$$G_\zeta(\tau)(w) = \int_0^\zeta e^{-(\zeta-\lambda)hA} g_\lambda^{(k)}(w) (G_\lambda(\tau_1)(w), \dots, G_\lambda(\tau_k)(w)) d\lambda$$

for $\tau = [\tau_1, \dots, \tau_k] \in \mathcal{T}$.

Our assumptions on g and A ensure that the integrand $\Psi_\zeta(\tau)(\cdot, w)$ is bounded if $\tau \in \mathcal{LT}_{p+1}$ for w in a neighborhood of the exact solution of (5) and h sufficiently small.

Remark 3.6 In the nonstiff situation, where $A \equiv 0$, all evaluations of g or its derivatives are at the fixed value w . Thus $G_\zeta(\tau)(w)$ reduces to a multivariate integral over the constant integrand $\Psi_\zeta(\tau)(\cdot, w) \equiv D(\tau)(w)$.

The following theorem shows how the expansion (7) can be expressed as a (truncated) B-series. Here we use the notation from [4, Section III.1].

Theorem 3.7 *The exact solution of (1) satisfies*

$$u(\zeta h) = e^{-\zeta hA} u_0 + B_p(u_0)(\zeta) + \mathcal{O}(h^{p+1}), \quad \zeta \in [0, 1], \quad (10)$$

where we define the B-series for $w \in X$ and $\zeta \in [0, 1]$ as

$$B_p(w)(\zeta) = \sum_{\tau \in \mathcal{T}_p} \frac{h^{\varrho(\tau)}}{\sigma(\tau)} G_\zeta(\tau)(w),$$

with the symmetry coefficients $\sigma(\bullet) = 1$ and

$$\sigma([\tau_1, \dots, \tau_k]) = \sigma(\tau_1) \cdots \sigma(\tau_k) \mu_1! \mu_2! \cdots$$

The integers μ_1, μ_2, \dots specify the number of equal trees among τ_1, \dots, τ_k .

Proof The proof is done by induction. For $p = 0$, the claim follows from the variation-of-constants formula and $B_0(u_0)(\zeta) = 0$, since

$$u(\zeta h) = e^{-\zeta hA} u_0 + h \int_0^\zeta e^{-(\zeta-\lambda)hA} g(u(\lambda h)) d\lambda = e^{-\zeta hA} u_0 + \mathcal{O}(h). \quad (11)$$

The induction step follows the lines of the proof of [4, Lemma III.1.9] with the following modifications: we use the variation-of-constants formula and truncate the series in such a way that only the first p derivatives of g enter the expansion. We omit the details. \square

Now we proceed analogously for the numerical solution starting with the definition of elementary quadrature rules.

Definition 3.8 For $\tau \in \mathcal{LT}$ we define the *multivariate quadrature operators* $\widehat{I}(\tau)$ and $\widehat{I}_i(\tau)$, $i = 1, \dots, s$, recursively in the following way.

(a) For $\tau = \bullet$ and a univariate function f , we set

$$\widehat{I}(\bullet)f = \sum_{j=1}^s b_j f(c_j), \quad \widehat{I}_i(\bullet)f = \sum_{j=1}^{i-1} a_{ij} f(c_j), \quad 1 \leq i \leq s.$$

(b) For $\tau = [\tau_1, \dots, \tau_k] \in \mathcal{LT}$ and a multivariate function f in $\rho(\tau)$ variables, we set

$$\begin{aligned} \widehat{I}(\tau)f &= \sum_{j=1}^s b_j \widehat{I}_j(\tau_1) \cdots \widehat{I}_j(\tau_k) f(c_j, \cdot, \dots, \cdot), \\ \widehat{I}_i(\tau)f &= \sum_{j=1}^{i-1} a_{ij} \widehat{I}_j(\tau_1) \cdots \widehat{I}_j(\tau_k) f(c_j, \cdot, \dots, \cdot), \quad 1 \leq i \leq s. \end{aligned}$$

As in Definition 3.3, the quadrature operator $\widehat{I}_j(\tau_\ell)$ acts on the variables corresponding to labels in τ_ℓ .

Finally, we define the *elementary quadrature rules* in the following way

$$\widehat{G}(\tau)(w) = \widehat{I}(\tau)\Psi(\tau)(\cdot, w), \quad \widehat{G}_i(\tau)(w) = \widehat{I}_i(\tau)\Psi_{c_i}(\tau)(\cdot, w), \quad 1 \leq i \leq s.$$

As for the elementary integrals, also the elementary quadrature rules do not depend on the numbering of the nodes in a tree.

Lemma 3.9 *The elementary quadrature rules $\widehat{G}(\tau)$ and $\widehat{G}_i(\tau)$ are invariant under renumbering of $\tau \in \mathcal{LT}$. Thus, they are well defined for unlabelled rooted trees $\tau \in \mathcal{T}$ by using one representative of the corresponding equivalence class.* \square

It is easy to see from the recursive definitions that the elementary quadrature rules satisfy

$$\begin{aligned} \widehat{G}(\bullet)(w) &= \sum_{j=1}^s b_j e^{-(1-c_j)hA} g_{c_j}(w), \\ \widehat{G}_i(\bullet)(w) &= \sum_{j=1}^{i-1} a_{ij} e^{-(c_i-c_j)hA} g_{c_j}(w), \quad 1 \leq i \leq s \end{aligned}$$

and

$$\begin{aligned} \widehat{G}(\tau)(w) &= \sum_{j=1}^s b_j e^{-(1-c_j)hA} g_{c_j}^{(k)}(w) (\widehat{G}_j(\tau_1)(w), \dots, \widehat{G}_j(\tau_k)(w)), \\ \widehat{G}_i(\tau)(w) &= \sum_{j=1}^{i-1} a_{ij} e^{-(c_i-c_j)hA} g_{c_j}^{(k)}(w) (\widehat{G}_j(\tau_1)(w), \dots, \widehat{G}_j(\tau_k)(w)), \quad 1 \leq i \leq s \end{aligned}$$

for $\tau = [\tau_1, \dots, \tau_k]$. This allows us to express the expansion (9) of the numerical solution in terms of elementary quadrature rules.

Theorem 3.10 *The numerical solution of (5) satisfies*

$$u_1 = e^{-hA}u_0 + \widehat{B}_p(u_0) + \mathcal{O}(h^{p+1}), \quad (12a)$$

$$U_{0i} = e^{-c_i hA}u_0 + \widehat{B}_p^{(i)}(u_0) + \mathcal{O}(h^{p+1}), \quad i = 1, \dots, s, \quad (12b)$$

where we define the numerical B-series for $w \in X$ as

$$\begin{aligned} \widehat{B}_p(w) &= \sum_{\tau \in \mathcal{T}_p} \frac{h^{\varrho(\tau)}}{\sigma(\tau)} \widehat{G}(\tau)(w), \\ \widehat{B}_p^{(i)}(w) &= \sum_{\tau \in \mathcal{T}_p} \frac{h^{\varrho(\tau)}}{\sigma(\tau)} \widehat{G}_i(\tau)(w), \quad i = 1, \dots, s. \end{aligned}$$

Proof The proof is done analogously to the proof of Theorem 3.7. \square

4 Order conditions and convergence

In this section we present a systematic way of deriving general stiff convergence results for Lawson methods based on trees.

The expansions of the exact and the numerical solution in terms of elementary integrals and elementary quadrature rules derived in the previous section allow us to study the local error in the same way as for classical Runge–Kutta methods. In fact we show that the orders of these quadrature rules determine the local error of the Lawson method. A similar strategy was used in the analysis of splitting methods by [10]. General stiff order conditions for exponential Runge–Kutta methods have been derived in [14] and for splitting methods in [6].

We say that the Lawson method is of (stiff) order p if the local error satisfies

$$\|u(h) - u_1\| \leq Ch^{p+1}$$

uniformly for smooth nonlinearities and operators A satisfying Assumption 3.1. This means that the constant C depends on the constant $C_{\mathcal{F}}$ defined in (6) but not on A itself.

Theorem 4.1 *The Lawson method (4) is of order p if*

$$\widehat{G}(\tau)(u_0) - G(\tau)(u_0) = \mathcal{O}(h^{p+1-\varrho(\tau)}) \quad \text{for all } \tau \in \mathcal{T}_p.$$

Proof From Theorems 3.7 and 3.10 we have

$$u(h) - u_1 = \sum_{\tau \in \mathcal{T}_p} \frac{h^{\varrho(\tau)}}{\sigma(\tau)} (G(\tau)(u_0) - \widehat{G}(\tau)(u_0)) + \mathcal{O}(h^{p+1}). \quad (13)$$

This proves the statement. \square

Remark 4.2 The above derivation can be easily generalized to exponential integrators with a fixed linearization

$$U_i = e^{-c_i h A} u_0 + h \sum_{j=1}^{i-1} a_{ij}(-hA)g(U_j), \quad i = 1, \dots, s,$$

$$u_1 = e^{-hA} u_0 + h \sum_{i=1}^s b_i(-hA)g(U_i),$$

cf. [9]. If one replaces $b_i e^{-(1-c_i)hA}$ by $b_i(-hA)$ and $a_{ij} e^{-(c_i-c_j)hA}$ by $a_{ij}(-hA)$ in Definition 3.8, Theorems 3.10 and 4.1 also hold for general exponential Runge–Kutta methods. If the stiff order conditions derived in [7] and [14] are satisfied up to order p , then $\widehat{G}(\tau)(u_0) - G(\tau)(u_0) = \mathcal{O}(h^{p+1-\varrho(\tau)})$ for all $\tau \in \mathcal{T}_p$.

Example 4.3 For the exponential Euler method, where $s = 1$, $c_1 = 0$, and $b_1(z) = \varphi_1(z)$, we have

$$\widehat{G}(\bullet)(u_0) = b_1(-hA)g(u_0) = \varphi_1(-hA)g(u_0) = \int_0^1 e^{-(1-\lambda)hA} g(u_0) d\lambda$$

and thus

$$\widehat{G}(\bullet)(u_0) - G(\bullet)(u_0) = \int_0^1 e^{-(1-\lambda)hA} (g(u_0) - g_\lambda(u_0)) d\lambda.$$

The condition for order one requires that $\|g(u_0) - g_\lambda(u_0)\| \leq Ch$. In the linear case, where $g(u) = Bu$, this can be written as

$$h^{-1}(g(u_0) - g_\lambda(u_0)) = h^{-1}B(I - e^{-\lambda hA})u_0 = B\varphi_1(-\lambda hA)\lambda Au_0. \quad (14)$$

Hence the condition is fulfilled if Au_0 is uniformly bounded, i.e., $u_0 \in \mathcal{D}(A)$. For the convergence, we thus need $u(t) \in \mathcal{D}(A)$ for $t \in [0, T]$.

It might be interesting to compare (14) to the condition given in [9, Lemma 2.13] which was proved by a Taylor series expansion of $g(u(t))$. For linear problems, it reads

$$\|B(Au(t) + Bu(t))\| \leq C. \quad (15)$$

Hence both results require the same regularity, namely that $Au(t)$ is uniformly bounded. Note, however, that (15) does not involve the φ_1 function. The latter decays like $1/z$ as $z \rightarrow \infty$ in the closed left half-plane, hence components corresponding to eigenvalues with large negative real part are damped.

Corollary 4.4 *If the underlying Runge–Kutta method is of (classical) order p then*

$$I(\tau)1 = \widehat{I}(\tau)1 \quad \text{for all } \tau \in \mathcal{LT}_p, \quad (16)$$

where $1 : [0, 1]^{\varrho(\tau)} \rightarrow \mathbb{R} : \lambda \mapsto 1$ denotes the multivariate constant function with value one.

Proof First note that for problems with $A \equiv 0$, we have

$$G(\tau)(w) = I(\tau)D(\tau)(w), \quad \widehat{G}(\tau)(w) = \widehat{I}(\tau)D(\tau)(w).$$

On the one hand, classical Runge–Kutta theory implies that the local error (13) behaves as $\mathcal{O}(h^{p+1})$ for any sufficiently smooth g . On the other hand, the elementary differentials $D(\tau)$ are known to be independent. Hence, we obtain that $G(\tau)(w) = \widehat{G}(\tau)(w)$. The statement follows because the integrand $\Psi(\tau)(\cdot, w) \equiv D(\tau)(w)$ is a constant. This yields (16). \square

Since the convergence analysis of Lawson methods also employs Taylor expansion, we next study quadrature of monomials. For $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_q) \in [0, 1]^q$ and a given vector $\boldsymbol{\kappa} = (\kappa_1, \dots, \kappa_q) \in \mathbb{N}_0^q$ of non-negative integers, we define as usual

$$\boldsymbol{\lambda}^\kappa = \lambda_1^{\kappa_1} \cdot \dots \cdot \lambda_q^{\kappa_q}$$

and set $\kappa! = \kappa_1! \cdot \dots \cdot \kappa_q!$ and $|\kappa| = \kappa_1 + \dots + \kappa_q$. Moreover, we denote the q -variate monomial function of degree $|\kappa|$ by

$$\mathbf{z}^\kappa : [0, 1]^q \rightarrow \mathbb{R} : \boldsymbol{\lambda} \mapsto \boldsymbol{\lambda}^\kappa.$$

We note that $\mathbf{z}^\kappa = \mathbf{z}_1^{\kappa_1} \cdot \dots \cdot \mathbf{z}_q^{\kappa_q}$ and $\mathbf{z}^0 = 1$.

It turns out that a multivariate integration (or quadrature) w.r.t. τ of such monomials corresponds to the integration (or quadrature) of the constant one function w.r.t. a particular higher order tree stemming from τ .

Lemma 4.5 *Let $\tau \in \mathcal{LT}$ and $\boldsymbol{\kappa} \in \mathbb{N}_0^{\varrho(\tau)}$. We denote by $\tau^{(\boldsymbol{\kappa})} \in \mathcal{LT}$ the tree stemming from τ where κ_j leaves are added to its j th node. Then $\varrho(\tau^{(\boldsymbol{\kappa})}) = \varrho(\tau) + |\boldsymbol{\kappa}|$ and*

$$I_\zeta(\tau)\mathbf{z}^\kappa = I_\zeta(\tau^{(\boldsymbol{\kappa})})1, \quad \widehat{I}(\tau)\mathbf{z}^\kappa = \widehat{I}(\tau^{(\boldsymbol{\kappa})})1, \quad \widehat{I}_i(\tau)\mathbf{z}^\kappa = \widehat{I}_i(\tau^{(\boldsymbol{\kappa})})1$$

for $i = 1, \dots, s$.

Before proving this lemma, we illustrate the proof by an example.

Example 4.6 Let $\tau = \bullet_1^{\bullet 2}$ and $\boldsymbol{\kappa} = (\kappa_1, \kappa_2) \in \mathbb{N}_0^2$. Here we have to consider

$$I(\tau)\mathbf{z}^\kappa = I(\bullet_1^{\bullet 2})\mathbf{z}^\kappa = \int_0^1 \int_0^{\lambda_1} \lambda_1^{\kappa_1} \lambda_2^{\kappa_2} d\lambda_2 d\lambda_1 = \frac{1}{(\kappa_1 + \kappa_2 + 2)(\kappa_2 + 1)},$$

$$\widehat{I}(\tau)\mathbf{z}^\kappa = \widehat{I}(\bullet_1^{\bullet 2})\mathbf{z}^\kappa = \sum_{i,j=1}^s b_i a_{ij} c_i^{\kappa_1} c_j^{\kappa_2}.$$

Writing $\lambda_j = I_{\lambda_j}(\bullet)1$, we observe that

$$I(\tau)\boldsymbol{\lambda}^\kappa = I(\tau^{(\boldsymbol{\kappa})})1, \quad \text{with} \quad \tau^{(\boldsymbol{\kappa})} = \begin{array}{c} \circ \quad \circ \quad \circ \\ \circ \quad \circ \quad \circ \\ \bullet_1 \quad \bullet_2 \end{array},$$

where $\tau^{(\kappa)}$ was obtained by adding κ_1 nodes to the root and κ_2 nodes to the leaf of τ , respectively. For illustration, all newly added nodes are white. They are labelled by $3, \dots, \kappa_1 + \kappa_2 + 2$, but we do not show these labels in the graph because they can be assigned arbitrarily to the white nodes.

Due to the simplifying assumptions, we also have $\widehat{I}(\tau)\mathbf{z}^\kappa = \widehat{I}(\tau^{(\kappa)})1$. For $\varrho(\tau^{(\kappa)}) = \kappa_1 + \kappa_2 + 2 \leq p$, the identity $I(\tau^{(\kappa)})1 = \widehat{I}(\tau^{(\kappa)})1$ holds by Corollary 4.4. In fact, this identity is equivalent to the (conventional) order condition corresponding to the tree $\tau^{(\kappa)}$.

Proof We prove the lemma by induction on $\varrho(\tau)$. The tree $\tau = \bullet$ is the unique tree with $\varrho(\tau) = 1$. For $k \in \mathbb{N}_0$, we have $\bullet^{(k)} = [\bullet]^k$, where $[\bullet]^k = [\bullet, \dots, \bullet]$ denotes the bush with k leaves. Using $\lambda = I_\lambda(\bullet)1$ and the recursive definition of $I_\zeta(\tau)$ we obtain

$$I_\zeta(\bullet)\mathbf{z}^k = \int_0^\zeta \lambda^k d\lambda = \int_0^\zeta (I_\lambda(\bullet)1)^k d\lambda = I_\zeta([\bullet]^k)1 = I_\zeta(\bullet^{(k)})1.$$

Analogously, for $i = 1, \dots, s$, the simplifying assumptions yield $c_i = \widehat{I}_i(\bullet)1$ and this gives

$$\widehat{I}_i(\bullet)\mathbf{z}^k = \sum_{j=1}^{i-1} a_{ij} c_j^k = \sum_{j=1}^{i-1} a_{ij} (\widehat{I}_j(\bullet)1)^k = \widehat{I}_i([\bullet]^k)1 = \widehat{I}_i(\bullet^{(k)})1$$

and $\widehat{I}(\bullet)\mathbf{z}^k = \widehat{I}(\bullet^{(k)})1$.

For the induction step, consider the tree $\tau = [\tau_1, \dots, \tau_m] \in \mathcal{LT}$ and assume that the statement holds true for all τ_j , $j = 1, \dots, m$.

We consider the monomial \mathbf{z}^κ with $|\kappa| = \rho(\tau)$ and write it according to the tree structure of τ as $\mathbf{z}^\kappa = \mathbf{z}_0^k \mathbf{z}_1^{\kappa_1} \dots \mathbf{z}_m^{\kappa_m}$. Here $\mathbf{z}_j^{\kappa_j}$ denotes the monomial containing the variables corresponding to the labels in τ_j with exponents given in the multiindex $\kappa_j \in \mathbb{N}_0^{\varrho(\tau_j)}$ and \mathbf{z}_0^k denotes the monomial corresponding to the root with exponent $k \in \mathbb{N}$.

The recursive definition of $\widehat{I}_i(\tau)$ and the induction hypothesis imply

$$\begin{aligned} \widehat{I}_i(\tau)\mathbf{z}^\kappa &= \sum_{j=1}^{i-1} a_{ij} \widehat{I}_j(\tau_1) \dots \widehat{I}_j(\tau_m) c_j^k \mathbf{z}_1^{\kappa_1} \dots \mathbf{z}_m^{\kappa_m} \\ &= \sum_{j=1}^{i-1} a_{ij} c_j^k (\widehat{I}_j(\tau_1)\mathbf{z}_1^{\kappa_1}) \dots (\widehat{I}_j(\tau_m)\mathbf{z}_m^{\kappa_m}) \\ &= \sum_{j=1}^{i-1} a_{ij} (\widehat{I}_j(\bullet)1)^k (\widehat{I}_j(\tau_1^{\kappa_1})1) \dots (\widehat{I}_j(\tau_m^{\kappa_m})1) \\ &= \widehat{I}_i(\tau^{(\kappa)})1 \end{aligned}$$

for $i = 1, \dots, s$, where we used that the sprouted tree can be cast recursively as

$$\tau^{(\kappa)} = [\underbrace{\bullet, \dots, \bullet}_k, \tau_1^{(\kappa_1)}, \dots, \tau_m^{(\kappa_m)}].$$

The assertion for $I_\zeta(\tau)$ and $\widehat{I}(\tau)$ can be shown analogously. \square

The following theorem provides a sufficient condition for Lawson methods being of (stiff) order p . Here, $C^{k,1}(\mathbb{R}^d, X)$ denotes the space of k times continuously differentiable functions which have a Lipschitz continuous k th derivative.

Theorem 4.7 *Let the integrand $\Psi(\tau)$ of $G(\tau)$ satisfy*

$$\Psi(\tau)(\cdot, u(t)) \in C^{p-\varrho(\tau),1}(\mathcal{D}(\tau), X) \quad \text{for all } \tau \in \mathcal{LT}_p, \quad (17)$$

where $u(t) \in X$ is the solution of (5), $0 \leq t \leq T$. If the underlying Runge–Kutta method is of (classical) order p , then the Lawson method (4) is of (stiff) order p .

Proof Let $\tau \in \mathcal{LT}_p$. We approximate $\Psi(\tau)(\cdot, u_0): \mathcal{D}(\tau) \rightarrow X$ by a multivariate Taylor polynomial of degree $p - \varrho(\tau)$. By assumption on $\Psi(\tau)$, the coefficients and the remainder of this Taylor polynomial are bounded. Using the linearity of the multivariate integrals and quadrature rules, we have by Lemma 4.5

$$\begin{aligned} G(\tau)(u_0) - \widehat{G}(\tau)(u_0) &= I(\tau)\Psi(\tau)(\cdot, u_0) - \widehat{I}(\tau)\Psi(\tau)(\cdot, u_0) \\ &= \sum_{|\kappa| \leq p-\varrho(\tau)} \frac{1}{\kappa!} D^\kappa(\Psi(\tau))(0, u_0) \left(I(\tau) - \widehat{I}(\tau) \right) \mathbf{z}^\kappa + \mathcal{O}(h^{p-\varrho(\tau)+1}) \\ &= \sum_{|\kappa| \leq p-\varrho(\tau)} \frac{1}{\kappa!} D^\kappa(\Psi(\tau))(0, u_0) \left(I(\tau^{(\kappa)})1 - \widehat{I}(\tau^{(\kappa)})1 \right) + \mathcal{O}(h^{p-\varrho(\tau)+1}). \end{aligned}$$

Here we used $\|D^\kappa \Psi(\tau)\| = \mathcal{O}(h^{|\kappa|})$ to bound the remainder term. Since the Runge–Kutta method is of order p , the claim now follows from $\varrho(\tau^{(\kappa)}) = \varrho(\tau) + |\kappa| \leq p$ and Corollary 4.4 which implies $I(\tau^{(\kappa)})1 = \widehat{I}(\tau^{(\kappa)})1$. \square

This result now allows us to prove an error bound for Lawson methods which is uniform for all problems (5) with A satisfying Assumption 3.1.

Theorem 4.8 *Let u be the solution of (5), let Assumption 3.1 and the assumptions of Theorem 4.7 be satisfied. If the underlying Runge–Kutta method is of (classical) order p , then there exists $h_0 > 0$ such that for all $0 < h \leq h_0$ sufficiently small,*

$$\|u(t_n) - u_n\| \leq Ch^p, \quad t_n = nh \leq T,$$

where C and h_0 are independent of n , h , and A .

Proof In the case of a group, we define an equivalent norm by

$$\|v\|_{\star} = \sup_{t \in \mathbb{R}} \|e^{-tA}v\|. \quad (18)$$

Then, e^{-tA} is a group of contractions in the corresponding operator norm

$$\|e^{-tA}\|_{\star} \leq 1 \quad \text{for all } t \in \mathbb{R}. \quad (19)$$

If $-A$ only generates a bounded semigroup, we take the supremum in (18) over $t \geq 0$ only. This shows that $-A$ generates a semigroup of contractions in this equivalent norm.

By assumption, g is locally Lipschitz continuous. Then (19) and Theorem 3.10 show that the Lawson method is locally Lipschitz with respect to the initial value with a Lipschitz constant of size $1 + \mathcal{O}(h)$. This implies the required stability.

The error bound follows in a standard way using Lady Windermere's fan; see [5, Fig. I.7.1]. \square

5 Regularity conditions and applications

It remains to discuss the regularity conditions (17) and to give some applications. We first examine the conditions for orders one and two, respectively. The extension to higher orders is a tedious but straightforward exercise. It turns out that these regularity conditions can all be expressed in terms of commutators, very much like in the case of splitting methods.

In order to obtain simple sufficient conditions, we replace the space $C^{k,1}(\Omega, X)$ in condition (17) by the subspace of $k + 1$ times partially differentiable functions with uniformly bounded partial derivatives on Ω in the following discussion. This is also justified by the fact that Lipschitz continuous functions are almost everywhere differentiable (Rademacher's theorem).

5.1 Condition for order one

Since $p = \varrho(\tau) = 1$, we only have to consider the tree $\tau = \bullet$ in (17). Differentiating

$$\Psi(\bullet)(\lambda, w) = e^{-(1-\lambda)hA} g_{\lambda}(w) = e^{-(1-\lambda)hA} g(e^{-\lambda hA} w) \quad (20)$$

with respect to λ yields

$$\begin{aligned} \partial_{\lambda} \Psi(\bullet)(\lambda, w) &= h e^{-(1-\lambda)hA} \left(A g_{\lambda}(w) - g'_{\lambda}(w) A e^{-\lambda hA} w \right) \\ &= h e^{-(1-\lambda)hA} [F_A, g](e^{-\lambda hA} w), \end{aligned} \quad (21)$$

where $[F_A, g]$ denotes the Lie commutator of g and $F_A(w) = Aw$, defined as

$$[F_A, g](w) = F'_A(w)g(w) - g'(w)F_A(w) = Ag(w) - g'(w)Aw. \quad (22)$$

From this calculation, we conclude the following result. If the bound

$$\sup_{0 \leq \lambda \leq 1} \sup_{0 \leq t \leq T} \left\| e^{-(1-\lambda)hA} [F_A, g](e^{-\lambda hA} u(t)) \right\| \leq C \quad (23)$$

holds with a constant C that is allowed to depend on $C_{\mathcal{F}}$, then a Lawson method of non-stiff order *one* has also *stiff* order *one*.

5.2 Conditions for order two

Stiff order two is achieved if we require the following two regularity conditions

$$\Psi(\bullet)(\cdot, u(t)) \in C^2([0, 1], X) \quad \text{and} \quad \Psi(\bullet\bullet)(\cdot, u(t)) \in C^1([0, 1]^2, X).$$

Here, we omit the numbering of the tree with two nodes because it is unique. We commence with the first condition and exploit the fact that $\partial_\lambda \Psi(\bullet)(\lambda, w)$ is of exactly the same form as (20) with g replaced by the vector field $[F_A, g]$. Hence from (21) we have

$$\begin{aligned} \partial_\lambda^2 \Psi(\bullet)(\lambda, w) & \frac{d}{dw} ([F_A, g])(e^{-\lambda hA} w) A e^{-\lambda hA} w \\ & = h^2 e^{-(1-\lambda)hA} [F_A, [F_A, g]](e^{-\lambda hA} w). \end{aligned}$$

Therefore, the bound

$$\sup_{0 \leq \lambda \leq 1} \sup_{0 \leq t \leq T} \left\| e^{-(1-\lambda)hA} [F_A, [F_A, g]](e^{-\lambda hA} u(t)) \right\| \leq C \quad (24)$$

should hold with a constant C that is independent of $\|A\|$.

Next, we move to the second condition. Differentiating

$$\Psi(\bullet\bullet)(\lambda_1, \lambda_2, w) = e^{-(1-\lambda_1)hA} g'_{\lambda_1}(w) e^{-(\lambda_1-\lambda_2)hA} g_{\lambda_2}(w)$$

with respect to λ_1 and λ_2 yields

$$\begin{aligned} \partial_{\lambda_1} \Psi(\bullet\bullet)(\lambda_1, \lambda_2, w) & = h e^{-(1-\lambda_1)hA} \left(A g'_{\lambda_1}(w) e^{-(\lambda_1-\lambda_2)hA} g_{\lambda_2}(w) \right. \\ & \quad - g''_{\lambda_1}(w) (e^{-\lambda_1 hA} A w, e^{-(\lambda_1-\lambda_2)hA} g_{\lambda_2}(w)) \\ & \quad \left. - g'_{\lambda_1}(w) A e^{-(\lambda_1-\lambda_2)hA} g_{\lambda_2}(w) \right) \\ & = h e^{-(1-\lambda_1)hA} [F_A, g]'(e^{-\lambda_1 hA} w) e^{-(\lambda_1-\lambda_2)hA} g_{\lambda_2}(w), \end{aligned}$$

since by definition (22) the derivative of the commutator satisfies

$$[F_A, g]'(w)v = \frac{d}{dw} ([F_A, g])(w)v = A g'(w)v - g''(w)(Aw, v) - g'(w)Av. \quad (25)$$

commutator of two operators, and Moreover, we have

$$\partial_{\lambda_2} \Psi(\bullet\bullet)(\lambda_1, \lambda_2, w) = h e^{-(1-\lambda_1)hA} g'_{\lambda_1}(w) e^{-(\lambda_1-\lambda_2)hA} [F_A, g](e^{-\lambda_2 hA} w),$$

respectively. From these two relations, we infer that the bounds

$$\sup_{0 \leq \lambda_2 \leq \lambda_1 \leq 1} \sup_{0 \leq t \leq T} \left\| e^{-(1-\lambda_1)hA} [F_A, g]'(e^{-\lambda_1 hA} u(t)) e^{-(\lambda_1 - \lambda_2)hA} g_{\lambda_2}(u(t)) \right\| \leq C, \quad (26a)$$

$$\sup_{0 \leq \lambda_2 \leq \lambda_1 \leq 1} \sup_{0 \leq t \leq T} \left\| e^{-(1-\lambda_1)hA} g'_{\lambda_1}(u(t)) e^{-(\lambda_1 - \lambda_2)hA} [F_A, g](e^{-\lambda_2 hA} u(t)) \right\| \leq C \quad (26b)$$

should hold with a constant C that is independent of $\|A\|$.

From the above calculations, we conclude the following result. If the conditions (23), (24), and (26) hold with a constant C that does not depend on $\|A\|$, then a Lawson method of non-stiff order *two* has also *stiff* order *two*.

5.3 Conditions for higher order

The following lemma provides the formulas to derive the order conditions for orders larger than two in a systematic way.

Lemma 5.1 *Let $m \geq 1$.*

(a) *For $\tau = \bullet$ we have*

$$\partial_\lambda^m \Psi_\zeta(\bullet)(\lambda, w) = h^m e^{-(\zeta - \lambda)hA} [F_A, g]_m(e^{-\lambda hA} w),$$

where $[F_A, g]_{m+1} = [F_A, [F_A, g]_m]$ with $[F_A, g]_1 = [F_A, g]$ denotes the $(m+1)$ -fold commutator.

(b) *For $\tau = [\tau_1, \dots, \tau_k] \in \mathcal{LT}$ we have*

$$\begin{aligned} & \partial_\lambda^m \Psi_\zeta(\tau)(\lambda, \lambda_2, \dots, \lambda_{\rho(\tau)}, w) \\ &= h e^{-(\zeta - \lambda)hA} [F_A, g]_m^{(k)}(e^{-\lambda hA} w) (\Psi_\lambda(\tau_1)(\cdot_1, w), \dots, \Psi_\lambda(\tau_k)(\cdot_k, w)) \end{aligned}$$

for $\lambda \in \mathbb{R}$, and where \cdot_j are the $\varrho(\tau_j)$ variables corresponding to the subtree τ_j , $j = 1, \dots, k$, sorted in increasing order.

Proof Both parts are proved by induction on m .

(a) For $m = 1$ the statement was proved in (21). The induction step is proved by the same arguments as were used for $m = 2$ above.

(b) To prove the statement for $m = 1$, we first note that for $\tau = [\tau_1, \dots, \tau_k]$ the integrand of $G_\zeta(\tau)$ is given recursively as

$$\Psi_\zeta(\tau)(\lambda, \lambda_2, \dots, \lambda_{\rho(\tau)}, w) = e^{-(\zeta - \lambda)hA} g_\lambda^{(k)}(w) (\Psi_\lambda(\tau_1)(\cdot_1, w), \dots, \Psi_\lambda(\tau_k)(\cdot_k, w)).$$

Since $\partial_\eta \Psi_\eta(\tau) = -hA\Psi_\eta(\tau)$ for any tree τ , we obtain

$$\begin{aligned} & \partial_\lambda \Psi_\zeta(\tau)(\lambda, \lambda_2, \dots, \lambda_{\rho(\tau)}, w) \\ &= h e^{-(\zeta-\lambda)hA} \left(Ag_\lambda^{(k)}(w)(\Psi_\lambda(\tau_1)(\cdot_1, w), \dots, \Psi_\lambda(\tau_k)(\cdot_k, w)) \right. \\ & \quad - g_\lambda^{(k+1)}(w)(Ae^{-\lambda hA}w, \Psi_\lambda(\tau_1)(\cdot_1, w), \dots, \Psi_\lambda(\tau_k)(\cdot_k, w)) \\ & \quad - g_\lambda^{(k)}(w)(A\Psi_\lambda(\tau_1)(\cdot_1, w), \Psi_\lambda(\tau_2)(\cdot_2, w), \dots, \Psi_\lambda(\tau_k)(\cdot_k, w)) \\ & \quad - \dots \\ & \quad \left. - g_\lambda^{(k)}(w)(\Psi_\lambda(\tau_1)(\cdot_1, w), \dots, \Psi_\lambda(\tau_{k-1})(\cdot_{k-1}, w), A\Psi_\lambda(\tau_k)(\cdot_k, w)) \right). \end{aligned}$$

On the other hand, by definition (22), we have

$$[F_A, g]^{(k)}(w)(v_1, \dots, v_k) = Ag^{(k)}(w)(v_1, \dots, v_k) - \frac{d^k}{dw^k}(g'(w)Aw)(v_1, \dots, v_k).$$

Using induction on k it is easy to see that

$$\begin{aligned} \frac{d^k}{dw^k}(g'(w)Aw)(v_1, \dots, v_k) &= g^{(k+1)}(w)(Aw, v_1, \dots, v_k) \\ & \quad + g^{(k)}(w)(Av_1, v_2, \dots, v_k) + \dots \\ & \quad + g^{(k)}(w)(v_1, \dots, Av_k). \end{aligned}$$

This proves the claim for $m = 1$. If it holds for some $m \geq 1$ then it does also for $m + 1$, since the same calculation can be done with $[F_A, g]^{(k)}$ in the role of $g^{(k)}$. \square

The lemma thus shows that all derivatives arising in the order conditions can be obtained recursively from the tree structure. Moreover, only commutators, iterated commutators and their derivatives appear.

5.4 Specialization to linear problems

For the linear evolution equation

$$u' + Au = Bu, \quad u(0) = u_0$$

with bounded operator B on X , the above conditions (23), (24), and (26) simplify a bit. Having $g(u) = Bu$, the Lie commutator coincides with the operator commutator of A and B

$$[F_A, g](w) = ABw - BA w = [A, B]w.$$

A first-order Lawson method is of *stiff* order one if

$$\sup_{0 \leq \lambda \leq 1} \sup_{0 \leq t \leq T} \left\| e^{-(1-\lambda)hA} [A, B] e^{-\lambda hA} u(t) \right\| \leq C. \quad (27a)$$

For second order, the conditions read

$$\sup_{0 \leq \lambda \leq 1} \sup_{0 \leq t \leq T} \left\| e^{-(1-\lambda)hA} [A, [A, B]] e^{-\lambda hA} u(t) \right\| \leq C, \quad (27b)$$

$$\sup_{0 \leq \lambda_2 \leq \lambda_1 \leq 1} \sup_{0 \leq t \leq T} \left\| e^{-(1-\lambda_1)hA} [A, B] e^{-(\lambda_1-\lambda_2)hA} B e^{-\lambda_2 hA} u(t) \right\| \leq C, \quad (27c)$$

$$\sup_{0 \leq \lambda_2 \leq \lambda_1 \leq 1} \sup_{0 \leq t \leq T} \left\| e^{-(1-\lambda_1)hA} B e^{-(\lambda_1-\lambda_2)hA} [A, B] e^{-\lambda_2 hA} u(t) \right\| \leq C. \quad (27d)$$

We recall that such conditions also arise in the analysis of splitting methods; see [10].

Using Lemma 5.1, the above analysis can easily be generalized to higher order, since for linear problems, only long trees have to be considered. For all other trees, which have at least one node with two branches, the integrand Ψ vanishes.

5.5 Nonlinear Schrödinger equations

For the time discretization of nonlinear Schrödinger equations

$$u' = i(\Delta u + f(|u|^2)u), \quad (28)$$

split-step methods are commonly viewed as the method of choice. In recent years, however, exponential integrators have been considered as a viable alternative for the solution of (28). For instance, [2] studied exponential integrators in the context of Bose–Einstein condensates; [3] and [1] reported favorable results for Lawson integrators of the form as discussed in this paper. Rigorous convergence results, however, are still missing for these methods.

As an application of our analysis, we will use the above regularity conditions (23), (24), and (26) to verify second-order convergence of Lawson methods. We refrain from any particular space discretization and argue in an abstract Hilbert space framework. Note, however, that our reasoning carries over to spatial discretizations (by spectral methods, e.g.) without any difficulty.

For this purpose, we consider (28) with periodic boundary conditions on the d dimensional torus and smooth potential. Then it is well known (see, e.g., [12, Thm. 4.1]) that the problem is well posed in H^m for $m > d/2$. The regularity of an initial value $u_0 \in H^m$ is thus preserved along the solution. Henceforth we choose $m > d/2$.

Second-order Strang splitting for (28) with $f(u) = \pm u$ was rigorously analysed in [15]. There it was shown that commutator relations similar to our conditions (23), (24), and (26) play a crucial role in the convergence proof for Strang splitting. The analysis given here shows that Lawson methods converge under the same regularity assumptions as splitting schemes. This will be worked out now in detail for first and second-order methods.

Let $A = -i\Delta$ and $g(u) = i\beta|u|^2u$, $\beta \in \mathbb{R}$, i.e. $f = \beta I$. By

$$g(u+w) = i\beta(u+w)^2(\overline{u+w}) = g(u) + i\beta(u^2\bar{w} + 2u\bar{w}u) + \mathcal{O}(|w|^2),$$

the Fréchet derivative of g is given by

$$g'(u)w = i\beta(u^2\bar{w} + 2|u|^2 w).$$

The first commutator $[F_A, g]$ then takes the form

$$\begin{aligned} [F_A, g](u) &= Ag(u) - g'(u)Au \\ &= -i\Delta g(u) + g'(u)i\Delta u \\ &= \beta\nabla \cdot \nabla(u^2\bar{u}) + i\beta(u^2(i\Delta u) + 2|u|^2 i\Delta u) \\ &= \beta\nabla \cdot (2u\bar{u}\nabla u + u^2\nabla\bar{u}) + \beta(u^2\Delta\bar{u} - 2|u|^2\Delta u) \\ &= \beta(2\bar{u}\nabla u \cdot \nabla u + 2u\nabla u \cdot \nabla\bar{u} + 2|u|^2\Delta u + 2u\nabla u \cdot \nabla\bar{u} \\ &\quad + 2u^2\Delta\bar{u} - 2|u|^2\Delta u) \\ &= 2\beta(\bar{u}\nabla u \cdot \nabla u + 2u\nabla u \cdot \nabla\bar{u} + u^2\Delta\bar{u}). \end{aligned} \quad (29)$$

We next show that the commutator can be bounded in H^m if the solution is in H^{m+2} for $m \geq 0$.

Lemma 5.2 *Let $\Omega \subset \mathbb{R}^d$, $d \leq 3$, be a bounded Lipschitz domain. Then there exists a constant C which only depends on Ω and d such that*

$$\|[F_A, g]u\|_m \leq C \|u\|_{m+2}^3. \quad (30)$$

Proof Note that by the Sobolev embedding theorem we have the following bounds

$$\|uvw\|_0 \leq C \|u\|_1 \|v\|_1 \|w\|_1, \quad (31a)$$

$$\|uvw\|_0 \leq C \|u\|_0 \|v\|_2 \|w\|_2, \quad (31b)$$

$$\|uvw\|_1 \leq C \|u\|_1 \|v\|_2 \|w\|_2, \quad (31c)$$

$$\|uvw\|_m \leq C \|u\|_m \|v\|_m \|w\|_m, \quad m \geq 2; \quad (31d)$$

see [15, Section 8].

For $m = 0$, the bound (30) follows from using (31a) for the first two terms and (31b) for the last one in the explicit expression (29) of $[F_A, g]$. For $m = 1$ we apply (31c) to all terms and for $m \geq 2$ the bound follows from (31d). \square

For Lawson methods, a first-order convergence bound in H^m thus requires H^{m+2} regularity of the exact solution, which is the same regularity as required for the first-order Lie splitting.

For second-order methods, one has to estimate the double commutator $[F_A, [F_A, g]]$. A simple calculation shows that a bound in H^m requires H^{m+4} regularity of the exact solution. This situation is exactly the same as for second-order Strang splitting (see [15]). Using (29) we conclude that the derivative of the commutator $[F_A, g]$ can be expressed as

$$\begin{aligned} [F_A, g]'(u)w &= 2\beta(\bar{w}\nabla u \cdot \nabla u + 2\bar{u}\nabla u \cdot \nabla w + 2w\nabla u \cdot \nabla\bar{u} + 2u\nabla w \cdot \nabla\bar{u} \\ &\quad + 2u\nabla u \cdot \nabla\bar{w} + u^2\Delta\bar{w} + 2uw\Delta\bar{u}). \end{aligned}$$

This commutator can again be bounded in H^m for $u, w \in H^{m+2}$. We thus conclude that Lawson methods require the same regularity for second-order convergence as Strang splitting.

5.6 Numerical examples

Lawson methods exhibit a strong order reduction, in general. For particular problems, however, they show full order of convergence (see [11], [2], [3], [1], and [16]). Most of the problems considered in these papers result from space discretizations of partial differential equations posed with periodic boundary conditions.

After space discretization (by finite differences, finite elements, or spectral methods) the evolution equation (5) becomes an ordinary differential equation

$$\mathbf{u}'(t) + A_N \mathbf{u}(t) = g_N(\mathbf{u}(t)), \quad \mathbf{u}(0) = \mathbf{u}_0 \quad (32)$$

with a matrix $A_N \in \mathbb{C}^{N \times N}$ and a discretization $g_N : \mathbb{C}^N \rightarrow \mathbb{C}^N$ of g , where N denotes the employed degrees of freedom. In order to satisfy Assumption 3.1 the space discretization is required to provide matrices A_N such that

$$\|e^{-tA_N}\| \leq C_{\mathcal{F}} \quad (33)$$

holds with a constant $C_{\mathcal{F}}$ being uniform in N and $t \in \mathbb{R}$.

In the previous sections we showed that full order of convergence is only guaranteed if certain regularity conditions are satisfied. The aim of the following numerical examples is to show that order reduction can also be observed numerically, if some of these regularity assumptions are violated. In fact, such order reductions can even be observed for linear problems. Hence we abstain from presenting numerical examples for semilinear problems here. Numerous such examples can be found in the literature mentioned above. We also restrict ourselves to the first order schemes covered by our analysis, the exponential Euler and the Lawson–Euler method, since they already show interesting (and different) convergence behavior.

We consider the linear Schrödinger equation

$$u_t = iu_{xx} + if(x)u, \quad x \in [-\pi, \pi], \quad u(0, \cdot) = u_0, \quad (34)$$

with periodic boundary conditions and discretize it using a Fourier spectral method on an equidistant grid. Let N be even and denote by \mathcal{F}_N the discrete Fourier matrix. Then matrix A_N is given as

$$A_N = i\mathcal{F}_N^{-1} D_N^2 \mathcal{F}_N, \quad \text{where} \quad D_N = \text{diag}\left(-\frac{N}{2} + 1, -\frac{N}{2} + 2, \dots, \frac{N}{2}\right),$$

and

$$g_N(\mathbf{u}) = B_N \mathbf{u}, \quad B_N = i \text{diag}\left(f(x_{-N/2+1}), \dots, f(x_{N/2})\right), \quad x_m = m \frac{2\pi}{N}.$$

With this notation, the exact solution of (32) is given by

$$\mathbf{u}(t) = e^{t(-A_N + B_N)} \mathbf{u}_0. \quad (35)$$

Example 5.3 The aim of this example is to explain that the concept of regularity is relevant even in the ODE context. In order to show what regularity means here, we carry out the following experiment. For each $N = 2^7, \dots, 2^{12}$ we choose a regularity parameter $\alpha \geq 0$ and a vector $\mathbf{r} = (r_m)_{m=-N/2+1}^{N/2} \in \mathbb{C}^N$ of Fourier coefficients whose entries contain random numbers uniformly distributed in the unit disc. Then we define an initial function as the trigonometric polynomial

$$\tilde{u}_{0;N}(x) = \sum_{m=-N/2+1}^{N/2} \nu_m e^{imx}, \quad \nu_m = \frac{r_m}{(1+m^2)^{\frac{1}{2}} \left(\frac{1}{2} + \alpha + \epsilon\right)}, \quad \epsilon = 10^{-6}.$$

In the limit $N \rightarrow \infty$, this sequence of trigonometric polynomials converges to a function in the Sobolev space $H_{\text{per}}^\alpha = H_{\text{per}}^\alpha((-\pi, \pi))$ equipped with the norm

$$\|u\|_\alpha^2 = 2\pi \sum_{m \in \mathbb{Z}} (1+m^2)^\alpha |\hat{u}_m|^2 \quad \text{for} \quad u = \sum_{m \in \mathbb{Z}} \hat{u}_m e^{imx}.$$

For $\alpha = 0$ we have the standard L^2 norm

$$\|u\|_0^2 = \int_{-\pi}^{\pi} |u(x)|^2 dx.$$

Then we define an initial vector $\mathbf{u}_0 \in \mathbb{C}^N$ for (32) corresponding to a function $u_0 \in H_{\text{per}}^\alpha$ by setting the j th component as

$$(\mathbf{u}_0)_j = u_{0;N}(x_j), \quad j = -\frac{N}{2} + 1, \dots, \frac{N}{2}, \quad (36)$$

where $u_{0;N} = \tilde{u}_{0;N} / \|\tilde{u}_{0;N}\|_0$ has unit L^2 norm. The discrete Sobolev norms in \mathbb{C}^N corresponding to $\|\cdot\|_\alpha$ can be computed via

$$\|\mathbf{u}\|_{\alpha,N}^2 = 2\pi \left\| (I + D_N^2)^{\alpha/2} \mathcal{F}_N \mathbf{u} \right\|_{\mathbb{C}^N}^2,$$

where $\|\cdot\|_{\mathbb{C}^N}$ denotes the Euclidean norm in \mathbb{C}^N . This yields $\|\mathbf{u}_0\|_{\alpha,N} = \|u_{0;N}\|_\alpha$.

In Figure 1 we plot $\|\mathbf{u}_0\|_{\mu,N}$ for different values of μ against the number of Fourier modes N . The three graphs clearly show that $\|\mathbf{u}_0\|_{\mu,N}$ is bounded independently of the number N of Fourier modes only for $\mu \leq \alpha$. This corresponds to the continuous case, where obviously, the Sobolev norm $\|u\|_\mu$ is bounded for all functions $u \in H_{\text{per}}^\alpha$ for $\mu \leq \alpha$.

The example clearly shows that regularity of the corresponding continuous function is crucial to obtain error bounds which do not deteriorate in the limit $N \rightarrow \infty$.

After these introductory explanations, we now fix the spatial discretization and set $N = 2048$. We consider (34) for two different functions f :

$$f(x) = \sin x, \quad (37a)$$

$$f(x) = (x/\pi)^2. \quad (37b)$$




Fig. 1 Illustration of discrete regularity: the discrete H_{per}^μ -Sobolev seminorm $\|\mathbf{u}_0\|_{\mu,N}$ is plotted against the number N of Fourier modes, where \mathbf{u}_0 is chosen as in (36) (and thus corresponds to a function in H_{per}^α).

Example 5.4 In Figure 2 we show the numerically observed orders of the exponential Euler and the Lawson–Euler method for the smooth, periodic potential (37a) for different values of α such that the corresponding initial function is contained in H_{per}^α . The leading error terms of the new analysis for the exponential Euler and the Lawson–Euler method are given in (14) and (27a), respectively.

Since $B : H_{\text{per}}^\alpha \rightarrow H_{\text{per}}^\alpha$ is a bounded perturbation of A , the exact solution of the continuous problem is guaranteed to stay in H_{per}^α for initial values in H_{per}^α for $\alpha \geq 0$. For the discrete problem, $e^{-\lambda h A_N}$ and $e^{\lambda h (-A_n + B_N)}$ are unitary matrices, which means that they leave all discrete Sobolev norms $\|\cdot\|_{\alpha,N}$ invariant. Thus the expression in (27a) can be bounded by

$$\begin{aligned} \left\| e^{-(1-\lambda)hA_N} [A_N, B_N] e^{-\lambda h A_N} \mathbf{u}(t) \right\|_{0,N} &\leq c_1 \left\| e^{-\lambda h A_N} \mathbf{u}(t) \right\|_{1,N} = c_1 \|\mathbf{u}(t)\|_{1,N} \\ &= c_1 \|\mathbf{u}_0\|_{1,N}. \end{aligned}$$

Here, the first inequality was proved in [10, Lemma 3.1] with a constant c_1 independent of N and A_N .

Hence, the (sufficient but not necessary) order condition (27a) for the Lawson–Euler method yields order one convergence for initial values bounded in $\|\cdot\|_\alpha$ for $\alpha \geq 1$. Numerically, we observe an order reduction for $\alpha = 0$ for the Lawson–Euler method, while the exponential Euler method, which requires initial values in $H_{\text{per}}^2 = D(A)$, cf. (14) or (15), shows order reduction for $\alpha \leq 1$. For $\alpha = 0$ the error of the exponential Euler method has an irregular behaviour for larger step sizes. To better visualise the order, we added thin




Fig. 2 Discrete $L^\infty((0, 1), L^2(\Omega))$ error (on the y -axis) of the numerical solution of (34) with periodic potential (37a) plotted against the step size h (on the x -axis) for starting values in H_{per}^α . The top graph shows the exponential Euler method and the bottom graph shows the Lawson–Euler method. The values of p in the legend show the numerically observed orders of the schemes.

lines (blue in the colored version) to all curves related to $\alpha = 0$. The slopes p of these lines are also given in the legends (blue in the colored version).

Example 5.5 In Figure 3 we present the same experiment for the quadratic potential (37b). Here, the commutator bound of [10, Lemma 3.1] does not apply, since it requires a C^5 smooth and periodic potential f . The situation differs considerably for the exponential Euler method which suffers from order reduction for all $\alpha \leq 2$ due to the nonsmooth potential f . In contrast, the Lawson–Euler method still converges with order one for $\alpha \geq 0.5$.

Note that for these examples, the convergence behavior is slightly better than predicted by our theory. This is not a contradiction, because the order conditions are only sufficient but not necessary. To be more precise, our analysis contains a worst case estimation of the error propagation from the local to the global error by using Lady Windermere’s fan in the proof of Theorem 4.8. Nevertheless, the examples clearly show the different behavior of the exponential Euler method and the Lawson–Euler method. Which of the two methods yields better results depends on the given problem, as reflected by our error analysis.




Fig. 3 Discrete $L^\infty((0, 1), L^2(\Omega))$ error (on the y -axis) of the numerical solution of (34) with quadratic potential (37b) plotted against the step size h (on the x -axis) for starting values in H_{per}^α . The top graph shows the exponential Euler method and the bottom graph shows the Lawson–Euler method. The values of p in the legend show the numerically observed orders of the schemes.

Acknowledgements The work of M. Hochbruck and J. Leibold was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Project-ID 258734477 SFB 1173.

References

1. Balac, S., Fernandez, A., Mahé, F., Méhats, F., Texier-Picard, R.: The interaction picture method for solving the generalized nonlinear Schrödinger equation in optics. *ESAIM: M2AN* **50**(4), 945–964 (2016). DOI 10.1051/m2an/2015060
2. Besse, C., Dujardin, G., Lacroix-Violet, I.: High order exponential integrators for nonlinear Schrödinger equations with application to rotating Bose–Einstein condensates. *SIAM J. Numer. Anal.* **55**(3), 1387–1411 (2017). DOI 10.1137/15M1029047
3. Cano, B., González-Pachón, A.: Projected explicit Lawson methods for the integration of Schrödinger equation. *Numer. Methods Partial Differential Equations* **31**(1), 78–104 (2015). DOI 10.1002/num.21895
4. Hairer, E., Lubich, C., Wanner, G.: Geometric numerical integration: Structure-preserving algorithms for ordinary differential equations, *Springer Series in Computational Mathematics*, vol. 31, 2nd ed., Springer-Verlag, Berlin (2006)
5. Hairer, E., Nørsett, S.P., Wanner, G.: Solving ordinary differential equations I: Nonstiff problems, *Springer Series in Computational Mathematics*, vol. 8, 2nd ed., Springer-Verlag, Berlin (1993)
6. Hansen, E., Ostermann, A.: High-order splitting schemes for semilinear evolution equations. *BIT* **56**(4), 1303–1316 (2016). DOI 10.1007/s10543-016-0604-2
7. Hochbruck, M., Ostermann, A.: Explicit exponential Runge–Kutta methods for semilinear parabolic problems. *SIAM J. Numer. Anal.* **43**(3), 1069–1090 (2005). DOI 10.1137/040611434

8. Hochbruck, M., Ostermann, A.: Exponential Runge–Kutta methods for parabolic problems. *Appl. Numer. Math.* **53**(2–4), 323–339 (2005)
9. Hochbruck, M., Ostermann, A.: Exponential integrators. *Acta Numer.* **19**, 209–286 (2010). DOI 10.1017/S0962492910000048
10. Jahnke, T., Lubich, C.: Error bounds for exponential operator splittings. *BIT* **40**(4), 735–744 (2000). DOI 10.1023/A:1022396519656
11. Kassam, A.K., Trefethen, L.N.: Fourth-order time-stepping for stiff PDEs. *SIAM J. Sci. Comput.* **26**(4), 1214–1233 (electronic) (2005). DOI 10.1137/S1064827502410633
12. Kato, T.: On nonlinear Schrödinger equations. II. H^s -solutions and unconditional well-posedness. *J. Anal. Math.* **67**, 281–306 (1995). DOI 10.1007/BF02787794
13. Lawson, J.D.: Generalized Runge–Kutta processes for stable systems with large Lipschitz constants. *SIAM J. Numer. Anal.* **4**(3), 372–380 (1967). DOI 10.1137/0704033
14. Luan, V.T., Ostermann, A.: Exponential B-series: the stiff case. *SIAM J. Numer. Anal.* **51**(6), 3431–3445 (2013). DOI 10.1137/130920204
15. Lubich, C.: On splitting methods for Schrödinger–Poisson and cubic nonlinear Schrödinger equations. *Math. Comp.* **77**(264), 2141–2153 (2008). DOI 10.1090/S0025-5718-08-02101-7
16. Montanelli, H., Bootland, N.: Solving stiff PDEs in 1D, 2D and 3D with exponential integrators. Preprint arXiv:1604.08900 (2016). URL <https://arxiv.org/abs/1604.08900>