

# Full discretization error analysis of exponential integrators for semilinear wave equations

Benjamin Dörich, Jan Leibold

CRC Preprint 2021/31, July 2021

KARLSRUHE INSTITUTE OF TECHNOLOGY

CRC 1173



Wave  
phenomena

## Participating universities



Universität Stuttgart

EBERHARD KARLS  
UNIVERSITÄT  
TÜBINGEN



Funded by

**DFG**

# FULL DISCRETIZATION ERROR ANALYSIS OF EXPONENTIAL INTEGRATORS FOR SEMILINEAR WAVE EQUATIONS

BENJAMIN DÖRICH AND JAN LEIBOLD

ABSTRACT. In this article we prove full discretization error bounds for semilinear second-order evolution equations. We consider exponential integrators in time applied to an abstract nonconforming semi discretization in space. Since the fully discrete schemes involve the spatially discretized semigroup, a crucial point in the error analysis is to eliminate the continuous semigroup in the representation of the exact solution. Hence, we derive a modified variation-of-constants formula driven by the spatially discretized semigroup which holds up to a discretization error. Our main results provide bounds for the full discretization errors for exponential Adams and explicit exponential Runge–Kutta methods. We show convergence with the stiff order of the corresponding exponential integrator in time, and errors stemming from the spatial discretization.

As an application of the abstract theory, we consider an acoustic wave equation with kinetic boundary conditions, for which we also present some numerical experiments to illustrate our results.

## 1. INTRODUCTION

In the present paper we analyze the full discretization of semilinear second-order evolution equations

$$(1.1) \quad \begin{aligned} u''(t) + Bu'(t) + Au(t) &= f(t, u(t)), \quad t \in (0, T], \\ u(0) &= u^0, \quad u'(0) = v^0, \end{aligned}$$

posed on a Hilbert space  $H$  with unbounded operators  $A$  and  $B$  and a smooth nonlinearity  $f$ . The numerical scheme is obtained by applying exponential Runge–Kutta and multistep methods to the first-order formulation of a spatially discretized version of (1.1).

Equation (1.1) covers a wide range of second-order semilinear wave equations. The most prominent example is the acoustic wave equation subject to homogeneous Dirichlet or Neumann boundary conditions. A more interesting example, however, is the acoustic wave equation with kinetic boundary conditions. This model was derived in [14] for a membrane with a boundary that carries a mass density and is subject to linear tension. We also refer to [33, 34], where analytical wellposedness of such equations was shown and further references are given.

---

2020 *Mathematics Subject Classification.* Primary: 65M12, 65M15; Secondary: 65M60, 65J08.

*Key words and phrases.* error analysis, full discretization, exponential integrators, wave equation, semilinear evolution equations, dynamic boundary conditions, nonconforming space discretization, a-priori error bounds.

Funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 258734477 – SFB 1173.

Kinetic boundary conditions are a special type of dynamic boundary conditions, given by a differential equation on the boundary and usually posed on smooth boundaries. In practice, such boundaries have to be approximated by piecewise polynomials, which leads to nonconforming space discretizations. This makes the error analysis much more involved, since the exact and the numerical solution are defined on different spatial domains. To tackle this difficulty, the so-called *unified error analysis* was introduced in [15] for linear wave-type equations, and extended in [16] to the semilinear case. The unified error analysis is a systematic approach, where an abstract space discretization of (1.1) is considered as an evolution equation on a finite dimensional space. In [15, 16], abstract error bounds were derived, which lead to optimal convergence rates for a large class of equations and corresponding conforming and nonconforming space discretizations.

In this paper, we combine the unified error analysis with exponential integrators. A review on those schemes can, e.g., be found in [19]. Compared to classical numerical integrators, where the fundamental theorem of calculus is used as the starting point for approximations, exponential integrators are based on the variation-of-constants formula. This way, the representation of the solution incorporates the unbounded linear part via semigroup theory, while only the smooth nonlinear part within the integral is approximated.

We focus on two different classes of methods. On the one hand, we consider exponential Adams methods. Here, the nonlinearity is replaced by an interpolation polynomial through preceding approximations and the polynomial is integrated exactly. Such methods were first proposed in [29] and a rigorous error analysis for the time discretization was performed in [7] and [20]. On the other hand, we employ explicit exponential Runge–Kutta methods, first considered in [11, 12, 24]. They extend the classical Runge–Kutta methods by the use of operator-valued coefficients which depend on the linear part of the problem. For the methods considered in this work, error bounds for the spatially continuous case were derived in [18]. In [10], related schemes have been applied to stiff kinetic equations. For the (semi-)linear Schrödinger equation, symmetric but implicit variants of these schemes and their geometric properties were derived in [8].

For a long time, these methods have been regarded as unpractical due to evaluation of a matrix exponential and related matrix functions. However, this view changed when in [17] for the first time an implementation was proposed that was competitive in certain scenarios. For quasilinear Maxwell’s equations, the numerical experiments in [31] confirm this computational potential. Further, their efficiency was demonstrated in [26, 28] in the application to molecular dynamics and nonlinear coupled oscillators. Besides the Krylov methods to compute a matrix function applied to a vector, a different approach is considered in [22]. Here, the analytic functions are expressed by the Cauchy integral formula which is discretized such that it remains to evaluate a fixed number of resolvents. Very recently, in [6] novel rational exponential integrators were derived which gain efficiency and accuracy by a parallel-in-time computation of resolvents applied to a vector in order to evaluate the matrix exponential.

So far, the error analysis has mostly been performed for abstract evolution equations or for systems of ordinary differential equations. Concerning the full discretization error analysis involving exponential integrators, we are only aware of the following works. In [5, 9], stochastic parabolic partial differential equations are

discretized by the exponential Euler method combined with a spectral method in space. This type of space discretization has very favorable properties in the analysis, however, one needs to know the relevant eigenvalues and eigenvectors of the differential operators exactly. This is for example ensured if one considers periodic boundary conditions and constant coefficients in the differential operator. In this spirit, the very elegant time discretization error analysis of a quasilinear wave equation by trigonometric integrators in [13] is easily extended to a full discretization due to the Fourier spectral methods in space. In [1, 2], exponential splitting methods are applied to parabolic evolution equations and the order reduction induced by the boundary conditions is investigated. For finite difference and spectral discretization, the results are extended to full discretization error bounds.

However, none of these error analyses is applicable to general wave equations with non-constant coefficients on general domains and, in particular, not to unstructured meshes and nonconforming space discretizations. The main contribution of our paper is to fill this gap. We provide full discretization error bounds in terms of the stiff order of the exponential integrator in time and the abstract space discretization error terms from the unified error analysis. Due to the general framework, these bounds apply to a large class of equations and space discretizations, e.g., those considered in [15, 16]. To prove these bounds we have to intertwine the techniques used in the proofs for the time and space discretization, respectively. A crucial difficulty of the error analysis is to come up with a suitable representation of the exact solution involving the discretized semigroup. This is in sharp contrast to spectral space discretizations mentioned above, where the projection onto the finite dimensional space usually commutes with the exact semigroup. Further, this issue has not occurred so far in the analyses of standard implicit time integration methods.

The rest of the paper is structured as follows: In Section 2, we introduce an abstract framework adapted to the first-order formulation of (1.1), and present the abstract space discretization. Further, we collect the properties of the discrete objects in order to perform the error analysis.

The fully discrete schemes are presented in Section 3. We discuss the methods and state our main results for the first-order system. The proofs are given in Section 4. We first derive the defects for the exact solution expressed by a modified variation-of-constants formula and provide bounds on those. From this, the fully discrete errors bounds are established. We then transfer our results in Section 5 to the abstract second-order evolution equation (1.1) to conclude the corresponding novel error bounds.

In Section 6, we discuss a concrete example of a wave equation with kinetic boundary conditions and present numerical experiments to confirm our theoretical findings.

## 2. GENERAL SETTING AND THE UNIFIED ERROR ANALYSIS

In this section, we introduce the analytical framework for the second-order evolution equation (1.1). Since the exponential integrators considered in this paper are applied to first-order systems, we rewrite (1.1) in a first-order form. For this, we present an abstract space discretization established in [15, 16], where a unified error analysis for space discretizations of linear and semilinear wave-type equations was derived. We recall the setting to keep this presentation self-contained.

**2.1. The continuous second-order problem.** Let  $V, H$  be Hilbert spaces and  $V$  be densely and continuously embedded in  $H$ . We consider the following variational differential equation (2.1) as a prototype for weak formulations of second-order wave equations: seek  $u \in C^2([0, T]; H) \cap C^1([0, T]; V)$  such that

$$(2.1) \quad \begin{aligned} m(u'', v) + b(u', v) + a(u, v) &= m(f(t, u), v) \quad \text{for all } v \in V, t \in (0, T], \\ u(0) &= u^0, \quad u'(0) = v^0. \end{aligned}$$

We pose the following assumptions.

*Assumption 2.1.*

- a) The bilinear form  $m$  is a scalar product on  $H$  with induced norm  $\|\cdot\|_m$ .
- b)  $a: V \times V \rightarrow \mathbb{R}$  is a symmetric bilinear form and there exists a constant  $c_G \geq 0$  such that

$$\tilde{a} := a + c_G m$$

is a scalar product on  $V$  with induced norm  $\|\cdot\|_{\tilde{a}}$ .

- c) The bilinear form  $b: V \times H \rightarrow \mathbb{R}$  is continuous and there exists a  $\beta_{\text{qm}} \geq 0$  such that

$$b(v, v) + \beta_{\text{qm}} \|v\|_m^2 \geq 0 \quad \text{for all } v \in V.$$

- d) The nonlinearity  $f$  satisfies  $f \in C^1([0, T] \times V; H)$  and is locally Lipschitz-continuous on  $V$  with Lipschitz-constant  $L_{T,M}$ , i.e., for all  $t \in [0, T]$  and all  $v, w \in V$  with  $\|v\|_{\tilde{a}}, \|w\|_{\tilde{a}} \leq M$  it holds

$$\|f(t, v) - f(t, w)\|_m \leq L_{T,M} \|v - w\|_{\tilde{a}}.$$

By the dense and continuous embedding of the Hilbert spaces there exists a constant  $C_{H,V} > 0$  such that

$$\|v\|_m \leq C_{H,V} \|v\|_{\tilde{a}} \quad \text{for all } v \in V.$$

We define operators  $A: D(A) \rightarrow H$  and  $B: V \rightarrow H$  induced by the bilinear forms  $a$  and  $b$  via

$$\begin{aligned} m(Av, w) &= a(v, w), & \text{for all } v \in D(A), w \in V, \\ m(Bv, w) &= b(v, w), & \text{for all } v \in V, w \in H, \end{aligned}$$

with

$$D(A) = \{v \in V \mid \exists C = C(v) > 0 \text{ such that } \forall w \in V : |a(v, w)| \leq C \|w\|_m\}.$$

By the construction of the operators  $A$  and  $B$ , a solution of (2.1) additionally satisfies  $u \in C([0, T]; D(A))$ . Hence, problem (2.1) is equivalent to the evolution equation (1.1). In order to rewrite (1.1) in a first-order formulation, we define  $X = V \times H$ ,  $u'(t) = v(t)$  and set

$$(2.2) \quad x(t) = \begin{bmatrix} u(t) \\ v(t) \end{bmatrix}, \quad S = \begin{bmatrix} 0 & -\mathbb{I} \\ A & B \end{bmatrix}, \quad g(t, x) = \begin{bmatrix} 0 \\ f(t, u) \end{bmatrix}, \quad x^0 = \begin{bmatrix} u^0 \\ v^0 \end{bmatrix}.$$

Then (1.1) is equivalent to the evolution equation

$$(2.3) \quad \begin{aligned} x'(t) + Sx(t) &= g(t, x(t)), & t \in (0, T], \\ x(0) &= x^0. \end{aligned}$$

**2.2. The continuous first-order problem.** In the following sections, we consider a general system of the form (2.3) on a Hilbert space  $X$  with scalar product  $p(\cdot, \cdot)$ . In this setting, we derive the fully discrete schemes and perform the error analysis. In Section 5, we transfer these results back to the formulation (1.1).

We pose the following classical assumptions, which are a direct consequence of Assumption 2.1, cf. [16, Sec. 3.2].

*Assumption 2.2.*

- a) The linear operator  $S: D(S) \rightarrow X$  is the generator of a  $C_0$ -semigroup with
- $$(2.4) \quad \left\| e^{-tS} \right\|_{X \leftarrow X} \leq e^{c_{\text{qm}} t}$$
- for some  $c_{\text{qm}} \geq 0$ .
- b) The nonlinearity  $g \in C^1([0, T] \times X; X)$  is locally Lipschitz continuous w.r.t. the second component with constant  $L_{T,M}$ .

Assumption 2.2 ensures local wellposedness of (2.3), cf. [30, Thm. 6.1.5].

**Lemma 2.3.** *If Assumption 2.2 holds true, then (2.3) is locally wellposed, i.e., for every  $x^0 \in X$  there exists  $t^*(x^0) > 0$  such that, for all  $T < t^*(x^0)$ , (2.3) has a unique solution*

$$x \in C^1([0, T]; X) \cap C([0, T]; D(S)).$$

For the rest of the paper we fix some  $T < t^*(x^0)$  in order to obtain uniform bounds of the solution on  $[0, T]$ .

**2.3. Abstract space discretization.** We now introduce an abstract space discretization of the evolution equation (2.3). We use the setting introduced in [15, 16] to analyze nonconforming space discretizations of linear and semilinear wave-type equations.

For this, let  $(X_h)_h$  be a family of finite dimensional vector spaces related to a discretization parameter  $h$ , e.g., the maximal mesh width of a finite element discretization. In  $X_h$ , we seek an approximation  $x_h$  to the solution  $x$  of (2.3). Let  $p_h(\cdot, \cdot)$  be a scalar product on  $X_h$  and let  $S_h \in \mathcal{L}(X_h, X_h)$  and  $g_h: [0, T] \times X_h \rightarrow X_h$  be discretizations of  $S$  and  $g$ , respectively. We impose the following conditions similar to Assumption 2.2.

*Assumption 2.4.*

- a) The linear operator  $S_h \in \mathcal{L}(X_h; X_h)$  is the generator of a  $C_0$ -semigroup with
- $$(2.5) \quad \left\| e^{-tS_h} \right\|_{X_h \leftarrow X_h} \leq e^{\widehat{c}_{\text{qm}} t}.$$
- b) The nonlinearity  $g_h: [0, T] \times X_h \rightarrow X_h$  is locally Lipschitz continuous w.r.t. the second component with constant  $\widehat{L}_{T,M}$ , i.e., for all  $x_h, y_h \in X_h$  with  $\|x_h\|_{X_h}, \|y_h\|_{X_h} \leq M$  and  $t \in [0, T]$ :

$$(2.6) \quad \|g_h(t, x_h) - g_h(t, y_h)\|_{X_h} \leq \widehat{L}_{T,M} \|x_h - y_h\|_{X_h}.$$

The constants  $\widehat{c}_{\text{qm}}$  and  $\widehat{L}_{T,M}$  are independent of  $h$ .

The discretized evolution equation (2.3) is then of the form

$$(2.7) \quad \begin{aligned} x_h'(t) + S_h x_h(t) &= g_h(t, x_h(t)), & t \in (0, T], \\ x_h(0) &= x_h^0, \end{aligned}$$




FIGURE 1. Overview of spaces and operators, taken from [16, Fig. 3.1].

where  $x_h^0 \in X_h$  is an approximation of  $x^0$ .

For exponential Runge–Kutta methods of order  $p \geq 3$  we require additional regularity of the discretized nonlinearity which is beyond the Lipschitz property given in (2.6).

*Assumption 2.5.* The nonlinearity  $g_h$  is in  $C^3([0, T] \times X_h; X_h)$  with derivatives bounded independent of  $h$ .

For a concrete example, we comment on this assumption in Remark 6.1.

**2.4. Abstract space discretization errors.** We explicitly allow for nonconforming space discretizations where  $X_h \not\subseteq X$ . Thus, to relate the continuous and discrete solution, we make the following assumptions:

*Assumption 2.6.*

- a) There exists a *lift operator*  $\mathcal{L}_h: X_h \rightarrow X$  which satisfies

$$(2.8) \quad \|\mathcal{L}_h y_h\|_X \leq \widehat{C}_X \|y_h\|_{X_h} \quad \text{for all } y_h \in X_h$$

with  $\widehat{C}_X$  independent of  $h$ .

By  $\mathcal{L}_h^*: X \rightarrow X_h$  we denote the adjoint of the lift operator defined via

$$p_h(\mathcal{L}_h^* y, y_h) = p(y, \mathcal{L}_h y_h) \quad \text{for all } y \in X, y_h \in X_h.$$

- b) There exists a Hilbert space  $Z$ , which is densely and continuously embedded in  $X$ , and a *reference operator*  $J_h \in \mathcal{L}(Z, X_h)$  which satisfies

$$(2.9) \quad \|J_h z\|_{X_h} \leq \widehat{C}_{J_h} \|z\|_Z \quad \text{for all } z \in Z$$

with a constant  $\widehat{C}_{J_h}$  independent of  $h$ .

The reference operator  $J_h$  could, e.g., be an interpolation operator defined on a subspace  $Z$  of the continuous functions, and should satisfy  $\mathcal{L}_h J_h z \approx z$  for all  $z \in Z$ . Figure 1 illustrates the relations of the spaces and operators.

In our error analysis, we will bound the space discretization errors in terms of the following quantities:

**Definition 2.7** (Space discretization errors).

- a) The *linear remainder operator* is defined via

$$(2.10) \quad R_h := \mathcal{L}_h^* S - S_h J_h: D(S) \cap Z \rightarrow X_h.$$

- b) The *nonlinear remainder operator*  $r_h: [0, T] \times Z \rightarrow X_h$  is given by

$$(2.11) \quad r_h(t, z) := \mathcal{L}_h^* g(t, z) - g_h(t, J_h z).$$

c) The space discretization errors are collected in the term

$$(2.12) \quad \begin{aligned} E_h(t) = & \left\| x_h^0 - J_h x^0 \right\|_{X_h} + t \left\| \Delta_{J_h} x' \right\|_{L^\infty([0,t]; X_h)} \\ & + t \left\| R_h x \right\|_{L^\infty([0,t]; X_h)} + t \left\| r_h(\cdot, x(\cdot)) \right\|_{L^\infty([0,t]; X_h)} \end{aligned}$$

where  $\Delta_{J_h} := J_h - \mathcal{L}_h^* : Z \rightarrow X_h$  is the *reference error*.

Additionally, for our analysis we need the following assumptions on the regularity of the exact solution and the consistency of the discretization.

*Assumption 2.8.* The solution of (2.3) satisfies  $x \in C^1([0, T]; Z)$  and the error terms in (2.12) converge to zero, i.e.,

$$E_h(T) \rightarrow 0 \quad \text{for } h \rightarrow 0.$$

*Remark 2.9.* The assumptions are satisfied for wave equations with various boundary conditions, in particular of dynamical and Dirichlet type, discretized by non-conforming finite elements, cf. [16, 25]. More details are provided in Sections 5 and 6. A further example covered by the framework, are discontinuous Galerkin methods applied to linear Maxwell's equations, cf. [15].

### 3. EXPONENTIAL INTEGRATORS AND MAIN RESULTS

In the following we present the fully discrete schemes obtained by applying exponential integrators to the spatially discretized evolution equation (2.7) and state our main results, the full discretization error bounds. We denote by  $\tau > 0$  the time step size and set  $t_n = n\tau$ . Further, we abbreviate

$$(3.1) \quad t_{n+s} := t_n + \tau s, \quad s \in [0, 1],$$

and denote the fully discrete approximation at time  $t_n$  by  $x_h^n \approx x(t_n)$ . Most exponential integrators are based on the variation-of-constants formula for the solution of (2.7)

$$(3.2) \quad x_h(t_{n+1}) = e^{-\tau S_h} x_h(t_n) + \tau \int_0^1 e^{-(1-s)\tau S_h} g_h(t_{n+s}, x_h(t_{n+s})) ds,$$

where only the nonlinearity  $g_h$  is approximated. Applying Taylor expansion to  $s \mapsto g_h(t_{n+s}, x_h(t_{n+s}))$ , the  $\varphi$ -functions given by

$$(3.3) \quad \varphi_{k+1}(z) = \int_0^1 e^{(1-s)z} \frac{s^k}{k!} ds, \quad k \geq 0,$$

appear naturally. They are analytic in  $\mathbb{C}$  and closely related to the exponential function. In order to expand the right-hand side, we assume the following differentiability.

*Assumption 3.1.* Let  $m \geq 1$  and  $x$  be the exact solution of (2.3). Then, the differentiability condition

$$s \mapsto g(s, x(s)) \in C^m([0, T]; X)$$

holds.

We start with the prototypical example, namely the exponential Euler method. It is constructed by freezing the right-hand side in (3.2) at the last approximation and is given by

$$(3.4) \quad x_h^{n+1} = e^{-\tau S_h} x_h^n + \tau \varphi_1(-\tau S_h) g_h(t_n, x_h^n).$$

Concerning the time integration, it is well known that the stiff order of (3.4) is one, cf. [18, Thm. 4.2]. This is actually a special case of Theorem 3.2. Higher order methods can be constructed for example in the following two ways.

**3.1. Exponential Adams methods.** On the one hand, we consider the exponential  $k$ -step Adams methods. They were proposed in [29] and time integration error bounds were derived in [20]. The schemes can be expressed as

$$(3.5) \quad x_h^{n+1} = e^{-\tau S_h} x_h^n + \tau \int_0^1 e^{-(1-s)\tau S_h} p_h^{n,k}(t_{n+s}) ds, \quad k \geq 1,$$

where  $p_h^{n,k}$  is the interpolation polynomial through the points

$$(3.6) \quad \left( (t_{n-k+1}, g_h^{n-k+1}), \dots, (t_n, g_h^n) \right) \quad \text{with} \quad g_h^j := g_h(t_j, x_h^j).$$

We note that the integral in (3.5) can also be written as a linear combination of the  $\varphi$ -functions  $\varphi_1, \dots, \varphi_k$  applied to  $g_h^j$ . The exponential Euler scheme (3.4) is obtained by setting  $k = 1$ .

Closely related methods, using the variation-of-constants formula (3.2) from  $t_n$  to  $t_{n+k}$ , have been constructed and analyzed in [7]. We expect that for these methods the results obtained in the following can be achieved fully analogously and, hence, we do not further comment on this.

**Theorem 3.2.** *Let Assumptions 2.2, 2.4, 2.6, 2.8, and 3.1 for  $m = k$  hold. Further, consider the exact solution  $x$  of (2.3) and the approximations  $x_h^n$  obtained from (3.5). Then, there are  $\tau_0, h_0 > 0$  and a constant  $C > 0$  such that for all  $\tau \leq \tau_0, h \leq h_0$ , and  $t_k \leq t_n \leq T$  it holds*

$$\begin{aligned} \|\mathcal{L}_h x_h^n - x(t_n)\|_X &\leq \|(\mathcal{L}_h J_h - \mathbf{I})x(t_n)\|_X + C(\tau^k + E_h(t_n)) \\ &\quad + C \sum_{j=1}^{k-1} \|x_h^j - J_h x(t_j)\|_{X_h}, \end{aligned}$$

where  $C$  is independent of  $\tau$  and  $h$ , and  $E_h(t_n)$  is defined in (2.12).

*Remark 3.3.*

- a) For  $k = 2$ , an exponential Euler step yields a sufficiently good starting value  $x_h^1$ . For  $k \geq 3$ , we suggest computing the starting values  $x_h^j, j = 1, \dots, k-1$ , via the starting procedure proposed in [7, 20]. There, one has to solve a nonlinear system of equations which is usually done by fixed point iteration. We expect that by our proofs, one can derive the same error bounds as for the following approximations, but we omit the details here.
- b) We note that the constant  $C$  can actually be written in the form

$$C = C(t_n) = c_1 t_n e^{c_2 t_n}$$

with constants  $c_1, c_2 > 0$  independent of the time  $t_n$ .

TABLE 1. Stiff order conditions for explicit exponential Runge–Kutta methods in (3.7), cf. [18, Table 2]. Here  $\mathcal{D}_h$  and  $\mathcal{K}_h$  denote arbitrary bounded operators on  $X_h$ . The functions  $\psi_i$  and  $\psi_{k,\ell}$  are defined in (3.8).

Order	Order condition
1	$\psi_1(-\tau S_h) = 0$
2	$\psi_2(-\tau S_h) = 0$ $\psi_{1,i}(-\tau S_h) = 0$
3	$\psi_3(-\tau S_h) = 0$ $\sum_{i=1}^s b_i(-\tau S_h) \mathcal{D}_h \psi_{2,i}(-\tau S_h) = 0$ $\psi_4(-\tau S_h) = 0$
4	$\sum_{i=1}^s b_i(-\tau S_h) \mathcal{D}_h \psi_{3,i}(-\tau S_h) = 0$ $\sum_{i=1}^s b_i(-\tau S_h) \mathcal{D}_h \sum_{j=2}^{i-1} a_{ij}(-\tau S_h) \mathcal{D}_h \psi_{2,j}(-\tau S_h) = 0$ $\sum_{i=1}^s b_i(-\tau S_h) c_i \mathcal{K}_h \psi_{2,i}(-\tau S_h) = 0$

**3.2. Exponential Runge–Kutta methods.** Further, we consider the class of explicit  $s$ -stage exponential Runge–Kutta methods introduced in [12]. Error bounds for the time integration were derived in [18]. Given analytic coefficients satisfying  $|a_{ij}(z)|, |b_i(z)| \leq C$  for  $\operatorname{Re} z \leq \widehat{c}_{\text{qm}}$ , they read

$$(3.7) \quad \begin{aligned} X_h^{ni} &= e^{-c_i \tau S_h} x_h^n + \tau \sum_{j=1}^{i-1} a_{ij}(-\tau S_h) g_h(t_{n+c_j}, X_h^{nj}), \quad i = 1, \dots, s, \\ x_h^{n+1} &= e^{-\tau S_h} x_h^n + \tau \sum_{i=1}^s b_i(-\tau S_h) g_h(t_{n+c_i}, X_h^{ni}). \end{aligned}$$

For the error analysis, several order conditions are necessary in order to ensure convergence also in the stiff case. They relate the  $\varphi$ -functions defined in (3.3) to certain combinations of the coefficients  $a_{ij}, b_i$  and use the following quantities

$$(3.8a) \quad \psi_j(z) = \varphi_j(z) - \sum_{k=1}^s b_k(z) \frac{c_k^{j-1}}{(j-1)!},$$

$$(3.8b) \quad \psi_{j,i}(z) = \varphi_j(c_i z) c_i^j - \sum_{k=1}^{i-1} a_{ik}(z) \frac{c_k^{j-1}}{(j-1)!}.$$

In Table 1, we collected the order conditions derived in [18]. In particular, we are interested in the methods satisfying the stiff order conditions for  $p = 2, 3, 4$ . For the methods, which we present in the following, the corresponding order conditions of Table 1 are satisfied for all  $h > 0$ .

In order to obtain schemes of order  $p = 2$  using  $s = 2$  stages, they lead to the choices

$$(3.9) \quad a_{21}(z) = c_2 \varphi_1(c_2 z), \quad b_1(z) = \varphi_1(z) - \frac{1}{c_2} \varphi_2(z), \quad b_2(z) = \frac{1}{c_2} \varphi_2(z).$$

To obtain order  $p = 3$ , we can for example use the method (5.16) from [23, eq. (51)]

$$(3.10) \quad \begin{array}{c|ccc} 0 & & & \\ \frac{1}{2} & \frac{1}{2}\varphi_{1,2} & & \\ \frac{1}{2} & \frac{1}{2}\varphi_{1,2} - \varphi_{2,2} & \varphi_{2,2} & \\ 1 & \varphi_1 - 2\varphi_2 & 0 & 2\varphi_2 \\ \hline & \varphi_1 - 3\varphi_2 + 4\varphi_3 & 2\varphi_2 - 4\varphi_3 & 2\varphi_2 - 4\varphi_3 \quad -\varphi_2 + 4\varphi_3 \end{array}$$

which satisfies the stiff order conditions up to order  $p = 3$  with  $\varphi_{j,2} := \varphi_j(\frac{1}{2})$ . Another example for a method of order  $p = 3$  is given in [32, Example 4.5.5]. Using 8 inner stages, a method satisfying the order conditions up to order  $p = 4$  is proposed in [27, Sec. 5], but we refrain from stating the coefficients.

For these methods, we derive the following error bounds of optimal order.

**Theorem 3.4.** *Let  $p \in \{2, 3, 4\}$  and let Assumptions 2.2, 2.4, 2.6, 2.8, and 3.1 hold for  $m = p$ . For  $p \in \{3, 4\}$  we additionally require Assumption 2.5. Further, consider the exact solution  $x$  of (2.3) and the approximations  $x_h^n$  obtained from (3.7) with coefficients satisfying the order conditions of Table 1 up to order  $p$ . Then, there are  $\tau_0, h_0 > 0$  and a constant  $C > 0$  such that for all  $\tau \leq \tau_0$  and  $h \leq h_0$  it holds*

$$(3.11) \quad \|\mathcal{L}_h x_h^n - x(t_n)\|_X \leq \|(\mathcal{L}_h J_h - \mathbf{I})x(t_n)\|_X + C(\tau^p + E_h(t_n)), \quad t_n \leq T,$$

where  $C$  is independent of  $\tau$  and  $h$ , and  $E_h(t_n)$  is defined in (2.12).

*Remark 3.5.* In the papers [7, 18, 20, 27], a special focus is put on parabolic problems such as the heat equation. When dealing with analytical semigroups the smoothing property can be heavily exploited and the order conditions can be severely relaxed. In this case, second-order convergence is also achieved by

$$(3.12) \quad a_{21}(z) = c_2\varphi_1(c_2z), \quad b_1(z) = \left(1 - \frac{1}{2c_2}\right)\varphi_1(z), \quad b_2(z) = \frac{1}{2c_2}\varphi_1(z),$$

which does not require the evaluation of the  $\varphi_2$ -function compared to (3.9). Furthermore, the parabolic smoothing allows to gain one order in  $\tau$  for the schemes of order  $p = 3$  and  $p = 4$  above, cf. [18, Thm. 4.7] and [27, Thm. 4.1].

#### 4. PROOFS OF THE MAIN RESULTS

In the following section, we aim to bound the fully discrete error which can be decomposed into

$$(4.1) \quad \begin{aligned} \mathcal{L}_h x_h^n - x(t_n) &= \mathcal{L}_h(x_h^n - J_h x(t_n)) + (\mathcal{L}_h J_h - \mathbf{I})x(t_n) \\ &=: \mathcal{L}_h e_h^n + e_{J_h}^n. \end{aligned}$$

The error  $e_{J_h}^n$  is an approximation error of the spatial discretization. It is independent of the time integration method. By (2.8) it holds

$$\|\mathcal{L}_h x_h^n - x(t_n)\|_X \leq \widehat{C}_X \|e_h^n\|_{X_h} + \|e_{J_h}^n\|_X,$$

and thus it is sufficient to bound the first term. To shorten the notation further, we introduce

$$(4.2) \quad \widehat{x}^{n+s} := x(t_{n+s}), \quad \widehat{g}^{n+s} := g(t_{n+s}, \widehat{x}^{n+s}), \quad \widehat{g}_h^{n+s} := g_h(t_{n+s}, J_h \widehat{x}^{n+s})$$

for the exact solution as well as the exact and discretized nonlinearity evaluated at the exact solution.

The first result of this section is concerned with the representation of the exact solution. Simply using the variation-of-constants formula, we are not able to provide suitable bounds of the defect, as we exemplify for the exponential Euler method. We insert  $J_h \tilde{x}^n$  into the numerical scheme (3.4) and obtain the defect

$$\delta^n = J_h \tilde{x}^{n+1} - (e^{-\tau S_h} J_h \tilde{x}^n + \tau \varphi_1(-\tau S_h) \tilde{g}_h^n).$$

Expanding  $J_h \tilde{x}^{n+1}$  by the variation-of-constants formula, one readily observes that the defect  $\delta^n$  contains the term

$$(4.3) \quad J_h e^{-\tau S} \tilde{x}^n - e^{-\tau S_h} J_h \tilde{x}^n = \int_0^\tau e^{-(\tau-s)S_h} (S_h J_h - J_h S) e^{-sS} \tilde{x}^n ds.$$

Writing

$$S_h J_h - J_h S = (S_h J_h - \mathcal{L}_h^* S) + (\mathcal{L}_h^* - J_h) S$$

and using  $R_h$  and  $\Delta_{J_h}$  from Definition 2.7, one can bound (4.3) in  $X_h$  in terms of

$$(4.4) \quad \|R_h e^{-sS} \tilde{x}^n\|_{X_h} + \|\Delta_{J_h} e^{-sS} S \tilde{x}^n\|_{X_h}, \quad s \in [0, \tau].$$

However, suitable bounds on these terms are in general not available, as we explain for the first term in the following counterexample.

**Example 4.1.** Let  $\Omega = (0, 1)$ ,  $V = H_0^1(\Omega)$ ,  $H = L^2(\Omega)$ ,  $A = -\Delta$ ,  $B = 0$  in (2.2). Then,  $e^{-\tau S} x^0$  is the solution at time  $t = \tau$  of the one-dimensional wave equation

$$(4.5) \quad u_{tt}(t, \mathbf{x}) - \Delta u(t, \mathbf{x}) = 0, \quad \mathbf{x} \in \Omega, \quad t \geq 0,$$

subject to homogeneous Dirichlet boundary conditions and initial values  $u^0, v^0$ .

The remainder operator  $R_h$  applied to  $x = [u, v]^\top$  contains, among others, interpolation errors of  $u$  and  $v$ , cf. [15, Lem. 4.7]. Hence, when discretizing with finite elements of order  $q$ , a bound for (4.4) of order  $\mathcal{O}(h^q)$  requires at least that  $e^{-tS} x^0 \in H^{q+1}(\Omega) \times H^{q+1}(\Omega)$ . However, choosing

$$u^0(\mathbf{x}) = u(0, \mathbf{x}) = \mathbf{x}(1 - \mathbf{x}), \quad v^0(\mathbf{x}) = u_t(0, \mathbf{x}) = 0,$$

we note that  $u^0, v^0 \in C^\infty(\bar{\Omega})$ . Extending the initial values to an odd function on  $[-1, 1]$  and then periodically to  $\mathbb{R}$ , the exact solution is given by d'Alembert's formula. Since the extended version of  $u^0$  is not  $C^2$  at  $\mathbf{x} = \ell$ ,  $\ell \in \mathbb{Z}$ , we obtain that  $u(t, \cdot) \notin H^{q+1}(\Omega)$ ,  $q \geq 2$ , for almost all  $t \geq 0$ .  $\diamond$

Hence, we take a different approach that only requires regularity of the exact solution  $x$ . The crucial point is to write  $J_h x$  in a modified variation-of-constants formula that is driven by the discretized operator  $S_h$  up to an error covered by (2.12).

**Lemma 4.2.** *Let Assumptions 2.2, 2.4, and 2.6 hold, and let  $x \in C^1([0, T]; Z)$  be the solution of (2.3). Then, we have the representation*

$$(4.6) \quad J_h \tilde{x}^{n+1} = e^{-\tau S_h} J_h \tilde{x}^n + \tau \int_0^1 e^{-(1-s)\tau S_h} \mathcal{L}_h^* \tilde{g}^{n+s} ds + E_{HG}^n(\tau)$$

with remainder  $E_{HG}^n(\tau)$  bounded by

$$(4.7) \quad \|E_{HG}^n(\tau)\|_{X_h} \leq \tau e^{\tau \hat{c}_{\text{qm}}} \left( \|\Delta_{J_h} x'\|_{L^\infty([t_n, t_{n+1}]; X_h)} + \|R_h x\|_{L^\infty([t_n, t_{n+1}]; X_h)} \right),$$

where  $\Delta_{J_h}$  and  $R_h$  are given in Definition 2.7.

*Proof.* We use equation (2.3) to obtain

$$\begin{aligned} J_h x'(t) &= \mathcal{L}_h^* x'(t) + \Delta_{J_h} x'(t) \\ &= -\mathcal{L}_h^* S x(t) + \mathcal{L}_h^* g(t, x(t)) + \Delta_{J_h} x'(t) \\ &= -S_h J_h x(t) + \mathcal{L}_h^* g(t, x(t)) - R_h x(t) + \Delta_{J_h} x'(t). \end{aligned}$$

Applying the variation-of-constants formula and (2.5) yields the assertion.  $\square$

The following two subsections treat the classes of exponential integrators separately. However, all the error estimates rely on the representation of the exact solution provided by Lemma 4.2. Throughout, we use a local in time version of  $E_h(t)$  defined in (2.12), given by

$$(4.8) \quad \begin{aligned} \varepsilon_h^n &= \|R_h x\|_{L^\infty([t_n, t_{n+1}]; X_h)} + \|r_h(\cdot, x(\cdot))\|_{L^\infty([t_n, t_{n+1}]; X_h)} \\ &\quad + \|\Delta_{J_h} x'\|_{L^\infty([t_n, t_{n+1}]; X_h)} \end{aligned}$$

to account for the spatial defects.

**4.1. Exponential Adams methods.** Using the notation defined in (4.2), for the analysis we introduce the two auxiliary interpolation polynomials  $\tilde{p}_h^{n,k}$  through the discretized nonlinearity evaluated at  $J_h x$

$$\left( (t_{n-k+1}, \tilde{g}_h^{n-k+1}), \dots, (t_n, \tilde{g}_h^n) \right),$$

and  $\tilde{p}_h^{n,k}$  through the continuous nonlinearity evaluated at exact solution  $x$

$$\left( (t_{n-k+1}, \tilde{g}^{n-k+1}), \dots, (t_n, \tilde{g}^n) \right).$$

**Lemma 4.3.** *Under the assumptions of Theorem 3.2, the exact solution satisfies*

$$(4.9) \quad J_h \tilde{x}^{n+1} = e^{-\tau S_h} J_h \tilde{x}^n + \tau \int_0^1 e^{-(1-s)\tau S_h} \tilde{p}_h^{n,k}(t_{n+s}) \, ds + \delta^n, \quad n \geq k,$$

where the defect is bounded by

$$\|\delta^n\|_{X_h} \leq C\tau \left( \tau^k + \sum_{j=0}^{k-1} \varepsilon_h^{n-j} \right)$$

with a constant  $C$  independent of  $\tau$  and  $h$ , and  $\varepsilon_h^n$  defined in (4.8).

*Proof.* Employing (4.6) and (4.9), we obtain

$$\begin{aligned} \delta^n &= \tau \int_0^1 e^{-(1-s)\tau S_h} (\mathcal{L}_h^* \tilde{g}^{n+s} - \tilde{p}_h^{n,k}(t_{n+s})) \, ds + E_{\text{HG}}^n(\tau) \\ &= \tau \int_0^1 e^{-(1-s)\tau S_h} \mathcal{L}_h^* (\tilde{g}^{n+s} - \tilde{p}_h^{n,k}(t_{n+s})) \, ds \\ &\quad + \tau \int_0^1 e^{-(1-s)\tau S_h} \mathcal{L}_h^* \tilde{p}_h^{n,k}(t_{n+s}) - \tilde{p}_h^{n,k}(t_{n+s}) \, ds + E_{\text{HG}}^n(\tau). \end{aligned}$$

The first term is bounded using the standard interpolation bounds and Assumption 3.1. The second term yields a linear combination of the nonlinear remainders

$r_h(t_{n-j}, \tilde{x}^{n-j})$ ,  $j = 0, \dots, k-1$ , given in (2.11), and for the last term we employ the bound (4.7).  $\square$

From the bounds on the defects, we conclude our first main result.

*Proof of Theorem 3.2.* We recall that by (4.1) it is sufficient to bound  $e_h^n$  and assume first that the numerical approximations  $x_h^n$  are bounded. Subtracting (4.9) from (3.5), we derive the error recursion

$$\begin{aligned} e_h^{n+1} &= e^{-\tau S_h} e_h^n + \tau \int_0^1 e^{-(1-s)\tau S_h} (p_h^{n,k}(t_{n+s}) - \tilde{p}_h^{n,k}(t_{n+s})) ds - \delta^n \\ &= e^{-\tau S_h} e_h^n + \tau \sum_{j=0}^{k-1} L^{n,j}(x_h^{n-j}, \tilde{x}^{n-j}) - \delta^n, \end{aligned}$$

where by Assumption 2.4 it holds with a constant  $C_{\hat{L}}$  independent of  $\tau$  and  $h$

$$\|L^{n,j}(x_h^{n-j}, \tilde{x}^{n-j})\|_{X_h} \leq C_{\hat{L}} e^{\hat{c}_{\text{qm}}\tau} \|e_h^{n-j}\|_{X_h}.$$

Hence, resolving the error recursion and taking norms, we obtain for  $n \geq k$

$$\|e_h^n\|_{X_h} \leq e^{\hat{c}_{\text{qm}}t_n} k \sum_{m=0}^{k-1} \|e_h^m\|_{X_h} + kC_{\hat{L}} \sum_{m=k}^{n-1} e^{\hat{c}_{\text{qm}}(t_n-t_m)} \|e_h^m\|_{X_h} + e^{\hat{c}_{\text{qm}}t_n} \sum_{m=1}^n \|\delta^m\|_{X_h}.$$

Multiplication with  $e^{-\hat{c}_{\text{qm}}t_n}$  and the application of a discrete Gronwall lemma yields the desired error bound. By an induction argument, Assumption 2.8 yields the existence of  $h_0, \tau_0 > 0$  such that the numerical approximations indeed remain bounded and the claim follows.  $\square$

**4.2. Exponential Runge–Kutta methods.** Our error analysis within this section is based on the analysis for the time discretization presented in [18]. In the following, we show how to extend these results in the presence of additional spatial errors. We first insert the reference operator  $J_h$  applied to the exact solution  $x$  into the numerical scheme and obtain with the defects  $\Delta_h^{n,i}, \delta_h^n$

$$\begin{aligned} (4.10) \quad J_h \tilde{x}^{n+c_i} &= e^{-c_i \tau S_h} J_h \tilde{x}^n + \tau \sum_{j=1}^{i-1} a_{ij}(-\tau S_h) \tilde{g}_h^{n+c_j} + \Delta_h^{n,i}, \\ J_h \tilde{x}^{n+1} &= e^{-\tau S_h} J_h \tilde{x}^n + \tau \sum_{i=1}^s b_i(-\tau S_h) \tilde{g}_h^{n+c_i} + \delta_h^n. \end{aligned}$$

Introducing the error of the inner stages  $E_h^{ni} = X_h^{ni} - J_h \tilde{x}^{n+c_i}$ , we derive the error recursions

$$\begin{aligned} (4.11) \quad E_h^{ni} &= e^{-c_i \tau S_h} e_h^n + \tau \sum_{j=1}^{i-1} a_{ij}(-\tau S_h) (g_h(t_{n+c_j}, X_h^{nj}) - \tilde{g}_h^{n+c_j}) - \Delta_h^{n,i}, \\ e_h^{n+1} &= e^{-\tau S_h} e_h^n + \tau \sum_{i=1}^s b_i(-\tau S_h) (g_h(t_{n+c_i}, X_h^{ni}) - \tilde{g}_h^{n+c_i}) - \delta_h^n. \end{aligned}$$

In the following, we establish bounds on  $E_h^{n,i}$  and  $e_h^{n+1}$ . In large parts, we follow the arguments in [18] and, use the shorthand notation

$$\tilde{g}^{(j),n} := \frac{d^j}{dt^j} g(t, x(t)) \Big|_{t=t_n}$$

which is well-defined by Assumption 3.1. As a first result, derive the structure of the defects. This is an extension of [18, Lem. 4.1] incorporating the spatial defects  $\Delta_{h,\text{sp}}^{n,i}$ ,  $\delta_{h,\text{sp}}^n$ .

**Lemma 4.4.** *Let Assumptions 2.2, 2.4, 2.6, and 3.1 for  $m = p$  hold. The defects can be expressed as*

$$\begin{aligned} \Delta_h^{n,i} &= \Delta_{h,\text{sp}}^{n,i} + \Delta_{h,\tau}^{n,i} + \sum_{j=1}^{p-1} \tau^j \psi_{j,i}(-\tau S_h) \mathcal{L}_h^* \tilde{g}^{(j-1),n}, \\ \delta_h^n &= \delta_{h,\text{sp}}^n + \delta_{h,\tau}^n + \sum_{j=1}^p \tau^j \psi_j(-\tau S_h) \mathcal{L}_h^* \tilde{g}^{(j-1),n}. \end{aligned}$$

They satisfy

$$(4.12) \quad \|\Delta_{h,\text{sp}}^{n,i}\|_{X_h} + \|\delta_{h,\text{sp}}^n\|_{X_h} \leq C\tau \varepsilon_h^n, \quad \|\Delta_{h,\tau}^{n,i}\|_{X_h} \leq C\tau^p, \quad \|\delta_{h,\tau}^n\|_{X_h} \leq C\tau^{p+1}$$

with constants  $C$  independent of  $\tau$  and  $h$ , and  $\varepsilon_h^n$  defined in (4.8).

*Proof.* We only prove the claim for  $\delta_h^n$  since the argument for  $\Delta_h^{n,i}$  is easily deduced from it. We obtain from (4.10), Lemma 4.2, and (2.11)

$$\begin{aligned} \delta_h^n &= J_h \tilde{x}^{n+1} - e^{-\tau S_h} J_h \tilde{x}^n - \tau \sum_{i=1}^s b_i(-\tau S_h) \tilde{g}_h^{n+c_i} \\ &= \tau \left( \int_0^1 e^{-(1-s)\tau S_h} \mathcal{L}_h^* \tilde{g}_h^{n+s} ds - \sum_{i=1}^s b_i(-\tau S_h) \tilde{g}_h^{n+c_i} \right) + E_{\text{HG}}^n(\tau) \\ &= \tau \left( \int_0^1 e^{-(1-s)\tau S_h} \mathcal{L}_h^* \tilde{g}_h^{n+s} ds - \sum_{i=1}^s b_i(-\tau S_h) \mathcal{L}_h^* \tilde{g}_h^{n+c_i} \right) \\ &\quad + \tau \sum_{i=1}^s b_i(-\tau S_h) r_h(t_{n+c_i}, \tilde{x}^{n+c_i}) + E_{\text{HG}}^n(\tau). \end{aligned}$$

The last two terms are collected in  $\delta_{h,\text{sp}}^n$  and satisfy (4.12). It remains to study the first difference. Applying Taylor expansion on  $\tilde{g}_h^{n+s}$  and  $\tilde{g}_h^{n+c_i}$ , we derive the desired representation and the bounds on the defects follow from Assumption 3.1.  $\square$

From this we derive the second main result.

*Proof of Theorem 3.4.* For the sake of readability, we only treat the case of a globally Lipschitz function. The local version is obtained by an induction argument.

(a) We first consider the case  $p = 2$ . As in [18, Sec. 4.3] we derive from (4.11)

$$\begin{aligned} \|e_h^{n+1}\|_{X_h} &\leq C\tau \sum_{j=0}^n \|e_h^j\|_{X_h} + C \sum_{j=0}^n \left( \|\delta_{h,\text{sp}}^j\|_{X_h} + \|\delta_{h,\tau}^j\|_{X_h} \right) \\ &\quad + C\tau \sum_{j=0}^n \left( \max_{i=1,\dots,s} \|\Delta_{h,\text{sp}}^{j,i}\|_{X_h} + \max_{i=1,\dots,s} \|\Delta_{h,\tau}^{j,i}\|_{X_h} \right), \end{aligned}$$

where we used Assumption 2.4 and the order conditions in Table 1 and Lemma 4.4 with  $p = 2$ . The bounds in (4.12) and a discrete Gronwall lemma yield the result.

(b) We now turn to the case  $p \in \{3, 4\}$ . We follow the steps of [18, Thm. 4.7] and explain the necessary modifications.

(1) Using Assumption 2.5, we obtain the discrete analogue to [18, Lem. 4.4]

$$(4.13) \quad \begin{aligned} g_h(t_{n+c_j}, X_h^{nj}) - \tilde{g}_h^{n+c_j} &= \mathcal{D}_h^n E_h^{ni} + c_i \tau \mathcal{K}_h^n E_h^{ni} + Q_h^{ni}, \\ g_h(t_n, x^n) - \tilde{g}_h^n &= \mathcal{D}_h^n e_h^n + Q_h^n, \end{aligned}$$

with the operators

$$\mathcal{D}_h^n := \frac{\partial g_h}{\partial x_h}(t_n, J_h \tilde{x}^n), \quad \mathcal{K}_h^n := \frac{\partial^2 g_h}{\partial t \partial x_h}(t_n, J_h \tilde{x}^n) + \frac{\partial^2 g_h}{\partial x_h^2}(t_n, J_h \tilde{x}^n) J_h x'(t_n).$$

In addition, there is a constant  $C > 0$  independent of  $\tau$  and  $h$  such that

$$\|Q_h^{ni}\|_{X_h} \leq C(\tau^2 + \|E_h^{ni}\|_{X_h}) \|E_h^{ni}\|_{X_h}, \quad \|Q_h^n\|_{X_h} \leq C \|e_h^n\|_{X_h}^2.$$

(2) We proceed as in [18, Lem. 4.5 and 4.6]. Using (4.13), we inductively eliminate the inner stages in (4.11). Employing the order conditions, which are satisfied up to order  $p$ , and Lemma 4.4, leads to the representation of the global error

$$(4.14) \quad e_h^{n+1} = e^{-\tau S_h} e_h^n + \tau \mathcal{N}_h^n(e_h^n) e_h^n + \tau(\tau^p + \varepsilon_h^n) \mathcal{R}_h^n,$$

with uniformly bounded operators

$$\|\mathcal{N}_h^n(e_h^n)\|_{X_h \leftarrow X_h}, \|\mathcal{R}_h^n\|_{X_h} \leq C.$$

(3) Resolving the error recursion (4.14), using (2.5), and the application of a discrete Gronwall lemma gives the result.  $\square$

## 5. APPLICATION TO SECOND-ORDER WAVE-TYPE EQUATIONS

In this section we explain how our results apply to second-order wave-type equations. For this purpose, we first present an abstract spatial discretization of (1.1) which was introduced in [15, 16]. In these references, it was proven that this discretization fits into the setting of Sections 2.3 and 2.4. We shortly recall the relevant results.

**5.1. Abstract space discretization.** Let  $V_h$  be a finite dimensional vector space. We consider the following discretized version of (2.1) on  $(0, T]$

$$(5.1) \quad \begin{aligned} m_h(u_h'', v_h) + b_h(u_h', v_h) + a_h(u_h, v_h) &= m_h(f_h(t, u_h), v_h) \quad \text{for all } v_h \in V_h, \\ u_h(0) &= u_h^0, \quad u_h'(0) = v_h^0. \end{aligned}$$

For the discretized quantities we require similar properties as for their continuous counterparts, cf. Assumption 2.1.

*Assumption 5.1.*

- a) The bilinear form  $a_h: V_h \times V_h \rightarrow \mathbb{R}$  is symmetric and there exists a constant  $\widehat{c}_G \geq 0$  such that

$$\tilde{a}_h := a_h + \widehat{c}_G m_h$$

is a scalar product on  $V_h$  with induced norm  $\|\cdot\|_{\tilde{a}_h}$ .

- b) The bilinear form  $m_h$  is also a scalar product on  $V_h$ . We denote  $V_h$  equipped with this scalar product  $m_h$  by  $H_h$  and the induced norm by  $\|\cdot\|_{m_h}$ .
- c) The bilinear form  $b_h: V_h \times H_h \rightarrow \mathbb{R}$  is bounded independent of  $h$  and there exists a  $\widehat{\beta}_{\text{qm}} \geq 0$  such that

$$b_h(v_h, v_h) + \widehat{\beta}_{\text{qm}} \|v_h\|_{m_h}^2 \geq 0 \quad \text{for all } v_h \in V_h.$$

- d) The nonlinearity  $f_h: [0, T] \times V_h \rightarrow H_h$  is locally Lipschitz-continuous on  $V_h$  with constant  $\widehat{L}_{T, M}$ .
- e) There exists a constant  $\widehat{C}_{H, V} > 0$  such that  $\|v_h\|_{m_h} \leq \widehat{C}_{H, V} \|v_h\|_{\tilde{a}_h}$  for all  $v_h \in V_h$ .

All constants in this assumption are independent of  $h$ .

Similar to the continuous case, we reformulate (5.1) as an evolution equation. For this purpose, we define  $A_h, B_h \in \mathcal{L}(V_h; V_h)$  via

$$m_h(A_h v_h, \varphi_h) = a_h(v_h, \varphi_h), \quad m_h(B_h v_h, \varphi_h) = b_h(v_h, \varphi_h) \quad \text{for all } v_h, \varphi_h \in V_h.$$

Then, (5.1) is equivalent to

$$(5.2) \quad \begin{aligned} u_h''(t) + B_h u_h'(t) + A_h u_h(t) &= f_h(t, u_h(t)), \quad t \in (0, T], \\ u_h(0) &= u_h^0, \quad u_h'(0) = v_h^0. \end{aligned}$$

**5.2. Framework for error analysis.** Similar to the first-order case we assume the existence of certain operators that are necessary for the error analysis.

*Assumption 5.2.*

- a) There exists a lift operator  $\mathcal{L}_h^V \in \mathcal{L}(V_h; V)$  satisfying

$$(5.3) \quad \|\mathcal{L}_h^V v_h\|_m \leq \widehat{C}_H \|v_h\|_{m_h}, \quad \|\mathcal{L}_h^V v_h\|_{\tilde{a}} \leq \widehat{C}_V \|v_h\|_{\tilde{a}_h}$$

for all  $v_h \in V_h$  with constants  $\widehat{C}_H, \widehat{C}_V > 0$  independent of  $h$ .

- b) There exists an interpolation operator  $I_h: Z^V \rightarrow H_h$  defined on a dense subspace  $Z^V$  of  $V$  with

$$(5.4) \quad \|I_h z\|_{m_h} \leq \widehat{C}_{I_h} \|z\|_{Z^V} \quad \text{for all } z \in Z^V$$

where the constant  $\widehat{C}_{I_h} > 0$  is independent of  $h$ .

Under Assumption 5.2, we can define the following operators and quantities that will appear in the latter error bounds.

**Definition 5.3.**

- a) The adjoints of the lift operator  $\mathcal{L}_h^{V*}: V \rightarrow V_h$  and  $\mathcal{L}_h^{H*}: H \rightarrow H_h$  w.r.t. the scalar products in  $V$  and  $H$  are defined via

$$\begin{aligned} m_h(\mathcal{L}_h^{H*} v, w_h) &= m(v, \mathcal{L}_h^V w_h) \quad \text{for all } v \in H, w_h \in H_h, \\ \tilde{a}_h(\mathcal{L}_h^{V*} v, w_h) &= \tilde{a}(v, \mathcal{L}_h^V w_h) \quad \text{for all } v \in V, w_h \in V_h. \end{aligned}$$

- b) Our error bounds rely on the errors in the scalar products that are defined for  $v_h, w_h \in V_h$  via

$$\begin{aligned}\Delta m(v_h, w_h) &:= m(\mathcal{L}_h^V v_h, \mathcal{L}_h^V w_h) - m_h(v_h, w_h), \\ \Delta \tilde{a}(v_h, w_h) &:= \tilde{a}(\mathcal{L}_h^V v_h, \mathcal{L}_h^V w_h) - \tilde{a}_h(v_h, w_h).\end{aligned}$$

**5.3. Fully discrete error bounds.** In this section we explain how the error bounds from Section 3 translate to the second-order equations considered in this section. In the same way as in (2.2), defining  $X_h = V_h \times H_h$  and

$$x_h(t) = \begin{bmatrix} u_h(t) \\ v_h(t) \end{bmatrix}, \quad S_h = \begin{bmatrix} 0 & -\mathbf{I} \\ A_h & B_h \end{bmatrix}, \quad g_h(t, x_h) = \begin{bmatrix} 0 \\ f_h(t, u_h) \end{bmatrix}, \quad x_h^0 = \begin{bmatrix} u_h^0 \\ v_h^0 \end{bmatrix},$$

the second-order equation (5.2) is equivalent to (2.7).

We now recall the results from [16], which show that the first-order formulations of (1.1) as well as (5.2), respectively, fit in the setting of Section 2. In [16, Sec. 3.2], it was shown that Assumption 5.1 implies Assumption 2.4. We define the first-order reference operator  $J_h: Z \rightarrow X_h$  via

$$J_h \begin{bmatrix} v \\ w \end{bmatrix} := \begin{bmatrix} \mathcal{L}_h^{V*} v \\ I_h w \end{bmatrix}$$

on  $Z = V \times Z^V \xrightarrow{d} X$ . Using this definition, it was shown in the proof of [16, Thm. 3.9] that Assumption 5.2 implies Assumption 2.6 and, that for all  $t \in [0, T]$  the error term  $E_h(t)$  defined in (2.12) is bounded by  $E_h(t) \leq \sum_{i=1}^5 E_i$  with

$$\begin{aligned}E_1 &:= \|u_h^0 - \mathcal{L}_h^{V*} u^0\|_{\tilde{a}_h} + \|v_h^0 - I_h v^0\|_{m_h}, \\ E_2 &:= \|\mathcal{L}_h^{H*} f(\cdot, u(\cdot)) - f_h(\cdot, \mathcal{L}_h^{V*} u(\cdot))\|_{L^\infty([0, T]; H_h)}, \\ E_3 &:= \|(\mathbf{I} - \mathcal{L}_h^V I_h)u\|_{L^\infty([0, T]; V)} + \|(\mathbf{I} - \mathcal{L}_h^V I_h)u'\|_{L^\infty([0, T]; V)} \\ &\quad + \|(\mathbf{I} - \mathcal{L}_h^V I_h)u''\|_{L^\infty([0, T]; H)}, \\ (5.5) \quad E_4 &:= \left\| \max_{\|\varphi_h\|_{\tilde{a}_h}=1} \Delta \tilde{a}(I_h u, \varphi_h) \right\|_{L^\infty(0, t)} + \left\| \max_{\|\psi_h\|_{m_h}=1} \Delta m(I_h u, \psi_h) \right\|_{L^\infty(0, t)} \\ &\quad + \left\| \max_{\|\varphi_h\|_{\tilde{a}_h}=1} \Delta \tilde{a}(I_h u', \varphi_h) \right\|_{L^\infty(0, t)} + \left\| \max_{\|\psi_h\|_{m_h}=1} \Delta m(I_h u'', \psi_h) \right\|_{L^\infty(0, t)}, \\ E_5 &:= \left\| \max_{\|\psi_h\|_{m_h}=1} |b(u', \mathcal{L}_h^V \psi_h) - b_h(I_h u', \psi_h)| \right\|_{L^\infty(0, t)}.\end{aligned}$$

Thus, the consistency Assumption 2.8 is implied by the following condition.

*Assumption 5.4.* The solution of (1.1) satisfies  $u \in C^2([0, T]; Z^V)$  and the error terms in (5.5) converge to zero, i.e.,

$$E_i \rightarrow 0, \quad \text{for } h \rightarrow 0, \quad i = 1, \dots, 5.$$

It remains to transfer Assumptions 2.5 and 3.1 into the second-order formulation. The Fréchet differentiability is easily obtained from the following reformulation.

*Assumption 5.5.* The nonlinearity  $f_h$  satisfies  $f_h \in C^3([0, T] \times V_h; H_h)$  with derivatives bounded independent of  $h$ .

Analogously, Assumption 3.1 takes the following form.

*Assumption 5.6.* Let  $m \geq 1$  and  $u$  be the exact solution of (1.1). The differentiability

$$s \mapsto f(s, x(s)) \in C^m([0, T]; H)$$

holds.

Employing the above assumptions, we rephrase the main theorems of Section 3 for the second-order wave equation (1.1).

**Corollary 5.7.** *Let Assumptions 2.1, 5.1, 5.2, 5.4, and 5.6 be satisfied. Then, Theorems 3.2 and 3.4 (for  $p = 2$ ) hold true for the first-order formulation of the second-order wave equation (1.1) with corresponding spatial discretization (5.2). If additionally Assumption 5.5 is satisfied, then Theorem 3.4 holds also true for  $p = 3$  and  $p = 4$ . In particular, we obtain the following two error bounds.*

a) *For the  $k$ -step Adams method (3.5), it holds for  $t_k \leq t_n \leq T$*

$$\begin{aligned} \|\mathcal{L}_h^V u_h^n - u(t_n)\|_{\bar{a}} + \|\mathcal{L}_h^V v_h^n - v(t_n)\|_m &\leq C(\tau^k + \sum_{i=1}^5 E_i) \\ &+ C \sum_{j=1}^{k-1} \left( \|u_h^j - \mathcal{L}_h^{V*} u(t_j)\|_{\bar{a}_h} + \|v_h^j - I_h v(t_j)\|_{m_h} \right). \end{aligned}$$

b) *For an exponential Runge–Kutta method (3.7) of order  $p$ , it holds for  $t_n \leq T$*

$$\|\mathcal{L}_h^V u_h^n - u(t_n)\|_{\bar{a}} + \|\mathcal{L}_h^V v_h^n - v(t_n)\|_m \leq C(\tau^p + \sum_{i=1}^5 E_i).$$

We emphasize that this result immediately implies full discretization error bounds for concrete examples if the space discretization admits suitable bounds on the errors  $E_i$ . An application is given in the following section.

## 6. APPLICATION: WAVE EQUATION WITH KINETIC BOUNDARY CONDITIONS

In this section, we apply the abstract theory of Section 5 to a semilinear wave equation with kinetic boundary conditions. By this example we illustrate our results and perform some numerical experiments.

**6.1. Continuous equation.** Let  $\Omega = B_1(0) \subset \mathbb{R}^2$  be the two-dimensional unit sphere with boundary  $\Gamma = \partial\Omega$  and corresponding outer normal vector  $\mathbf{n}$ . By  $\Delta_\Gamma$  we denote the Laplace–Beltrami operator on  $\Gamma$  and consider the wave equation with kinetic boundary conditions and nonlinear forcing terms given by

$$(6.1a) \quad u_{tt} + (1 + \mathbf{x} \cdot \nabla)u_t - \Delta u = |u|u + \eta_\Omega(t, \mathbf{x}), \quad \text{for } t \geq 0, \mathbf{x} \in \Omega,$$

$$(6.1b) \quad u_{tt} + \partial_{\mathbf{n}} u - \Delta_\Gamma u = u^3 + \eta_\Gamma(t, \mathbf{x}), \quad \text{for } t \geq 0, \mathbf{x} \in \Gamma,$$

$$(6.1c) \quad u(0, \mathbf{x}) = u^0(\mathbf{x}), \quad u_t(0, \mathbf{x}) = v^0(\mathbf{x}), \quad \text{in } \bar{\Omega}$$

with inhomogeneities  $\eta_\Omega, \eta_\Gamma$ . In [16, Sec. 2.1], it was shown that a more general version of this equation fits into the setting of Section 2.1 with

$$\begin{aligned} H &= L^2(\Omega) \times L^2(\Gamma), \quad V = H^1(\Omega; \Gamma), \quad Z = D(A) = H^2(\Omega; \Gamma), \\ A &= \begin{bmatrix} -\Delta & 0 \\ \partial_{\mathbf{n}} & -\Delta_\Gamma \end{bmatrix}, \quad B = \begin{bmatrix} (1 + \mathbf{x} \cdot \nabla) & 0 \\ 0 & 0 \end{bmatrix}, \quad f(t, u) = \begin{bmatrix} |u|u + \eta_\Omega(t, \mathbf{x}) \\ (\gamma u)^3 + \eta_\Gamma(t, \mathbf{x}) \end{bmatrix}, \end{aligned}$$

where

$$H^k(\Omega; \Gamma) := \{v \in H^k(\Omega) \mid \gamma(v) \in H^k(\Gamma)\} \subset H^k(\Omega) \times H^k(\Gamma)$$

and  $\gamma: H^1(\Omega) \rightarrow L^2(\Gamma)$  denotes the trace operator.

**6.2. Discretization.** To discretize (6.1) in space we use the bulk-surface finite element method as presented in [16, Sec. 2.2]. We denote the maximal mesh width by  $h$ . This discretization is nonconforming since the computational domain  $\Omega_h$  does not coincide exactly with  $\Omega$ . Further, in [16, Sec. 2.2] it was proven, that the discretization fits into the setting of Section 5. Moreover, in the proof of [16, Thm. 2.7] it was shown that, for finite elements of order  $q$ , the space discretization error terms from (5.5) are bounded by

$$(6.2) \quad E_i \leq Ch^q.$$

*Remark 6.1.* Note that compared to [16, Sec. 2.2] the only additional assumptions that we require in Section 5 in order to prove the full discretization bounds are Assumptions 5.5 and 5.6 which correspond to Assumptions 2.5 and 3.1 in the first-order formulation.

- a) The proof of the Lipschitz continuity of the discretized nonlinearity  $f_h$  in [16, Lem. 2.6] is non-trivial and relies on discrete norms that are equivalent to the  $L^q$  norms on the finite elements space. This is necessary to prove that the local Lipschitz constant is independent of  $h$ . Similarly, one can prove the differentiability of  $f_h$ , and bounds for the derivatives independent of  $h$  required in Assumption 5.5. In fact, we can also treat more general nonlinearities in (6.1) if the underlying function and its derivatives satisfy certain growth conditions.
- b) Assumption 5.6 is a regularity assumption on the exact solution and was already necessary to prove convergence of the time discretization, cf. [7, 18, 20]. The exact solution in our numerical experiments satisfies this assumption.

**6.3. Numerical results.** We implemented the numerical experiments in C++ using the finite element library `deal.II` (version 9.2) [3, 4]. The codes to reproduce the experiments are available at <https://doi.org/10.5445/IR/1000134973>.

The finite element discretization was implemented as described in [25, Ch. 6.5.1]. For the time discretization, we consider the exponential Euler scheme (3.4), the exponential two-step Adams method (3.5) (for  $k = 2$ ), and the exponential trapezoidal rule of order  $p = 2$  given by (3.9) with  $c_2 = 1$ . To evaluate the matrix functions appearing in the exponential integrators, we used a rational Krylov method including mass matrices as described in [21, Alg. 2].

For our computations, we set

$$\begin{aligned} u^0(\mathbf{x}) &= 0, & v^0(\mathbf{x}) &= 2\pi\mathbf{x}_1\mathbf{x}_2, \\ \eta_\Omega(t, \mathbf{x}) &= -(4\pi^2 + |\sin(2\pi t)\mathbf{x}_1\mathbf{x}_2|) \sin(2\pi t)\mathbf{x}_1\mathbf{x}_2 + 6\pi \cos(2\pi t)\mathbf{x}_1\mathbf{x}_2, \\ \eta_\Gamma(t, \mathbf{x}) &= -4\pi^2 \sin(2\pi t)\mathbf{x}_1\mathbf{x}_2 + 6 \sin(2\pi t)\mathbf{x}_1\mathbf{x}_2 - (\sin(2\pi t)\mathbf{x}_1\mathbf{x}_2)^3. \end{aligned}$$

Then, the exact solution of (6.1) is given by

$$u(t, \mathbf{x}) = \sin(2\pi t)\mathbf{x}_1\mathbf{x}_2.$$

Since the computation of the lift of a finite element function is very laborious, we do not compute the full error  $\mathcal{L}_h^V u_h^n - u(t_n)$ . Instead, in our numerical examples




FIGURE 2. Error  $\mathbf{E}(0.8)$  of the exponential trapezoidal rule (with time step size  $\tau = 5.33 \cdot 10^{-4}$ ) combined with finite elements of order  $q = 1$  and  $q = 2$  plotted against the mesh width  $h$ .

we consider the error

$$\mathbf{E}(t) := \|u_h(t) - u(t)|_{\Omega_h}\|_{H^1(\Omega_h; \Gamma_h)} + \|u'_h(t) - u'(t)|_{\Omega_h}\|_{L^2(\Omega_h) \times L^2(\Gamma_h)},$$

cf. [25, Ch. 6] for more details. We evaluate the integrals appearing in  $\mathbf{E}$  with a quadrature rule of higher order, such that the quadrature error is negligible. The restriction of  $u$  to  $\Omega_h$  is possible since in our case we have  $\Omega_h \subset \Omega$ .

In Figure 2 the convergence of the error w.r.t. the spatial mesh width  $h$  is shown when using the exponential trapezoidal rule. We observe that for finite elements of order  $q$  the error converges with order  $q$  in space for  $q = 1, 2$  as predicted by Corollary 5.7 combined with (6.2).

In Figure 3, we consider the convergence of the error w.r.t. the time step size  $\tau$ . For all three time discretization schemes we observe the order predicted by Theorems 3.2 and 3.4, respectively, until the error is dominated by the error of the spatial approximation.

#### ACKNOWLEDGMENTS

We thank our colleagues Constantin Carle and Marlis Hochbruck for their careful reading of this manuscript.

#### REFERENCES

- [1] I. Alonso-Mallo, B. Cano, and N. Reguera, *Avoiding order reduction when integrating linear initial boundary value problems with exponential splitting methods*, IMA J. Numer. Anal. **38** (2018), no. 3, 1294–1323. MR3829162
- [2] ———, *Avoiding order reduction when integrating reaction-diffusion boundary value problems with exponential splitting methods*, J. Comput. Appl. Math. **357** (2019), 228–250. MR3924553
- [3] D. Arndt, W. Bangerth, B. Blais, T. C. Clevenger, M. Fehling, A.V. Grayver, T. Heister, L. Heltai, M. Kronbichler, M. Maier, P. Munch, J.-L. Pelteret, R. Rastak, I. Thomas, B. Turcksin, Z. Wang, and D. Wells, *The deal.II library, Version 9.2*, J. Numer. Math. **28** (2020), no. 3, 131–146. MR4151106
- [4] W. Bangerth, R. Hartmann, and G. Kanschat, *deal.II—a general-purpose object-oriented finite element library*, ACM Trans. Math. Software **33** (2007), no. 4, Art. 24, 27. MR2404402
- [5] M. Cai, S. Gan, and X. Wang, *Weak convergence rates for an explicit full-discretization of stochastic Allen-Cahn equation with additive noise*, J. Sci. Comput. **86** (2021), no. 3, Paper No. 34, 30. MR4200962




FIGURE 3. Error  $\mathbf{E}(0.8)$  of the exponential Euler, the exponential trapezoidal, and the two-step exponential Adams scheme combined with finite elements of order  $q = 1$  and  $q = 2$  ( $h \approx 0.014$ ) plotted against the time step size  $\tau$ .

- [6] N. Caliari, L. Einkemmer, A. Moriggl, and A. Ostermann, *An accurate and time-parallel rational exponential integrator for hyperbolic and oscillatory PDEs*, J. Comput. Phys. **437** (2021), 110289. MR4242415
- [7] M. P. Calvo and C. Palencia, *A class of explicit multistep exponential integrators for semi-linear problems*, Numer. Math. **102** (2006), no. 3, 367–381. MR2207266
- [8] E. Celledoni, D. Cohen, and B. Owren, *Symmetric exponential integrators with an application to the cubic Schrödinger equation*, Found. Comput. Math. **8** (2008), no. 3, 303–317. MR2413146
- [9] Z. Chen, S. Gan, and X. Wang, *A full-discrete exponential Euler approximation of the invariant measure for parabolic stochastic partial differential equations*, Appl. Numer. Math. **157** (2020), 135–158. MR4113805
- [10] G. Dimarco and L. Pareschi, *Exponential Runge-Kutta methods for stiff kinetic equations*, SIAM J. Numer. Anal. **49** (2011), no. 5, 2057–2077. MR2861709
- [11] B. L. Ehle and J. D. Lawson, *Generalized Runge-Kutta processes for stiff initial-value problems*, J. Inst. Math. Appl. **16** (1975), no. 1, 11–21. MR391524
- [12] A. Friedli, *Verallgemeinerte Runge-Kutta Verfahren zur Lösung steifer Differentialgleichungssysteme*, Numerical treatment of differential equations (Proc. Conf., Math. Forschungsinst., Oberwolfach, 1976), 1978, pp. 35–50. Lecture Notes in Math., Vol. 631. MR0494950
- [13] L. Gauckler, J. Lu, J. L. Marzuola, F. Rousset, and K. Schratz, *Trigonometric integrators for quasilinear wave equations*, Math. Comp. **88** (2019), no. 316, 717–749. MR3882282
- [14] G. R. Goldstein, *Derivation and physical interpretation of general boundary conditions*, Adv. Differential Equations **11** (2006), no. 4, 457–480. MR2215623
- [15] D. Hipp, M. Hochbruck, and C. Stohrer, *Unified error analysis for nonconforming space discretizations of wave-type equations*, IMA J. Numer. Anal. **39** (2019), no. 3, 1206–1245. MR3984056
- [16] M. Hochbruck and J. Leibold, *Finite element discretization of semilinear acoustic wave equations with kinetic boundary conditions*, Electron. Trans. Numer. Anal. **53** (2020), 522–540. MR4137273

- [17] M. Hochbruck, C. Lubich, and H. Selhofer, *Exponential integrators for large systems of differential equations*, SIAM J. Sci. Comput. **19** (1998), no. 5, 1552–1574. MR1618808
- [18] M. Hochbruck and A. Ostermann, *Explicit exponential Runge-Kutta methods for semilinear parabolic problems*, SIAM J. Numer. Anal. **43** (2005), no. 3, 1069–1090. MR2177796
- [19] ———, *Exponential integrators*, Acta Numer. **19** (2010), 209–286. MR2652783
- [20] ———, *Exponential multistep methods of Adams-type*, BIT **51** (2011), no. 4, 889–908. MR2855432
- [21] M. Hochbruck, T. Pažur, A. Schulz, E. Thawinan, and C. Wieners, *Efficient time integration for discontinuous Galerkin approximations of linear wave equations [Plenary lecture presented at the 83rd Annual GAMM Conference, Darmstadt, 26th–30th March, 2012]*, ZAMM Z. Angew. Math. Mech. **95** (2015), no. 3, 237–259. MR3324777
- [22] A.-K. Kassam and L. N. Trefethen, *Fourth-order time-stepping for stiff PDEs*, SIAM J. Sci. Comput. **26** (2005), no. 4, 1214–1233 (electronic). MR2143482
- [23] S. Krogstad, *Generalized integrating factor methods for stiff PDEs*, J. Comput. Phys. **203** (2005), no. 1, 72–88. MR2104391
- [24] J. D. Lawson, *Generalized Runge-Kutta processes for stable systems with large Lipschitz constants*, SIAM J. Numer. Anal. **4** (1967), 372–380. MR221759
- [25] J. Leibold, *A unified error analysis for the numerical solution of nonlinear wave-type equations with application to kinetic boundary conditions*, Ph.D. Thesis, 2021. <https://doi.org/10.5445/IR/1000130222>.
- [26] V. T. Luan and D. L. Michels, *Efficient exponential time integration for simulating nonlinear coupled oscillators*, J. Comput. Appl. Math. **391** (2021), 113429, 18. MR4215003
- [27] V. T. Luan and A. Ostermann, *Explicit exponential Runge-Kutta methods of high order for parabolic problems*, J. Comput. Appl. Math. **256** (2014), 168–179. MR3095695
- [28] D. L. Michels and M. Desbrun, *A semi-analytical approach to molecular dynamics*, J. Comput. Phys. **303** (2015), 336–354. MR3422717
- [29] S. P. Nørsett, *An A-stable modification of the Adams-Bashforth methods*, Conf. on Numerical Solution of Differential Equations (Dundee, 1969), 1969, pp. 214–219. MR0267771
- [30] A. Pazy, *Semigroups of linear operators and applications to partial differential equations*, Applied Mathematical Sciences, vol. 44, Springer-Verlag, New York, 1983. MR710486
- [31] M. Pototschnig, J. Niegemann, L. Tkeshelashvili, and K. Busch, *Time-domain simulations of the nonlinear Maxwell equations using operator-exponential methods*, IEEE Trans. Antennas and Propagation **57** (2009), no. 2, 475–483. MR2517334
- [32] K. Strehmel and R. Weiner, *Linear-implizite Runge-Kutta-Methoden und ihre Anwendung*, Teubner-Texte zur Mathematik [Teubner Texts in Mathematics], vol. 127, B. G. Teubner Verlagsgesellschaft mbH, Stuttgart, 1992. With English, French and Russian summaries. MR1171219
- [33] E. Vitillaro, *Strong solutions for the wave equation with a kinetic boundary condition*, Recent trends in nonlinear partial differential equations. I. Evolution problems, 2013, pp. 295–307. MR3155916
- [34] ———, *On the wave equation with hyperbolic dynamical boundary conditions, interior and boundary damping and source*, Arch. Ration. Mech. Anal. **223** (2017), no. 3, 1183–1237. MR3594355

INSTITUTE FOR APPLIED AND NUMERICAL MATHEMATICS, KARLSRUHE INSTITUTE OF TECHNOLOGY, 76131 KARLSRUHE, GERMANY

Email address: benjamin.doerich@kit.edu, jan.leibold@kit.edu