


On the well-posedness of magnetic Schrödinger equations with unbounded potentials

Dorothee Frey, Siliang Weng

CRC Preprint 2026/10, March 2026

KARLSRUHE INSTITUTE OF TECHNOLOGY

CRC 1173



Wave
phenomena

Participating universities



Funded by



ON THE WELL-POSEDNESS OF MAGNETIC SCHRÖDINGER EQUATIONS WITH UNBOUNDED POTENTIALS

DOROTHEE FREY AND SILIANG WENG

ABSTRACT. We consider magnetic Schrödinger equations with sublinear magnetic potentials and subquadratic electric potentials on \mathbb{R}^d , as well as generalizations thereof. We obtain new results on the global well-posedness of the Cauchy problem with initial data in magnetic modulation spaces $M_A^p(\mathbb{R}^d)$. Our results are achieved by approximating the solution in phase space using the magnetic Hamiltonian flow. This method includes the potentials as part of the generalized Schrödinger operator instead of treating them as perturbations, and thereby allows us to deal with unbounded potentials. For $A \equiv 0$, the space $M_A^p(\mathbb{R}^d)$ reduces to the usual modulation space $M^p(\mathbb{R}^d)$, for which relevant known results for the usual Schrödinger equation can be recovered.

1. INTRODUCTION

Let $d \geq 2$. In this article, we study the global well-posedness of the Cauchy problem for the Schrödinger equation with unbounded electromagnetic potential in $\mathbb{R} \times \mathbb{R}^d$,

$$(1) \quad i\partial_t u(t; y) = (H^A + V(t))u(t; y), \quad u(0) = u_0$$

on magnetic modulation spaces $M_A^p(\mathbb{R}^d)$ as defined below. Here H^A denotes the magnetic Laplacian with the magnetic vector potential A , that is

$$H^A := \frac{1}{2} \sum_{j=1}^d (-i\partial_j - A_j(y))^2, \quad \partial_k A_j - \partial_j A_k = B_{jk}, \quad 1 \leq j, k \leq d,$$

and $V(t) \in C^\infty(\mathbb{R}^d, \mathbb{R})$ denotes the electric scalar potential. We define the magnetic field B to be a smooth, closed 2-form

$$B(y) := \sum_{1 \leq j, k \leq d} B_{jk}(y) dy_j \wedge dy_k$$

Date: March 20, 2026.

Key words and phrases. Schrödinger equation, modulation spaces, magnetic pseudodifferential operators, unbounded potentials, phase space transforms, magnetic Gaussian wavepackets.

Funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 258734477 – SFB 1173. The second author wishes to thank H. Cornean for very helpful discussions.

with $B_{jk} = -B_{kj} \in C^\infty(\mathbb{R}^d, \mathbb{R})$, and a magnetic vector potential A associated to B is a vector-valued function $A = (A_1, A_2, \dots, A_d)$ satisfying $dA = B$. We consider unbounded potentials A with at most linear growth and V with at most quadratic growth, that is for multi-indices α and β , the potentials satisfy

$$|\partial^\alpha A(y)| \leq C_{A,\alpha}, \quad |\partial_y^\beta V(t; y)| \leq C_{V,\beta}, \quad |\alpha|, |\beta| \geq 2.$$

For differential operators with long-range magnetic potentials, a pseudo-differential calculus adapted to the magnetic potential has been developed by Iftimie, Măntoiu, and Purice in [12], [8]. The theory extends the usual Weyl calculus to the magnetic Weyl calculus $h \mapsto \text{Op}^A(h)$, mapping a symbol $h \in \mathcal{S}'(\mathbb{R}^{2d})$ to a magnetic pseudo-differential operator (magnetic Ψ DO) $\text{Op}^A(h)$, acting on $u \in \mathcal{S}(\mathbb{R}^d)$ as

$$(2) \quad \text{Op}^A(h)u(y) := (2\pi)^{-d} \int_{\mathbb{R}^d \times \mathbb{R}^d} e^{i(y-y') \cdot \eta} e^{-i\varphi^A(y', y)} h\left(\frac{y+y'}{2}, \eta\right) u(y') dy' d\eta.$$

The precise meaning of the above expression will be recalled in the next section. We note here that this adapted formulation differs from the usual Weyl calculus only by a magnetic-dependent phase term $e^{-i\varphi^A(y', y)}$, which has the property

$$(3) \quad e^{-i\varphi^A(y', y)} \Big|_{A=0} \equiv 1.$$

We thus recover the usual Weyl calculus as the special case $A \equiv 0$. With this formulation, the magnetic Schrödinger operator in (1) can be expressed as

$$(4) \quad H^A + V = \text{Op}^A(h), \quad h(t; y, \eta) = |\eta|^2 + V(t; y).$$

We show under suitable conditions, the Cauchy problems associated to a family of equations that contains (1)

$$(5) \quad i\partial_t u(t, y) = \text{Op}^A(h)u(t; y), \quad u(0) = u_0$$

is well-posed on the magnetic modulation space $M_A^p(\mathbb{R}^d)$, for $1 \leq p \leq \infty$. Here the space $M_A^p(\mathbb{R}^d)$ can be seen as the A -adapted version of the usual modulation space, defined through the magnetic wavepacket transform \mathcal{T}^A for $u \in \mathcal{S}'(\mathbb{R}^d)$

$$\mathcal{T}^A u(x, \xi) := \int e^{-i\xi \cdot (y-x)} e^{-i\varphi^A(y, x)} g(y-x) u(y) dy, \quad \|u\|_{M_A^p(\mathbb{R}^d)} := \|\mathcal{T}^A u\|_{L^p(\mathbb{R}^{2d})},$$

where g denotes the normalized Gaussian function on \mathbb{R}^d (a window function). We also note, although the space M_A^p is formulated explicitly with the magnetic potential A , it in fact intrinsically depends only on the magnetic field B . In other words, M_A^p is gauge-independent, that is for another magnetic potential A' with $dA' = dA = B$, we have $M_A^p \simeq M_{A'}^p$ isometrically.

This choice of space is inspired by the well-posedness of the usual Schrödinger equation in modulation spaces $M^{p,q}$, see [1], [2] and the references therein. Also due to the phase term $e^{-i\varphi^A}$ and its property (3), the magnetic modulation space M_A^p naturally extends the usual modulation space $M^{p,p}$, and is well-behaved interacting with the adapted magnetic Ψ DOs (2). One of the early ideas of defining magnetic versions of modulation spaces can be found in Măntoiu-Purice [13], where they considered the case $p = 2$. To the best of our knowledge, the definition and discussion of M_A^p and its weighted version $M_A^{m,p}$ in this article is new, and so are its applications to magnetic Schrödinger equations.

Previously well-posedness results for the Cauchy problem of (1) with unbounded potentials are mostly obtained on L^2 -based spaces. One foundational work is Yajima [16], where unbounded, non-smooth potentials with t -dependence are considered. More generally, Doi in [6] showed under certain conditions the variable-coefficient version of (1) with unbounded potentials is also well-posed in L^2 -based weighted Sobolev spaces. On the other hand, results for L^p -based spaces with $p \neq 2$ seem out of reach, since the non-magnetic Schrödinger group $e^{-it\Delta}$ is already unbounded on L^p for $p \neq 2$. For the non-magnetic $A \equiv 0$ case, the introduction of the modulation space $M^{p,q}$ as a substitute for L^p has been very successful, since the Schrödinger group $e^{-it\Delta}$ has been shown to be bounded on the modulation spaces $M^{p,q}$ [1]. This suggests taking spaces of modulation type as candidates for wellposedness of magnetic Schrödinger equations. In this direction, Keiichi-Muramatsu [10] obtained a-priori estimates on M^p , but there the existence of the solution is still restricted to L^2 .

By introducing the magnetic modulation space M_A^p and the weighted version $M_A^{m,p}$, we obtain a global well-posedness result for (5). Our assumptions on the potentials are the following.

Assumption 1.1. Fix $\epsilon > 0$. We assume the magnetic field B has smooth components $\{B_{jk}\}_{1 \leq j,k \leq d}$, which are all bounded with $\|B_{jk}\|_\infty \leq C_B$ for some $C_B > 0$. Furthermore, we assume the derivatives of B_{jk} satisfy a decay condition, that is for multi-indices α , there exists $C_{\alpha,B} > 0$ such that

$$(6) \quad |\partial^\alpha B_{jk}(y)| \leq C_{\alpha,B} \langle y \rangle^{-1-\epsilon}, \quad |\alpha| \geq 1.$$

For such given magnetic field B , we define the magnetic vector potential A by the transversal gauge

$$(7) \quad A_j(y) := - \sum_{k=1}^d \int_0^1 s y_k B_{jk}(sy) ds, \quad y \in \mathbb{R}^d.$$

For the electric scalar potential V , we allow t -dependence and assume that $V \in C(\mathbb{R}, C^\infty(\mathbb{R}))$, with quadratic growth bounds that is uniform in t ,

$$(8) \quad |\partial_y^\alpha V(t; y)| \leq C_{V, \alpha}, \quad |\alpha| \geq 2, \quad t \in \mathbb{R}.$$

For the magnetic Ψ DO $\text{Op}^A(h)$ generalizing the magnetic Schrödinger operator, we define the symbol class $S^{0, (k)}$, which aligns with the symbol class used in Tataru [14].

Definition 1.2. For $k \in \mathbb{N}$ and multi-indices α, β , a symbol $h \in C^\infty(\mathbb{R}^{2d})$ is in the symbol class $S^{0, (k)}(\mathbb{R}^{2d})$ if it satisfies

$$|\partial_y^\alpha \partial_\eta^\beta h(y, \eta)| \leq C_{\alpha, \beta}, \quad |\alpha| + |\beta| \geq k.$$

We make the following assumption on the symbols.

Assumption 1.3. We assume the time-dependent symbol $h \in C^1(\mathbb{R}, C^\infty(\mathbb{R}^{2d}))$ is real-valued, and $h(t), \dot{h}(t) \in S^{0, (2)}(\mathbb{R}^{2d})$ uniformly in t . Furthermore, we assume h is of second-order in η , that is, for all $t \in \mathbb{R}$ there exists $C > 0$ such that $|h(t; 0, \eta)| \geq C|\eta|^2$.

Remark 1.4. While the second-order part of the assumption guarantees the operator $\text{Op}^A(h)$ to be of magnetic Schrödinger operator type, our result also holds free of this part of the assumption. In fact, for h that is first-order in η , the arguments will be simpler.

The above assumption gives a family of symbols which contains the symbol $h(t; y, \eta) = |\eta|^2 + V(t; y)$ in (4) with V satisfying Assumption 1.1. Our main result is the following.

Theorem 1.5. Let $T > 0$ and $1 \leq p \leq \infty$. For a given magnetic field B satisfying Assumption 1.1 and a real-valued symbol h satisfying Assumption 1.3, the Cauchy problem with initial value $u_0 \in M_A^{2, p}(\mathbb{R}^d)$

$$(9) \quad \begin{cases} i\partial_t u(t, y) = \text{Op}^A(h)u(t, y), \\ u(0, y) = u_0(y), \end{cases}$$

has a unique strong solution $u \in W^{1, 1}([0, T], M_A^p(\mathbb{R}^d))$ with $u(t) \in M_A^{2, p}(\mathbb{R}^d)$ for all $t \in [0, T]$. Moreover, we have the bound

$$\|u(t)\|_{M_A^{2, p}(\mathbb{R}^d)} \leq C_T \|u_0\|_{M_A^{2, p}(\mathbb{R}^d)}.$$

This result also gives a simple corollary.

Corollary 1.6. *Let $T > 0$ and $1 \leq p \leq \infty, s \geq 2$. For the same magnetic Schrödinger operator $\text{Op}^A(h)$ as in Theorem 1.5 and $f \in L^1([0, T], M_A^{s,p}(\mathbb{R}^d))$, the Cauchy problem with initial value $u_0 \in M_A^{2,p}(\mathbb{R}^d)$*

$$\begin{cases} i\partial_t u(t, y) = \text{Op}^A(h)u(t, y) + f(t, y), \\ u(0, y) = u_0(y) \end{cases}$$

has a unique strong solution $u \in W^{1,1}([0, T], M_A^p(\mathbb{R}^d))$ with $u(t) \in M_A^{2,p}(\mathbb{R}^d)$ for all $t \in [0, T]$.

Our approach is also flexible enough to treat the case of time-dependent magnetic potentials $A(t)$, which we explain in Section 8.

Theorem 1.7. *Let $T > 0$ and $1 \leq p \leq \infty$. Let $B(t)$ be a time-dependent magnetic field with components $B_{jk} \in C^1(\mathbb{R}, C^\infty(\mathbb{R}^d))$. For some fixed $\epsilon > 0$, assume for all $t \in \mathbb{R}$, $B_{jk}(t)$ satisfies $\|B(t)\|_\infty + \|\dot{B}(t)\|_\infty \leq C_B$ and the decay condition*

$$(10) \quad |\partial^\alpha B_{jk}(t; y)| + |\partial^\alpha \dot{B}_{jk}(t; y)| \leq C_{\alpha, B} \langle y \rangle^{-1-\epsilon}, \quad |\alpha| \geq 1.$$

The Cauchy problem with initial value $u_0 \in M_{A(0)}^{2,p}(\mathbb{R}^d)$

$$(11) \quad \begin{cases} i\partial_t u(t, y) = \text{Op}^{A(t)}(h)u(t, y), \\ u(0, y) = u_0(y), \end{cases}$$

has a unique strong solution $u \in W^{1,1}([0, T], M_{A(t)}^p(\mathbb{R}^d))$ with $u(t) \in M_{A(t)}^{2,p}(\mathbb{R}^d)$ for all $t \in [0, T]$. Moreover, we have the bound

$$\|u(t)\|_{M_{A(t)}^{2,p}(\mathbb{R}^d)} \leq C_T \|u_0\|_{M_{A(0)}^{2,p}(\mathbb{R}^d)}.$$

With the identification $M_{A(t)}^2 \simeq L^2$, the above result can be seen as a generalization of the classical results in Yajima [16].

Our strategy for the proof is inspired by the work of Tataru [14], Cordero-Gröchenig-Nicola-Rodino [3] and Cordero-Nicola-Rodino [4]. The main idea of the strategy is to use the wavepacket transform \mathcal{T}^A to take a phase space perspective, so that the quantum evolution $u(t)$ satisfying (9) can be related to a classical evolution in the phase space (a magnetic Hamiltonian flow) with initial data $\tilde{u}_0 := \mathcal{T}^A u_0$. The flow property is then exploited through a generalized version of Yajima in [16, Lemma 2.1], which only requires the mild decay assumption on B as in (6). For $A \equiv 0$, our result recovers the modulation

space well-posedness results for the non-magnetic Schrödinger equation with unbounded electric potential V .

2. PRELIMINARIES

2.1. Notation. For the configuration space \mathbb{R}^d , the phase space is the cotangent bundle $T^*\mathbb{R}^d \simeq \mathbb{R}^{2d}$. For differential operators, we use the notation $D := -i\partial$, and for a scalar function f on \mathbb{R}^d , we use $\partial_k f(x) := \partial_{x_k} f(x)$ where $x := (x_1, \dots, x_k, \dots, x_d)$. If f is time-dependent, we also use the shorthand notation $\dot{f} := df/dt$. For scalar products of $x, y \in \mathbb{R}^d$ we use $x \cdot y$ or $\langle x, y \rangle$. For the L^2 pairing we denote $\langle f, g \rangle_{L^2} := \int \bar{f}g$, with the complex conjugation on the left.

We use the Japanese bracket $\langle x \rangle := (1 + |x|^2)^{1/2}$ for $x \in \mathbb{R}^d$, and note Peetre's inequality, for $s \in \mathbb{R}$,

$$\langle x + y \rangle^s \leq 2^s \langle x \rangle^s \langle y \rangle^{|s|}, \quad x, y \in \mathbb{R}^d.$$

For inequalities, we use the notation \lesssim to hide absolute constants, and also the constants that only depend on invariant objects such as the Gaussian g . We use subscripts to denote the dependence of relevant objects or parameters, for example C_B denotes constants that can be determined from a given magnetic field B .

2.2. Magnetic Potential and Flux. Let the magnetic field B with components $B_{jk} \in C^\infty(\mathbb{R}^d, \mathbb{R})$ satisfy $\|\partial^\alpha B_{jk}\|_\infty \leq C_\alpha$, and let $A(\cdot)$ denote the magnetic vector potential obtained through the transversal gauge (7). For fixed $x \in \mathbb{R}^d$, we define the vector potential $A(\cdot, x)$ by

$$A_j(y, x) := - \sum_{k=1}^d \int_0^1 s(y_k - x_k) B_{jk}(x + s(y - x)) ds, \quad y \in \mathbb{R}^d.$$

The above potential satisfies $dA(\cdot, x) = B$ and also recovers $A(y)$ by $A(y, 0) = A(y)$. The vector-valued function $A(\cdot) - A(\cdot, x)$ is related to the desired gauge-covariant property of the magnetic Ψ DOs. Notice we have $dA(\cdot) - dA(\cdot, x) = 0$, so $A(\cdot) - A(\cdot, x)$ is in fact a closed 1-form. We can thus find a function φ^A such that

$$(12) \quad \partial_{y_j} \varphi^A(y, x) = A_j(y) - A_j(y, x), \quad y, x \in \mathbb{R}^d, \quad 1 \leq j \leq d.$$

A straightforward choice is to integrate both sides above from x to y (cf. [5] equation (1.4)),

$$(13) \quad \varphi^A(y, x) := \int_{\Gamma_{x,y}} (A(\cdot) - A(\cdot, x)) = \int_{\Gamma_{x,y}} A, \quad x, y \in \mathbb{R}^d,$$

where $\Gamma_{x,y}$ denotes the line segment connecting x to y . Explicitly, we have

$$\int_{\Gamma_{x,y}} A := \int_{\Gamma_{x,y}} A(\cdot, 0) = (y - x) \cdot \int_0^1 A((1-s)x + sy, 0) ds,$$

and the $\int_{\Gamma_{x,y}} A(\cdot, x)$ term in (13) vanishes, because B is skew-symmetric, so

$$\int_{\Gamma_{x,y}} A(\cdot, x) = \sum_{j,k=1}^d (y_j - x_j) \int_0^1 \int_0^1 \theta s (y_k - x_k) B_{jk}(x + \theta((1-s)x + sy)) d\theta ds = 0.$$

From this construction, we see that φ^A is anti-symmetric in its variables, i.e.

$$(14) \quad \varphi^A(y, x) = -\varphi^A(x, y), \quad x, y \in \mathbb{R}^d.$$

Because the value of φ^A equals integrating the vector field A along a line segment, we denote by $\Gamma^B(y, x, z)$ the flux given by the magnetic field B through the oriented triangle with vertices y, x, z . Using Stokes' theorem, we have

$$(15) \quad \Gamma^B(y, x, z) = \varphi^A(y, x) + \varphi^A(x, z) + \varphi^A(z, y).$$

By [9, Lemma 2.10], we can formulate a matrix $C(y, x, z) := (c_{jk}(y, x, z))_{1 \leq j, k \leq d}$ with

$$c_{jk}(y, x, z) := \int_0^1 \int_0^s B_{jk}(ty + (s-t)x + (1-s)z) dt ds, \quad (y, x, z) \in \mathbb{R}^{3d},$$

it then allows us to re-write Γ^B as

$$\Gamma^B(y, x, z) = \langle C(y, x, z)(y - x), (y - z) \rangle.$$

With the above formulation, the following estimates hold by direct calculation, which bounds the derivatives of the magnetic flux by the length of the line segments $|y - x|$ and $|y - z|$.

Lemma 2.1. *For a magnetic field B with $B_{jk} \in C^\infty(\mathbb{R}^d, \mathbb{R})$ and $\|\partial^\alpha B_{jk}\|_\infty \leq C_\alpha$, the associated magnetic flux Γ^B and phase function $e^{i\Gamma^B}$ satisfy*

$$(16) \quad \begin{aligned} |\Gamma^B(y, x, z)| &\leq C_B |y - x| |y - z|, \\ |\partial^\alpha e^{i\Gamma^B(y, x, z)}| &\leq C_{\alpha, B} (|y - x| + |y - z|)^{|\alpha|}, \quad |\alpha| \geq 1. \end{aligned}$$

3. MAGNETIC PSEUDO-DIFFERENTIAL OPERATOR

In this section, we recall the construction of magnetic pseudo-differential operators (Ψ DOs) with a given magnetic field B satisfying $\sup_{x \in \mathbb{R}^d} |\partial^\alpha B_{jk}(x)| \leq C_\alpha$. With φ^A constructed in (13), we can now give a definition of magnetic Ψ DOs following the construction in [8, Section 1]), for general symbols h in the space of tempered distributions $\mathcal{S}'(\mathbb{R}^{2d})$. Since tempered distributions can grow at most polynomially at infinity, the resulting magnetic Ψ DOs as integral operators are well-defined for Schwartz functions $\mathcal{S}(\mathbb{R}^d)$.

Definition 3.1. Given a symbol $h \in \mathcal{S}'(\mathbb{R}^{2d})$, the magnetic pseudo-differential operator $\text{Op}^A(h) \in \mathcal{L}(\mathcal{S}(\mathbb{R}^d); \mathcal{S}'(\mathbb{R}^d))$ is defined by

$$(17) \quad \text{Op}^A(h)u(y) := (2\pi)^{-d} \int_{\mathbb{R}^{2d}} e^{i(y-y') \cdot \eta} e^{-i\varphi^A(y', y)} h\left(\frac{y+y'}{2}, \eta\right) u(y') \, dy' d\eta, \quad y \in \mathbb{R}^d,$$

where $u \in \mathcal{S}(\mathbb{R}^d)$.

This mapping of symbols to magnetic Ψ DOs $h \mapsto \text{Op}^A(h)$ is also called magnetic Weyl quantization. This formulation has the advantage that the magnetic potential A is fully decoupled from the symbol h . For comparison, the magnetic Schrödinger operator $H^A + V$ in (1) can also be formulated with the usual Ψ DO theory, which requires a much complicated symbol

$$h_A(t; y, \eta) := |\eta - A(y)|^2 + V(t; y)$$

such that

$$H^A + V = \text{Op}(h_A) := (2\pi)^{-d} \int_{\mathbb{R}^{2d}} e^{i(y-y') \cdot \eta} h_A\left(\frac{y+y'}{2}, \eta\right) u(y') \, dy' d\eta.$$

Furthermore, the magnetic Ψ DO formulation enjoys a gauge-covariant property. That is, for $A' = A + dv$ with $v \in C_{\text{pol}}^\infty(\mathbb{R}^d, \mathbb{R})$ a real-valued smooth function of at most polynomial growth, A and A' are both associated to the magnetic field $B = dA' = dA$. Then the magnetic Ψ DOs $\text{Op}^A(h)$ and $\text{Op}^{A'}(h)$ are unitarily equivalent in L^2 (cf. [12, Proposition 3.6]). Thus this formulation depends more intrinsically on the object B , which allows us to formulate Assumption 1.1 in terms of the magnetic field B instead of the magnetic potentials A . In particular, the adapted magnetic modulation space $M_A^p(\mathbb{R}^d)$ we formulate also depends intrinsically on B , which we show in the next section as Proposition 4.5.

Another important feature of the above Weyl quantization is that for real symbols h , the operator $\text{Op}^A(h)$ is symmetric on $\mathcal{S}(\mathbb{R}^d)$, thanks to the balanced choice $\frac{y+y'}{2}$ in (17). But this can also make explicit calculations a bit awkward. To simplify calculations, a slightly different quantization is also useful, that is the magnetic Kohn-Nirenberg quantization

Op_{KN}^A (or the magnetic “left” quantization)

$$\text{Op}_{KN}^A(h)u(y) := (2\pi)^{-d} \int_{\mathbb{R}^{2d}} e^{i(y-y')\cdot\eta} e^{-i\varphi^A(y',y)} h(y,\eta) u(y') dy' d\eta, \quad y \in \mathbb{R}^d.$$

The Weyl quantization can be converted to the Kohn-Nierenberg quantization by modifying the symbol h . [9, Proposition 2.13] provides such a conversion for a symbol class S^m , and the proof there can be used with obvious changes to give a conversion for our symbol class $S^{0,(2)}(\mathbb{R}^{2d})$.

Lemma 3.2. *Given $a \in S^{0,(2)}(\mathbb{R}^{2d})$, there exists a symbol $b \in S^{0,(2)}(\mathbb{R}^{2d})$ such that $\text{Op}^A(a) = \text{Op}_{KN}^A(b)$, with*

$$b(y,\eta) = a(y,\eta) + a_r(y,\eta), \quad a_r \in S^{0,(0)}(\mathbb{R}^{2d}).$$

By [9, Proposition 2.13], we also have an explicit expression for the symbol a_r given by

$$a_r(y,\eta) := \sum_{|\alpha| \geq 1} \frac{1}{|\alpha|!} \left(\frac{1}{2}\right)^{|\alpha|} (D_y^\alpha \partial_\eta^\alpha a)(y,\eta), \quad (y,\eta) \in \mathbb{R}^{2d}.$$

This implies that if the symbol a depends only on y or η , then the two quantizations $\text{Op}^A(a) = \text{Op}_{KN}^A(a)$ coincide, because all cross-derivatives vanish. In particular, the symbols of the basic magnetic differentiation and multiplication operator

$$P_j^A u(y) := (-i\partial_{y_j} - A_j(y))u(y), \quad Q_j u(y) := y_j u(y)$$

coincide in the two quantizations. That is, for $u \in \mathcal{S}(\mathbb{R}^d)$ we have

$$(18) \quad \begin{aligned} P_j^A u(y) &= \text{Op}^A(\eta_j)u(y) = \text{Op}_{KN}^A(\eta_j)u(y); \\ Q_j u(y) &= \text{Op}^A(y_j)u(y) = \text{Op}_{KN}^A(y_j)u(y). \end{aligned}$$

This can be shown with direct calculations using Fourier inversion. We showcase the calculation with $\text{Op}^A(\eta_j)u(y)$, and the other identity is similar. Denote by $\mathcal{F}_{y'}$ and \mathcal{F}_η^{-1} the Fourier transform and inversion regarding the variables y' and η . We have

$$\begin{aligned} \text{Op}^A(\eta_j)u(y) &= \text{Op}_{KN}^A(\eta_j)u(y) \\ &= (2\pi)^{-d} \int_{\mathbb{R}^{2d}} e^{i(y-y')\cdot\eta} e^{-i\varphi^A(y',y)} \eta_j u(y') dy' d\eta \\ &= \mathcal{F}_\eta^{-1} \mathcal{F}_{y'} \left(-i\partial_{y'_j} (e^{-i\varphi^A(\cdot,y)} u) \right) (y) \\ &= (-i\partial_{y_j} - A_j(y))u(y), \end{aligned}$$

where the last equality is based on (12) and $\varphi^A(y,y) = 0$, that is

$$-i\partial_{y'_j} e^{-i\varphi^A(y',y)} \Big|_{y'=y} = -(A_j(y') - A_j(y',y)) e^{-i\varphi^A(y',y)} \Big|_{y'=y} = A_j(y).$$

4. THE MAGNETIC WAVEPACKET TRANSFORM AND MODULATION SPACES

For a given magnetic field B and the associated magnetic potential A , we define a family of magnetic wavepackets using the magnetic phase function φ^A constructed in (13).

Definition 4.1. We denote the L^2 -normalized Gaussian by

$$g(y) := (2\pi)^{-\frac{d}{2}} \pi^{-\frac{d}{4}} e^{-\frac{1}{2}y^2}, \quad y \in \mathbb{R}^d.$$

For fixed $\lambda \geq 1$, the family of λ -rescaled magnetic wavepackets $\{g_{x,\xi}^{A,\lambda}\}_{(x,\xi) \in \mathbb{R}^{2d}}$ are Schwartz functions parameterized by $(x, \xi) \in \mathbb{R}^{2d}$, defined by

$$(19) \quad g_{x,\xi}^{A,\lambda}(y) := \lambda^{\frac{d}{2}} e^{i\xi \cdot (y-x)} g(\lambda(y-x)) e^{i\varphi^A(y,x)}, \quad y \in \mathbb{R}^d.$$

By the above definition, for $A \equiv 0$ and $\lambda = 1$, we recover the standard Gaussian wavepackets $\{g_{x,\xi}\}_{(x,\xi) \in \mathbb{R}^{2d}}$,

$$g_{x,\xi}(y) := e^{i\xi \cdot (y-x)} g(y-x) = \pi^{-\frac{d}{4}} e^{i\xi \cdot (y-x)} e^{-\frac{1}{2}(y-x)^2}, \quad y \in \mathbb{R}^d,$$

which are translated and modulated Gaussians, centered around x in the configuration space and around ξ in the frequency space. Thus, the magnetic wavepacket transform we define below can be seen as a generalization of the usual wavepacket transform (cf. [14], equation (1)).

Definition 4.2. For a tempered distribution $u \in \mathcal{S}'(\mathbb{R}^d)$, its magnetic wavepacket transform $\mathcal{T}_\lambda^A u \in \mathcal{S}'(\mathbb{R}^{2d})$ is obtained by integrating against wavepackets $g_{x,\xi}^{A,\lambda}$ defined in (19), i.e.,

$$\begin{aligned} \mathcal{T}_\lambda^A u(x, \xi) &:= \int_{\mathbb{R}^d} \overline{g_{x,\xi}^{A,\lambda}}(y) u(y) dy \\ &= \int_{\mathbb{R}^d} \lambda^{\frac{d}{2}} e^{-i\xi \cdot (y-x)} g(\lambda(y-x)) e^{-i\varphi^A(y,x)} u(y) dy, \quad (x, \xi) \in \mathbb{R}^{2d}. \end{aligned}$$

This magnetic wavepacket transform \mathcal{T}_λ^A is an example of a phase space transform, since it takes a function (distribution) on the configuration space \mathbb{R}^d to the phase space (the cotangent bundle) $T^*\mathbb{R}^d \simeq \mathbb{R}^{2d}$. We also define a formal adjoint \mathcal{T}_λ^{A*} acting on phase space distributions $\tilde{u} \in \mathcal{S}'(\mathbb{R}^{2d})$ by

$$\mathcal{T}_\lambda^{A*} \tilde{u}(y) := \int_{\mathbb{R}^{2d}} g_{x,\xi}^{A,\lambda}(y) \tilde{u}(x, \xi) dx d\xi.$$

Lemma 4.3. *The magnetic wavepacket transform \mathcal{T}_λ^A and its formal adjoint \mathcal{T}_λ^{A*} have the following properties:*

- (i) If $u \in \mathcal{S}(\mathbb{R}^d)$, then $\mathcal{T}_\lambda^A u \in \mathcal{S}(\mathbb{R}^{2d})$. If $\tilde{u} \in \mathcal{S}(\mathbb{R}^{2d})$ then $\mathcal{T}_\lambda^{A*} \tilde{u} \in \mathcal{S}(\mathbb{R}^d)$.
- (ii) $\mathcal{T}_\lambda^A, \mathcal{T}_\lambda^{A*}$ are L^2 isometries.
- (iii) For $u \in \mathcal{S}(\mathbb{R}^d)$, we have the inversion formula $u = \mathcal{T}_\lambda^{A*} \mathcal{T}_\lambda^A u$.

The first two properties can be checked by direct calculation, or proved similarly as in the non-magnetic case, such as Proposition 1.82 in [2] (also cf. Proposition 3.3 in [13]). The inversion formula can be obtained by

$$\begin{aligned}
\mathcal{T}_\lambda^{A*} \mathcal{T}_\lambda^A u(y) &= \lambda^d \int_{\mathbb{R}^{3d}} e^{i\xi \cdot (y-y')} g(\lambda(y-x)) g(\lambda(y'-x)) e^{i\varphi^A(y,x)} e^{-i\varphi^A(y',x)} u(y') dy' dx d\xi \\
&= \lambda^d (2\pi)^{\frac{d}{2}} \int_{\mathbb{R}^{2d}} \delta(y-y') g(\lambda(y-x)) g(\lambda(y'-x)) e^{i\Gamma^B(y,x,y')} e^{-i\varphi^A(y',y)} u(y') dy' dx \\
&= \lambda^d (2\pi)^{\frac{d}{2}} u(y) \int_{\mathbb{R}^d} g(\lambda(y-x))^2 dx \\
&= u(y),
\end{aligned}$$

where we used the evaluations

$$\Gamma^B(y, x, y') \Big|_{y'=y} = \varphi^A(y, x) + \varphi^A(x, y) = 0, \quad e^{-i\varphi^A(y',y)} \Big|_{y'=y} = 0.$$

From the above calculation we can also see the nice interaction between two wavepackets.

We then define the magnetic modulation space $M_A^{p,q}(\mathbb{R}^d)$ adapted to the magnetic potential A to be the space of functions whose \mathcal{T}^A transform has $L_x^p L_\xi^q$ integrability.

Definition 4.4. For $1 \leq p, q < \infty$, we define the magnetic modulation space $M_A^{p,q}(\mathbb{R}^d)$ by

$$M_A^{p,q}(\mathbb{R}^d) := \left\{ u \in \mathcal{S}'(\mathbb{R}^d) : \|u\|_{M_A^{p,q}(\mathbb{R}^d)} := \left(\int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} |\mathcal{T}_\lambda^A u(x, \xi)|^p dx \right)^{q/p} d\xi \right)^{1/q} < \infty \right\}.$$

For $p = \infty$ or $q = \infty$, the integral should be replaced by L^∞ norm and the space $M_A^{p,q}(\mathbb{R}^d)$ is defined as the closure of the Schwartz space with respect to the norm.

By this definition, when $A \equiv 0$ we recover the usual modulation space $M^{p,q}(\mathbb{R}^d)$. General information and properties of the non-magnetic modulation space $M^{p,q}(\mathbb{R}^d)$ can be found in [7] or Chapter 2 in [2]. As far as this article is concerned, we focus on the case $p = q$, and denote $M_A^p(\mathbb{R}^d) := M_A^{p,p}(\mathbb{R}^d)$. We can prove analogously the following properties.

Proposition 4.5. *Let $1 \leq p \leq \infty$. We have the following.*

- (i) $M_A^p(\mathbb{R}^d)$ is independent of the choice $\lambda \geq 1$.
- (ii) $M_A^p(\mathbb{R}^d)$ is a Banach space.

(iii) $\mathcal{S}(\mathbb{R}^d)$ is dense in $M_A^p(\mathbb{R}^d)$.

(iv) For $u \in M_A^p(\mathbb{R}^d)$, we have $u = \mathcal{T}_\lambda^{A*} \mathcal{T}_\lambda^A u$.

In particular, for $p = 2$ we have $M_A^2(\mathbb{R}^d) \simeq L^2(\mathbb{R}^d)$ isometrically.

Up to now, we have used the magnetic vector potential A from the transversal gauge (7) to define the wavepacket transform and the modulation spaces. Here, we remark on the general magnetic potentials $A'(y) := A(y) + \partial_y v(y)$. By replacing A with A' , we can define accordingly $\mathcal{T}^{A'}$ and $M_{A'}^p(\mathbb{R}^d)$, which actually leads to the same space.

Proposition 4.6. For $v \in C_{\text{pol}}^\infty(\mathbb{R}^d, \mathbb{R})$ and $A'(y) := A(y) + \partial_y v(y)$, which satisfies $dA' = dA$, we have $M_{A'}^p(\mathbb{R}^d) \simeq M_A^p(\mathbb{R}^d)$ isometrically.

Proof. Notice that $A'(y) := A(y) + \partial_y v(y)$ gives

$$\varphi^{A'}(y, x) = \int_{\Gamma_{x,y}} A' = \int_{\Gamma_{x,y}} (A + \partial v) = \varphi^A(y, x) + v(y) - v(x).$$

Consider the injective map $u \mapsto e^{iv} u$ for $u \in M_A^p(\mathbb{R}^d)$. Then for $w := e^{iv} u$, we have

$$\mathcal{T}^{A'} w(x, \xi) = \int e^{-i\xi \cdot (y-x)} g(y-x) e^{-i\varphi^{A'}(y,x)} w(y) dy = e^{iv(x)} \mathcal{T}^A u(x, \xi),$$

so $w \in M_{A'}^p(\mathbb{R}^d)$ with $\|w\|_{M_{A'}^p(\mathbb{R}^d)} = \|u\|_{M_A^p(\mathbb{R}^d)}$. The map $u \mapsto e^{iv} u$ is also surjective, since for any $w' \in M_{A'}^p(\mathbb{R}^d)$, we can take $u' := e^{-iv} w'$, and by a similar calculation as above we have $u' \in M_A^p(\mathbb{R}^d)$. \square

We also introduce the weighted version $M_A^{m,p}(\mathbb{R}^d)$.

Definition 4.7. Let $1 \leq p \leq \infty$ and $m \in \mathbb{R}$. We denote the standard weights $\nu_m(x, \xi) := (1 + |x|^2 + |\xi|^2)^{m/2}$ and the re-scaled weights $\nu_{\lambda,m}(x, \xi) := (1 + |\lambda^{-1}x|^2 + |\lambda^{-1}\xi|^2)^{m/2}$. The weighted magnetic modulation space $M_A^{m,p}(\mathbb{R}^d)$ is defined by

$$M_A^{m,p}(\mathbb{R}^d) := \left\{ u \in \mathcal{S}'(\mathbb{R}^d) : \|u\|_{M_A^{m,p}(\mathbb{R}^d)} := \|\nu_{\lambda,m} \mathcal{T}_\lambda^A u\|_{L^p(\mathbb{R}^{2d})} < \infty \right\},$$

where the case $p = \infty$ is also understood as the closure of the Schwartz space with respect to the associated norm.

By the above definition, $M_A^{m,p}(\mathbb{R}^d)$ is essentially a weighted L^p space from a phase space perspective, so it also enjoys properties analogous to the ones in Proposition 4.5. For $m_1 \geq m_0$, we have naturally $M_A^{m_1,p}(\mathbb{R}^d) \subseteq M_A^{m_0,p}(\mathbb{R}^d)$. For $p = 2$, by the isomorphism

$M_A^{0,2}(\mathbb{R}^d) \simeq L^2(\mathbb{R}^d)$, it is easy to see for $k \in \mathbb{N}$, the space $M_A^{k,2}(\mathbb{R}^d)$ contains the L^2 -based weighted Sobolev space $\Sigma(k)$ used in Yajima [16]

$$\Sigma(k) = \{u \in L^2(\mathbb{R}^d) : \sum_{|\alpha|+|\beta| \leq k} \|x^\alpha \partial_x^\beta u\|_{L^2(\mathbb{R}^d)}^2 = \|u\|_{\Sigma(k)}^2 < \infty\}.$$

Moreover, for our generalized magnetic Schrödinger operator $\text{Op}^A(h)$ with symbol h satisfying Assumption 1.3, the space $M_A^{2,p}$ is a natural domain for $\text{Op}^A(h)$ in M_A^p .

Lemma 4.8. *Let $1 \leq p \leq \infty, m \in \mathbb{R}$, and let symbol h satisfies the Assumption 1.3. Then the magnetic pseudo-differential operator $\text{Op}^A(h) : M_A^{m,p}(\mathbb{R}^d) \rightarrow M_A^{m-2,p}(\mathbb{R}^d)$ is bounded.*

This lemma can be seen as a consequence of the following more general lemma.

Lemma 4.9. *Let $1 \leq p \leq \infty$ and $m, s \in \mathbb{R}$. If a linear operator $\mathcal{L} : \mathcal{S}(\mathbb{R}^d) \rightarrow \mathcal{S}'(\mathbb{R}^d)$ satisfies for all $N, N' \in \mathbb{N}$*

$$(20) \quad |\langle g_{z,\zeta}^A, \mathcal{L}g_{x,\xi}^A \rangle| \lesssim \nu_s(z, \zeta) \langle z - x \rangle^{-N} \langle \zeta - \xi \rangle^{-N'},$$

then $\mathcal{L} : M_A^{m,p}(\mathbb{R}^d) \rightarrow M_A^{m-s,p}(\mathbb{R}^d)$ is bounded.

Proof. By density of the Schwartz space $\mathcal{S}(\mathbb{R}^d)$ in $M_A^{m,p}(\mathbb{R}^d)$, it suffices to consider $u \in \mathcal{S}(\mathbb{R}^d)$. Denote $G_{\mathcal{L}}(z, \zeta, x, \xi) := \langle g_{z,\zeta}^A, \mathcal{L}g_{x,\xi}^A \rangle$. By the reproducing formula $u = \mathcal{T}^{A*} \mathcal{T}^A u$ we can write

$$\begin{aligned} & \|\mathcal{L}u\|_{M_A^{m-s,p}(\mathbb{R}^d)}^p \\ &= \int_{\mathbb{R}^{2d}} \left| \nu_{m-s}(z, \zeta) \int_{\mathbb{R}^d} \overline{g_{z,\zeta}^A}(y) \mathcal{L} \left(\int_{\mathbb{R}^{2d}} g_{x,\xi}^A(y) \mathcal{T}^A u(x, \xi) dx d\xi \right) dy \right|^p dz d\zeta \\ &= \int_{\mathbb{R}^{2d}} \left| \int_{\mathbb{R}^{2d}} \nu_{m-s}(z, \zeta) G_{\mathcal{L}}(z, \zeta, x, \xi) \nu_{-m}(x, \xi) \nu_m(x, \xi) \mathcal{T}^A u(x, \xi) dx d\xi \right|^p dz d\zeta. \end{aligned}$$

Notice that the inner integral on RHS above can be seen as integrating a phase space function $\nu_m \mathcal{T}^A u$ against an integral kernel involving $G_{\mathcal{L}}$. By the property (20), the integral kernel can be bounded for all $N, N' \in \mathbb{N}$ by

$$\begin{aligned} & |\nu_{m-s}(z, \zeta) G_{\mathcal{L}}(z, \zeta, x, \xi) \nu_{-m}(x, \xi)| \\ & \lesssim \nu_m(z, \zeta) \langle z - x \rangle^{-N} \langle \zeta - \xi \rangle^{-N'} \nu_{-m}(x, \xi) \\ & \lesssim \nu_m(z - x, \zeta - \xi) \nu_m(x, \xi) \langle z - x \rangle^{-N} \langle \zeta - \xi \rangle^{-N'} \nu_{-m}(x, \xi) \\ & \lesssim \langle z - x \rangle^{-N+m} \langle \zeta - \xi \rangle^{-N'+m}. \end{aligned}$$

With the sufficient off-diagonal decay of the kernel above, we obtain the boundedness of \mathcal{L} by Young's inequality for integral operators

$$\|\mathcal{L}u\|_{M_A^{m-s,p}(\mathbb{R}^d)} \lesssim \|\nu_m \mathcal{T}^A u\|_{L^p(\mathbb{R}^{2d})} = \|u\|_{M_A^{m,p}(\mathbb{R}^d)}.$$

□

5. MAGNETIC PHASE SPACE FLOW

The magnetic Weyl quantization $h \mapsto \text{Op}^A(h)$ gives a correspondence between the classical Hamiltonian h and the quantum observable $\text{Op}^A(h)$. In the spirit of the correspondence principle, it is natural to relate the quantum evolution $u_0 \mapsto u(t)$ on \mathbb{R}^d to a classical phase flow on the phase space $T^*\mathbb{R}^d \simeq \mathbb{R}^{2d}$. Since our choice of quantization depends intrinsically on the magnetic field B , the corresponding flow should also depend on B instead of the vector potential A . This is achieved by equipping the phase space $T^*\mathbb{R}^d$ with the twisted symplectic form ω_B , given by

$$\omega_{B(x)} := \sum_{j=1}^d dx_j \wedge d\xi_j + \frac{1}{2} \sum_{j,k} B_{jk}(x) dx_j \wedge dx_k,$$

which then gives rise to the associated Poisson bracket $\{\cdot, \cdot\}_B$

$$\{f, g\}_B := \sum_{j=1}^d (\partial_{\xi_j} f \partial_{x_j} g - \partial_{x_j} f \partial_{\xi_j} g) + \sum_{j,k=1}^d B_{jk}(\cdot) \partial_{\xi_j} f \partial_{\xi_k} g.$$

With the magnetic Poisson bracket above, we define the magnetic Hamiltonian flow induced by a time-dependent Hamiltonian h .

Definition 5.1. For a given magnetic field B , we define $\chi_B(h)(t, s) : (x^s, \xi^s) \mapsto (x^t, \xi^t)$ the magnetic Hamiltonian flow induced by the Hamiltonian $h \in C(\mathbb{R}, C^\infty(\mathbb{R}^{2d}))$, satisfying the equations

$$(21) \quad \begin{cases} \dot{x}_j^t = \{h(t), x_j^t\}_B, \\ \dot{\xi}_j^t = \{h(t), \xi_j^t\}_B. \end{cases}$$

By definition of the Poisson bracket, the flow (x^t, ξ^t) satisfies more explicitly

$$(22) \quad \dot{x}_j^t = \partial_{\xi_j} h(t; x^t, \xi^t), \quad \dot{\xi}_j^t = -\partial_{x_j} h(t; x^t, \xi^t) + \sum_{k=1}^d B_{kj}(x^t) \partial_{\xi_k} h(t; x^t, \xi^t).$$

In the following, we focus on such magnetic flows associated to symbols h satisfying $h(t) \in S^{0,(2)}(\mathbb{R}^{2d})$.

Lemma 5.2. *Assume $h \in C(\mathbb{R}, C^\infty(\mathbb{R}^{2d}))$ with $h(t) \in S^{0,(2)}(\mathbb{R}^{2d})$ uniformly in t , and let the magnetic field B satisfy $\|B_{jk}\|_\infty \leq C_B$. Then the flow $\chi_B(h)$ exists globally in time and $\chi_B(h)(t, s)$ is volume-preserving for all $t, s \in \mathbb{R}$.*

Proof. For this lemma, the t -dependence of $h(t)$ does not play a role, because we only need the bounds for the x, ξ derivatives of h , which by our assumption are uniform in t . We thus suppress the notation and just take $h(x, \xi) := h(t; x, \xi)$. For the existence, we firstly note that the vector field in (22) associated to $h \in S^{0,(2)}(\mathbb{R}^{2d})$

$$\Phi_h := \left(\partial_{\xi_1} h, \dots, \partial_{\xi_d} h, -\partial_{x_1} h + \sum_k B_{k1}(x) \partial_{\xi_k} h, \dots, -\partial_{x_d} h + \sum_k B_{kd}(x) \partial_{\xi_k} h \right)$$

is smooth and thus locally Lipschitz, so $\chi_B(h)(t, s)$ exists locally in time. To extend it globally, we rule out finite time blow-up by showing, for $s, t \in \mathbb{R}$ with $s < t$, we can take $T > 0$ such that $t - s < T$ and bound

$$(23) \quad 1 + |x^t| + |\xi^t| \leq C_T(1 + |x^s| + |\xi^s|).$$

By the boundedness of the second-order derivatives of $h \in S^{0,(2)}(\mathbb{R}^{2d})$, we can bound its first-order derivatives by

$$|\partial_x h(x, \xi)| + |\partial_\xi h(x, \xi)| \leq C_h(1 + |x| + |\xi|).$$

Take $C_B \geq 1$ such that $|B(x)| \leq C_B$, and denote $C_{h,B} := C_h C_B \geq C_h$. The above bounds for the derivatives of h and the expression (22) for \dot{x}^t and $\dot{\xi}^t$ imply

$$(24) \quad |\dot{x}^t| \leq C_h(1 + |x^t| + |\xi^t|), \quad |\dot{\xi}^t| \leq C_{h,B}(1 + |x^t| + |\xi^t|).$$

We then calculate

$$(25) \quad \begin{aligned} \frac{d|x^t|}{dt} &= \frac{\dot{x}^t \cdot x^t}{|x^t|} \leq |\dot{x}^t| \leq C_h(1 + |x^t| + |\xi^t|), \\ \frac{d|\xi^t|}{dt} &= \frac{\dot{\xi}^t \cdot \xi^t}{|\xi^t|} \leq |\dot{\xi}^t| \leq C_{h,B}(1 + |x^t| + |\xi^t|). \end{aligned}$$

Combining the above bounds, we see that

$$\frac{d(1 + |x^t| + |\xi^t|)}{dt} \leq 2C_{h,B}(1 + |x^t| + |\xi^t|).$$

By Gronwall's inequality, we obtain the desired upper bound (23)

$$(26) \quad 1 + |x^t| + |\xi^t| \leq e^{2TC_{h,B}}(1 + |x^s| + |\xi^s|).$$

Moreover, since the backward-in-time flow $(x^t, \xi^t) \mapsto (x^s, \xi^s)$ can be seen as a flow driven by the vector field $-\Phi_h$, a similar argument as above also gives a reversed bound

$$(27) \quad 1 + |x^s| + |\xi^s| \leq e^{2TC_{h,B}}(1 + |x^t| + |\xi^t|).$$

For the volume-preserving property, we use the fact that a flow generated by a divergence-free vector field is volume-preserving, see Proposition 16.33 in [11]. For the vector field Φ_h we can compute

$$\operatorname{div} \Phi_h = \sum_{j,k=1}^d \partial_{x_j} \partial_{\xi_k} h - \partial_{\xi_j} \partial_{x_k} h + B_{kj}(x) \partial_{\xi_j} \partial_{\xi_k} h = 0,$$

so the flow $\chi_B(h)$ is volume-preserving. \square

The following lemma shows that the magnitude of the position and momentum vectors of certain magnetic flows are comparable in a time-averaged sense. The lemma is inspired by [16, Lemma 2.1], and is one of the crucial technical ingredients for dealing with unbounded magnetic potentials.

Lemma 5.3. *Let $\chi_B(h)$ be the magnetic Hamiltonian flow associated to a symbol h satisfying Assumption 1.3 and a magnetic field B satisfying $\|B_{jk}\|_\infty \leq C_B$. Let $I \subset \mathbb{R}$ be a compact interval and fix an arbitrary $\epsilon > 0$. We have for $(x^t, \xi^t), t \in I$, satisfying the flow $\chi_B(h)$ in (21),*

$$(28) \quad \int_I \langle x^\tau \rangle^{-1-\epsilon} |\xi^\tau| d\tau \leq C_{h,B,\epsilon}(1 + |I|), \quad (x, \xi) \in \mathbb{R}^{2d}.$$

Proof. We prove this in a similar way as [16, Lemma 2.1]. Again the t -dependence of $h(t)$ only makes a difference at the very end of the proof, so we suppress the notation for now. By Assumption 1.3, $h \in S^{0,(2)}(\mathbb{R}^{2d})$ and is second-order in ξ . We thus have a lower bound for its ξ -derivative, that is, there exist $b \in \mathbb{R}^d, C \in \mathbb{R}$ such that

$$(29) \quad |\partial_\xi h(x, \xi)| \geq \|\xi\| - |b \cdot x + C|.$$

We denote a new constant $C_{h,B}$ based on C_h, C_B and $|b|$ by

$$C_{h,B} := \max(|b|, C_h C_B).$$

Without loss of generality, we can assume the constant $C_{h,B} \geq 1$, since otherwise the arguments below hold by replacing $C_{h,B}$ everywhere with 1. Sub-divide the interval $I = \sum_j I_j$

into compact intervals with length $|I_j| = T$, such that T is sufficiently small satisfying

$$(30) \quad 200T^2C_{h,B}^3 + 60TC_{h,B}^2 \leq \frac{1}{4}.$$

It suffices to prove (28) separately for each I_j . We fix I_j and suppress the index j , denoting it again by I . We also fix the starting point of the flow as $x, \xi \in \mathbb{R}^d$. Firstly, for the case of $TC_{h,B}|\xi^t| \leq 1 + |x^t|$ for all $t \in I$, the integral is bounded by

$$(31) \quad \int_I \langle x^\tau \rangle^{-1-\epsilon} |\xi^\tau| d\tau \leq (TC_{h,B})^{-1} \int_I \langle x^\tau \rangle^{-1-\epsilon} (1 + |x^\tau|) d\tau \leq C_{h,B}^{-1}.$$

For the other case, there exists $r \in I$ such that $TC_{h,B}|\xi^r| > 1 + |x^r|$. We claim the following bounds hold for all $t \in I$:

$$(32) \quad 1 + |x^t| < 10TC_{h,B}|\xi^r|, \quad \frac{2}{3}|\xi^r| \leq |\xi^t| \leq \frac{3}{2}|\xi^r|, \quad t \in I.$$

We can in fact prove a stronger claim, that the above bounds are satisfied for all t with $|t - r| \leq T$. This claim can be verified by contradiction. Denote $T^* := \sup\{\sigma \leq T : (32) \text{ is satisfied for } |t - r| \leq \sigma\}$ and suppose $T^* < T$. For t with $|t - r| \leq T^*$, we can bound $1 + |x^t|$ by

$$1 + |x^t| \leq 1 + |x^r| + T^* \sup_{\tau \in I} \left| \frac{d|x^\tau|}{d\tau} \right|.$$

Then by $TC_{h,B}|\xi^r| > 1 + |x^r|$, the bound for $\frac{d|x^t|}{dt}$ in (25) and the bound (26), we obtain

$$\begin{aligned} 1 + |x^t| &\leq 1 + |x^r| + T^*C_{h,B}e^{2TC_{h,B}}(1 + |x^r| + |\xi^r|) \\ &< TC_{h,B}|\xi^r| + 3TC_{h,B}(TC_{h,B}|\xi^r| + |\xi^r|) \\ &= 4TC_{h,B}|\xi^r| + 3T^2C_{h,B}^2|\xi^r| \\ &< 9TC_{h,B}|\xi^r|, \end{aligned}$$

where we used $e^{2TC_{h,B}} < 3$ because by the smallness assumption (30) of T we have $2TC_{h,B} < 1$. Similarly by the bound for $\frac{d|\xi^t|}{dt}$ in (25) and the above, we have

$$\begin{aligned} |\xi^t| &\leq |\xi^r| + T^*C_{h,B}e^{2TC_{h,B}}(1 + |x^r| + |\xi^r|) \\ &< |\xi^r| + (3T^2C_{h,B}^2 + 3TC_{h,B})|\xi^r| \\ &< \frac{11}{10}|\xi^r|, \end{aligned}$$

where the smallness of T as in (30) is also used to obtain the last inequality. The lower bound of $|\xi^t|$ holds analogously, that is

$$|\xi^t| > |\xi^r| - (3T^2C_{h,B}^2 + 3TC_{h,B})|\xi^r| \geq \frac{9}{10}|\xi^r|.$$

Because the flow (x^t, ξ^t) is continuous in t , such T^* cannot be maximal, thus contradicting its definition.

Now for the constant C in (29), we distinguish two cases based on the size of $|\xi^r|$. Firstly for the case that $|\xi^r| \leq 10|C|$, we directly have a desired bound

$$\int_I \langle x^\tau \rangle^{-1-\epsilon} |\xi^\tau| d\tau \leq \int_I |\xi^r| d\tau \leq 10|C|T.$$

For the other case $|\xi^r| > 10|C|$, using (29), (25) and (32) we can calculate a lower bound for the second-order t -derivative of $|x^t|$. Here we will be taking t -derivatives on $h(t)$, so the t -dependence makes a difference, but things are under control since by our Assumption 1.3, we have $\dot{h}(t) \in S^{0,(2)}(\mathbb{R}^{2d})$ uniformly in t . In particular, the assumption implies

$$|\partial_\xi \dot{h}(t; x^t, \xi^t)| \leq C_h(1 + |x^t| + |\xi^t|).$$

Using bounds (24) and (32), we have

$$\begin{aligned} \frac{d^2|x^t|^2}{dt^2} &= 2|\partial_\xi h(t; x^t, \xi^t)|^2 + 2\frac{d\partial_\xi h(t; x^t, \xi^t)}{dt} \cdot x^t \\ &= 2|\partial_\xi h(t; x^t, \xi^t)|^2 + 2\left(\partial_\xi \dot{h}(t; x^t, \xi^t) + \partial_x \partial_\xi h(t; x^t, \xi^t) \dot{x}^t + \partial_\xi^2 h(t; x^t, \xi^t) \xi^t\right) \cdot x^t \\ &\geq 2\left(\frac{9}{10}|\xi^t| - |b \cdot x^t|\right)^2 - 2C_{h,B}(1 + |x^t| + |\xi^t|)|x^t| \\ &\geq 2\left(\frac{3}{5}|\xi^r| - 10TC_{h,B}^2|\xi^r|\right)^2 - 2C_{h,B}(10TC_{h,B}|\xi^r| + \frac{3}{2}|\xi^r|)10TC_{h,B}|\xi^r| \\ &\geq \left(\frac{1}{2} - (200T^2C_{h,B}^3 + 60TC_{h,B}^2)\right)|\xi^r|^2, \end{aligned}$$

so by the smallness of T as in (30), we have $\frac{d^2|x^t|^2}{dt^2} \geq \frac{1}{4}|\xi^r|^2$. Then we show for $\tau \in I$, $|x^\tau|$ can be bounded from below. Suppose at a time $\rho \in I$, x^ρ attains the minimum value $|x^\rho| = \inf_{t \in I} |x^t|$. Then by a Taylor expansion of $|x^\tau|^2$ in the t -variable at $|x^\rho|^2$, we have

$$\begin{aligned} |x^\tau|^2 &= |x^\rho|^2 + \frac{d|x^t|^2}{dt} \Big|_{t=\rho} (\tau - \rho) + \frac{1}{2} \frac{d^2|x^t|^2}{dt^2} \Big|_{t=\rho} (\tau - \rho)^2 \\ &\geq |x^\rho|^2 + \frac{1}{8}|\xi^r|^2(\tau - \rho)^2, \end{aligned}$$

where in the last inequality we used the fact that $|x^\rho|^2$ is a minimum, so $\frac{d|x^t|^2}{dt} \Big|_{t=\rho} (\tau - \rho) \geq 0$.

Then we have the desired bound for the integral

$$\begin{aligned} \int_I \langle x^\tau \rangle^{-1-\epsilon} |\xi^\tau| d\tau &\lesssim \int_I (1 + |x^\tau|)^{-1-\epsilon} |\xi^\tau| d\tau \\ &\lesssim \int_I (1 + |\xi^r| |\tau - \rho|)^{-1-\epsilon} |\xi^r| d\tau \\ &\leq C' \epsilon^{-1}, \end{aligned}$$

where we used the lower bound of $|x^\tau|$ and the fact that $|\xi^\tau|, |\xi^\rho|$ are comparable to $|\xi^r|$ by (32). Combining this with the bound (31) for the other case, we obtain (28). \square

6. PHASE SPACE APPROXIMATION

The evolution $u_0 \mapsto u(t)$ satisfying (9) can be seen as a quantum evolution associated to the operator $\text{Op}^A(h)$. Our goal in this section is to approximate such an evolution with a classical evolution associated to the Hamiltonian h in the phase space. This is in some sense a manifestation of the quantum-classical correspondence principle, and the relevant classical evolution turns out to follow a magnetic Hamiltonian flow $\chi_B(h)$ in phase space. So by taking a phase space perspective using the wavepacket transform \mathcal{T}_λ^A , we can obtain an approximate evolution in phase space that is more transparent and tractable. This change of perspective reveals that to some extent, the quantum evolution $u_0 \mapsto u(t)$ preserves phase space localizations in the sense of magnetic wavepackets, which leads to the magnetic modulation space boundedness of $u_0 \mapsto u(t)$.

6.1. Conjugating magnetic Ψ DOs. The first step in approximating the generalized magnetic Schrödinger evolution in phase space is to approximate the effect of the operator $\text{Op}^A(h)$ in phase space. That is, we need to understand how the conjugated operator $\mathcal{T}_\lambda^A \text{Op}^A(h) \mathcal{T}_\lambda^{A*}$ behaves on phase space functions. For real symbols h , the operator $\text{Op}^A(h)$ is symmetric for the pairing of $u \in \mathcal{S}(\mathbb{R}^d)$ and $g_{x,\xi}^{A,\lambda}$,

$$\mathcal{T}_\lambda^A \text{Op}^A(h)u(x, \xi) = \left\langle g_{x,\xi}^{A,\lambda}, \text{Op}^A(h)u \right\rangle_{L^2} = \left\langle \text{Op}^A(h)g_{x,\xi}^{A,\lambda}, u \right\rangle_{L^2},$$

so it boils down to determining an approximation of the term $\text{Op}^A(h)g_{x,\xi}^{A,\lambda}$ in phase space. The following proposition relates differentiation and multiplication on the configuration space and the phase space.

Lemma 6.1. *For magnetic wavepackets $g_{x,\xi}^{A,\lambda}(y)$, multiplication and differentiation with respect to $y \in \mathbb{R}^d$ can be approximated by multiplication and differentiation with respect to $(x, \xi) \in \mathbb{R}^{2d}$, that is,*

$$(33) \quad \begin{aligned} -i\partial_{y_j} g_{x,\xi}^{A,\lambda}(y) &= i\partial_{x_j} g_{x,\xi}^{A,\lambda}(y) - \sum_{k=1}^d B_{jk}(x) i\partial_{\xi_k} g_{x,\xi}^{A,\lambda}(y) \\ &\quad + [A_j(y) - A_j(x) - r_x^{(1)}(A_j(y, x)) + r_x^{(1)}(A_j(x, y))] g_{x,\xi}^{A,\lambda}(y), \end{aligned}$$

where $r_x^{(1)}$ denotes the remainder of the first-order Taylor expansion of y at x .

Proof. By property (12) and (14) of $\varphi^A(y, x)$, we have

$$\begin{aligned} \partial_{y_j} e^{i\varphi^A(y,x)} &= i(A_j(y, 0) - A_j(y, x)) e^{i\varphi^A(y,x)}, \\ \partial_{x_j} e^{i\varphi^A(y,x)} &= i(-A_j(x, 0) + A_j(x, y)) e^{i\varphi^A(y,x)}. \end{aligned}$$

A direct computation gives

$$-i\partial_{y_j}g_{x,\xi}^{A,\lambda}(y) = i\partial_{x_j}g_{x,\xi}^{A,\lambda}(y) + [A_j(y) - A_j(x) - A_j(y, x) + A_j(x, y)]g_{x,\xi}^{A,\lambda}(y).$$

Next, we use Taylor's formula to expand the term $-A_j(y, x) + A_j(x, y)$ with respect to the y variable at the value x . Recall the definition of the potential

$$A_j(y, x) := -\sum_{k=1}^d \int_0^1 s(y_k - x_k) B_{jk}(x + s(y - x)) ds,$$

so evaluating A_j at $y = x$ we have $A_j(x, x) = 0$. The first-order derivatives of A_j are

$$\partial_{y_l}A_j(y, x) = -\int_0^1 sB_{jl}(x + s(y - x)) ds - \sum_{k=1}^d \int_0^1 s^2(y_k - x_k) \partial_l B_{jk}(x + s(y - x)) ds.$$

The derivatives at $y = x$ evaluate to

$$(34) \quad \partial_{y_l}A_j(x, x) = -\int_0^1 sB_{jl}(x) ds = -\frac{1}{2}B_{jl}(x).$$

Further, the second-order derivatives are, for $1 \leq m, l \leq d$,

$$(35) \quad \begin{aligned} \partial_{y_m}\partial_{y_l}A_j(y, x) &= -\int_0^1 s^2\partial_m B_{jl}(x + s(y - x)) ds - \int_0^1 s^2\partial_l B_{jm}(x + s(y - x)) ds \\ &\quad - \sum_{k=1}^d \int_0^1 s^3(y_k - x_k) \partial_m\partial_l B_{jk}(x + s(y - x)) ds, \end{aligned}$$

so by (34) and (35), Taylor's formula of A_j takes the form

$$A_j(y, x) = -\sum_{l=1}^d \frac{1}{2}B_{jl}(x)(y_l - x_l) + r_x^{(1)}(A_j(y, x)),$$

where, for some $\theta \in (0, 1)$, the remainder can be expressed by

$$(36) \quad r_x^{(1)}(A_j(y, x)) := \sum_{m,l=1}^d (y_m - x_m)(y_l - x_l) \partial_{y_m}\partial_{y_l}A_j(x + \theta(y - x), x).$$

Expanding also $A_j(x, y)$ and combining the terms, we obtain

$$-A_j(y, x) + A_j(x, y) = \sum_{l=1}^d B_{jl}(x)(y_l - x_l) - r_x^{(1)}(A_j(y, x)) + r_x^{(1)}(A_j(x, y)).$$

The lemma then follows from the identity $\partial_{\xi_j}g_{x,\xi}^{A,\lambda}(y) = i(y_j - x_j)g_{x,\xi}^{A,\lambda}(y)$. \square

We build upon the above lemma to show that the action of $\text{Op}^A(h)$ on wavepackets $g_{x,\xi}^{A,\lambda}$ can be approximated, up to some error terms, as a phase space differentiation along the

magnetic Hamiltonian vector field associated to h . The lemma below can hold for more general h , but for simplicity we formulate it for the most relevant symbol class $S^{0,(2)}(\mathbb{R}^{2d})$.

Lemma 6.2. *Let $h \in S^{0,(2)}(\mathbb{R}^{2d})$. The magnetic pseudo-differentiation $\text{Op}^A(h)$ on wavepackets $g_{x,\xi}^{A,\lambda}$ can be approximated by a phase space differentiation \tilde{H}^B . That is,*

$$(37) \quad \text{Op}^A(h)g_{x,\xi}^{A,\lambda}(y) = \left(i\tilde{H}^B + \mathcal{R} \right) g_{x,\xi}^{A,\lambda}(y), \quad y \in \mathbb{R}^d,$$

where the components of \tilde{H}^B are

$$\tilde{H}_j^B := \left(\partial_{\xi_j} h \partial_{x_j}, -\partial_{x_j} h \partial_{\xi_j} + \sum_{k=1}^d B_{kj}(x) \partial_{\xi_k} h \partial_{\xi_j} \right), \quad 1 \leq j \leq d.$$

The approximation error \mathcal{R} term consists of the three parts

$$\mathcal{R} = m^A(h) + \mathcal{R}_1 + \mathcal{R}_2,$$

as defined in (43), (44), and (45), where $m^A(h)$ denotes a multiplier that only depends on the phase space variables x, ξ , while $\mathcal{R}_1, \mathcal{R}_2$ denotes respectively a multiplicative operator and a magnetic pseudo-differential operator on the y variable.

Proof. For convenience of later proofs, we first convert to the magnetic Kohn-Nirenberg quantization $\text{Op}_{KN}^A(h^*) = \text{Op}^A(h)$. By Lemma 3.2, the symbol $h^* \in S^{0,(2)}(\mathbb{R}^{2d})$ is given by

$$(38) \quad h^*(y, \eta) := h(y, \eta) + h_r(y, \eta), \quad h_r \in S^{0,(0)}(\mathbb{R}^{2d}),$$

and we can represent $\text{Op}^A(h)$ using the Kohn-Nirenberg quantization

$$\text{Op}^A(h) g_{x,\xi}^{A,\lambda}(y) = \left(\text{Op}_{KN}^A(h) + \text{Op}_{KN}^A(h_r) \right) g_{x,\xi}^{A,\lambda}(y).$$

The $\text{Op}_{KN}^A(h_r)$ will be included in the pseudo-differential remainder \mathcal{R}_2 , since its symbol h_r is nicely bounded. Our focus here is to approximate $\text{Op}_{KN}^A(h)$, and we start with a Taylor expansion on h at the point (x, ξ) , that is,

$$(39) \quad \begin{aligned} & \text{Op}_{KN}^A(h)g_{x,\xi}^{A,\lambda}(y) \\ &= \text{Op}_{KN}^A \left[h(x, \xi) + \partial_y h(x, \xi) \cdot (y - x) + \partial_\eta h(x, \xi) \cdot (\eta - \xi) + r_{x,\xi}^{(1)}(h)(y, \eta) \right] g_{x,\xi}^{A,\lambda}(y) \\ &= \text{Op}_{KN}^A \left[\partial_y h(x, \xi) \cdot (y - x) + \partial_\eta h(x, \xi) \cdot \eta + r_{x,\xi}^{(1)}(h)(y, \eta) \right] g_{x,\xi}^{A,\lambda}(y) \\ & \quad + (h(x, \xi) - \partial_\eta h(x, \xi) \cdot \xi) g_{x,\xi}^{A,\lambda}(y). \end{aligned}$$

Here $r_{x,\xi}^{(1)}(h)$ denotes the remainder term from the first-order expansion:

$$(40) \quad r_{x,\xi}^{(1)}(h)(y, \eta) := \int_0^1 (1-s) \partial_s^2 h(x + s(y-x), \xi + s(\eta - \xi)) ds.$$

By the basic quantizations (18) and the identity (33) in Lemma 6.1, we can calculate two magnetic Ψ DO terms on the right hand side of (39) as

$$(41) \quad \begin{aligned} \text{Op}_{KN}^A [\partial_y h(x, \xi) \cdot (y-x)] g_{x,\xi}^{A,\lambda}(y) &= \partial_y h(x, \xi) \cdot (y-x) g_{x,\xi}^{A,\lambda}(y) \\ &= -i \partial_y h(x, \xi) \cdot \partial_\xi g_{x,\xi}^{A,\lambda}(y), \end{aligned}$$

and

$$(42) \quad \begin{aligned} &\text{Op}_{KN}^A [\partial_\eta h(x, \xi) \cdot \eta] g_{x,\xi}^{A,\lambda}(y) \\ &= \partial_\eta h(x, \xi) \cdot (-i \partial_y - A(y)) g_{x,\xi}^{A,\lambda}(y) \\ &= \sum_{j=1}^d \partial_{\eta_j} h(x, \xi) \left(i \partial_{x_j} g_{x,\xi}^{A,\lambda}(y) - \sum_{k=1}^d B_{jk}(x) i \partial_{\xi_k} g_{x,\xi}^{A,\lambda}(y) \right) + \\ &\quad + \sum_{j=1}^d \partial_{\eta_j} h(x, \xi) \left(-A_j(x) - r_x^{(1)}(A_j(y, x)) + r_x^{(1)}(A_j(x, y)) \right) g_{x,\xi}^{A,\lambda}(y). \end{aligned}$$

Then substituting (41), (42) into (39) yields

$$\begin{aligned} &\text{Op}^A(h) g_{x,\xi}^{A,\lambda}(y) \\ &= i \sum_{j=1}^d \partial_{\eta_j} h(x, \xi) \partial_{x_j} g_{x,\xi}^{A,\lambda}(y) - i \sum_{j=1}^d \left(\partial_{y_j} h(x, \xi) - \sum_{k=1}^d B_{kj}(x) \partial_{\eta_k} h(x, \xi) \right) \partial_{\xi_j} g_{x,\xi}^{A,\lambda}(y) + \\ &\quad + (m^A(h) + \mathcal{R}_1 + \mathcal{R}_2) g_{x,\xi}^{A,\lambda}(y) \end{aligned}$$

with a multiplicative factor $m^A(h)$ which only depends on the variable x, ξ

$$(43) \quad m^A(h)(x, \xi) := h(x, \xi) - \partial_\eta h(x, \xi) \cdot (\xi + A(x)),$$

and two other error terms involving the y variable, including one multiplicative \mathcal{R}_1 ,

$$(44) \quad \mathcal{R}_1 g_{x,\xi}^{A,\lambda}(y) := \partial_\eta h(x, \xi) \cdot \left(-r_x^{(1)}(A(y, x)) + r_x^{(1)}(A(x, y)) \right) g_{x,\xi}^{A,\lambda}(y)$$

and one magnetic pseudo-differential \mathcal{R}_2 ,

$$(45) \quad \mathcal{R}_2 g_{x,\xi}^{A,\lambda}(y) := \text{Op}_{KN}^A \left(r_{x,\xi}^{(1)}(h) + h_r \right) g_{x,\xi}^{A,\lambda}(y).$$

□

We observe that in the above lemma, the phase space vector field \tilde{H}^B coincides with the Hamiltonian vector field of the magnetic flow $\chi_B(h)$ in (22). This makes it possible for

us to relate the quantum evolution associated to $\text{Op}^A(h)$ with the classical flow $\chi_B(h)$ associated to h .

6.2. Propagation along the phase space flow. Based on the representation (37) in Lemma 6.2, we see that if a time-dependent wavepacket w satisfies

$$D_t w(t; x, \xi) + (i\tilde{H}^B + m^A(h) + \mathcal{R})w(t; x, \xi) = 0, \quad w(t; x, \xi) = g_{x, \xi}^{A, \lambda},$$

on the phase space \mathbb{R}^{2d} , then it will also satisfy

$$(D_t + \text{Op}^A(h))w(t, x, \xi; y) = 0$$

on the configuration space \mathbb{R}^d . A solution $u(t)$ to (9) can then be constructed as a superposition of such wavepackets w with the initial data $\tilde{u}_0 := \mathcal{T}_\lambda^A u_0$, that is

$$u(t; y) = \int_{\mathbb{R}^{2d}} w(t, y; x, \xi) \tilde{u}_0(x, \xi) dx d\xi.$$

However, the existence of such w is unclear due to the complicated error term $\mathcal{R}w$. We instead first show the existence of a wavepacket solution to a simpler equation.

Lemma 6.3. *For h satisfying Assumption 1.3, with associated magnetic Hamiltonian vector field \tilde{H}^B and $m^A(h)$ defined in (43), there exists a wavepacket $w \in C(\mathbb{R}, C^\infty(\mathbb{R}^{2d}))$, which satisfies the equation*

$$\begin{cases} D_t w(t; x, \xi) + (i\tilde{H}^B + m^A(h))w(t; x, \xi) = 0, \\ w(0; x, \xi) = g_{x, \xi}^{A, \lambda}. \end{cases}$$

Proof. By Lemma 5.2, the magnetic Hamiltonian flow $\chi_B(h)(t, 0) : (x, \xi) \mapsto (x^t, \xi^t)$ associated to h exists globally, and we can construct a time-dependent wavepacket by setting

$$g_{x^t, \xi^t}^{A, \lambda}(y) := \chi_B(h)(t, 0) g_{x, \xi}^{A, \lambda}(y), \quad y \in \mathbb{R}^d.$$

Then this wavepacket $g_{x^t, \xi^t}^{A, \lambda}$ satisfies

$$\partial_t g_{x^t, \xi^t}^{A, \lambda}(y) = \tilde{H}^B g_{x^t, \xi^t}^{A, \lambda}(y), \quad g_{x^t, \xi^t}^{A, \lambda}|_{t=0}(y) = g_{x, \xi}^{A, \lambda}(y).$$

We can also construct a phase term $e^{i\psi(t)}$, with a phase function ψ based on $m^A(h)$, by

$$\psi(t; x, \xi) := \int_0^t -m^A(h)(\tau; x^\tau, \xi^\tau) d\tau,$$

so that $\psi(t)$ satisfies

$$\partial_t \psi(t) = -m^A(h)(t; x^t, \xi^t), \quad e^{i\psi(t)}|_{t=0} = 1.$$

We modulate the wavepacket with the phase term $e^{i\psi(t)}$ and define

$$g_\psi^{A,\lambda}(t; y, x, \xi) := e^{i\psi(t)} g_{x^t, \xi^t}^{A,\lambda}(y).$$

The modulated wavepacket $g_\psi^{A,\lambda}$ now satisfies

$$\begin{aligned} \partial_t g_\psi^{A,\lambda}(t; y, x, \xi) &= i\partial_t \psi(t) e^{i\psi(t)} g_{x^t, \xi^t}^A(y) + e^{i\psi(t)} \partial_t g_{x^t, \xi^t}^{A,\lambda}(y) \\ (46) \qquad \qquad \qquad &= -im^A(h) e^{i\psi(t)} g_{x^t, \xi^t}^{A,\lambda}(y) + e^{i\psi(t)} \tilde{H}^B g_{x^t, \xi^t}^{A,\lambda}(y) \\ &= \left(-im^A(h) + \tilde{H}^B \right) g_\psi^{A,\lambda}(t; y, x, \xi), \end{aligned}$$

with $g_\psi^{A,\lambda}(0; y, x, \xi) = g_{x, \xi}^{A,\lambda}(y)$ as desired. \square

Using the wavepacket solution $g_\psi^{A,\lambda}(t)$ from the above lemma, we can construct a parametrix to equation (9).

Proposition 6.4. *Let $0 \leq s, t \leq T$. We define the operator $\tilde{S}(t, s)$ by*

$$(47) \qquad \tilde{S}(t, s)u_0(y) := \int_{\mathbb{R}^{2d} \times \mathbb{R}^d} g_\psi^{A,\lambda}(t; y, x, \xi) \overline{g_\psi^{A,\lambda}(s; y', x, \xi)} u_0(y') \, dy' dx d\xi,$$

for $u_0 \in \mathcal{S}(\mathbb{R}^d)$. We also define the operator $K(t, s)$ by

$$(48) \qquad K(t, s)u_0(y) := \int (\mathcal{R}_1 + \mathcal{R}_2) g_\psi^{A,\lambda}(t; y, x, \xi) \overline{g_\psi^{A,\lambda}(s; y', x, \xi)} u_0(y') \, dy' dx d\xi,$$

for $u_0 \in \mathcal{S}(\mathbb{R}^d)$. We then have

$$(49) \qquad (D_t + \text{Op}^A(h)) \tilde{S}(t, s)u_0 = K(t, s)u_0, \quad u_0 \in \mathcal{S}(\mathbb{R}^d).$$

Moreover, for $1 \leq p \leq \infty, m \in \mathbb{R}$ and fixed t, s , the operator $\tilde{S}(t, s)$ is bounded on $M_A^{m,p}(\mathbb{R}^d)$. The operators $\tilde{S}(t, \cdot)$ and $\tilde{S}(\cdot, s)$ are also strongly continuous on $M_A^{m,p}(\mathbb{R}^d)$.

Proof. By construction of \tilde{S} based on the wavepacket solution $g_\psi^{A,\lambda}(t)$, we can calculate using Lemma 6.2 and (46)

$$\begin{aligned} \text{Op}^A(h) g_\psi^{A,\lambda}(t; y, x, \xi) &= e^{i\psi(t)} (i\tilde{H}^B + \mathcal{R}) g_{x^t, \xi^t}^{A,\lambda}(y) \\ &= (i\tilde{H}^B + m^A(h) + \mathcal{R}_1 + \mathcal{R}_2) g_\psi^{A,\lambda}(t; y, x, \xi), \end{aligned}$$

and

$$D_t(g_\psi^{A,\lambda})(t; y, x, \xi) = -(m^A(h) + i\tilde{H}^B) g_\psi^{A,\lambda}(t; y, x, \xi).$$

Then the equation (49) can be verified based on the above calculations

$$\begin{aligned}
& (D_t + \text{Op}^A(h))\tilde{S}(t, s)u_0(y) \\
&= \int_{\mathbb{R}^{2d} \times \mathbb{R}^d} (D_t + \text{Op}^A(h))(g_\psi^{A,\lambda})(t; y, x, \xi) \overline{g_\psi^{A,\lambda}}(s; y', x, \xi) u_0(y') dy' dx d\xi \\
&= \int_{\mathbb{R}^{2d} \times \mathbb{R}^d} (\mathcal{R}_1 + \mathcal{R}_2)(g_\psi^{A,\lambda})(t; y, x, \xi) \overline{g_\psi^{A,\lambda}}(s; y', x, \xi) u_0(y') dy' dx d\xi \\
&= K(t, s)u_0(y).
\end{aligned}$$

For the $M_A^{m,p}(\mathbb{R}^d)$ boundedness of $\tilde{S}(t, s)$, in view of Lemma 4.9, it suffices to show the phase space kernel of $\tilde{S}(t, s)$ has sufficient off-diagonal decay of the form $\langle \lambda(z - x^t) \rangle^{-N} \langle \lambda^{-1}(\zeta - \xi^t) \rangle^{-N'}$. Note that by definition the $M_A^{m,p}(\mathbb{R}^d)$ norm of $\tilde{S}(t, s)u_0$ equals the weighted $L^p(\mathbb{R}^{2d})$ norm on the phase space

$$\|\tilde{S}(t, s)u_0\|_{M_A^{m,p}(\mathbb{R}^d)} = \|\nu_{\lambda,m} \mathcal{T}_\lambda^A \tilde{S}(t, s)u_0\|_{L^p(\mathbb{R}^{2d})}.$$

We fully write out the wavepacket transform $\mathcal{T}_\lambda^A \tilde{S}(t, s)u_0$ to see that

$$\begin{aligned}
& \mathcal{T}_\lambda^A \tilde{S}(t, s)u_0(z, \zeta) \\
&= \int_{\mathbb{R}^{2d}} \int_{\mathbb{R}^d} \overline{g_{z,\zeta}^{A,\lambda}}(y) g_{x^t, \xi^t}^{A,\lambda}(y) dy \left(e^{i\psi(t) - i\psi(s)} \int_{\mathbb{R}^d} \overline{g_{x^s, \xi^s}^{A,\lambda}}(y') u_0(y') dy' \right) dx d\xi.
\end{aligned}$$

Denote by $G^{A,\lambda}$ the integral kernel

$$(50) \quad G^{A,\lambda}(z, \zeta, x^t, \xi^t) := \int_{\mathbb{R}^d} \overline{g_{z,\zeta}^{A,\lambda}}(y) g_{x^t, \xi^t}^{A,\lambda}(y) dy.$$

Then $\mathcal{T}_\lambda^A \tilde{S}(t, s)u_0$ can be seen as associated to the phase space kernel $G^{A,\lambda}$

$$\mathcal{T}_\lambda^A \tilde{S}(t, s)u_0(z, \zeta) = \int_{\mathbb{R}^{2d}} G^{A,\lambda}(z, \zeta, x^t, \xi^t) e^{i\psi(t) - i\psi(s)} \tilde{u}_0(x^s, \xi^s) dx d\xi,$$

We can then write out the weighted version as

$$\begin{aligned}
& \nu_{\lambda,m}(z, \zeta) \mathcal{T}_\lambda^A \tilde{S}(t, s)u_0(z, \zeta) \\
&= \int_{\mathbb{R}^{2d}} \nu_{\lambda,m}(z, \zeta) G^{A,\lambda}(z, \zeta, x^t, \xi^t) \nu_{\lambda,-m}(x^s, \xi^s) e^{i\psi(t) - i\psi(s)} \nu_{\lambda,m}(x^s, \xi^s) \tilde{u}_0(x^s, \xi^s) dx d\xi,
\end{aligned}$$

and note the weighted phase space kernel is related to the unweighted $G^{A,\lambda}$ by

$$\begin{aligned}
(51) \quad & |\nu_{\lambda,m}(z, \zeta) G^{A,\lambda}(z, \zeta, x^t, \xi^t) \nu_{\lambda,-m}(x^s, \xi^s)| \\
& \lesssim \langle \lambda^{-1}(z - x^t) \rangle^m \langle \lambda^{-1}(\zeta - \xi^t) \rangle^m \nu_{\lambda,m}(x^t, \xi^t) \nu_{\lambda,-m}(x^s, \xi^s) |G^{A,\lambda}(z, \zeta, x^t, \xi^t)| \\
& \lesssim C_{T,h,B}^m \langle \lambda^{-1}(z - x^t) \rangle^m \langle \lambda^{-1}(\zeta - \xi^t) \rangle^m |G^{A,\lambda}(z, \zeta, x^t, \xi^t)|,
\end{aligned}$$

where we used the flow property (26) to bound $\nu_{\lambda,m}(x^t, \xi^t) \nu_{\lambda,-m}(x^s, \xi^s) \lesssim C_{T,h,B}^m$.

By the volume-preservation property in Lemma 5.2, we can replace $dx d\xi$ with $dx^s d\xi^s$, so that

$$\begin{aligned} \|\nu_{\lambda,m} e^{i\psi(t)-i\psi(s)} \tilde{u}_0\|_{L^p(\mathbb{R}^{2d})} &= \left(\int_{\mathbb{R}^{2d}} |\nu_{\lambda,m}(x^s, \xi^s) e^{i\psi(t)-i\psi(s)} \tilde{u}_0(x^s, \xi^s)|^p dx^s d\xi^s \right)^{1/p} \\ &= \|u_0\|_{M_A^{m,p}(\mathbb{R}^d)}. \end{aligned}$$

So by Young's inequality for integral operators and the relation (51), the $M_A^{m,p}(\mathbb{R}^d)$ boundedness of $\tilde{S}(t, s)$ will be implied by the off-diagonal decay of the kernel

$$(52) \quad |G^{A,\lambda}(z, \zeta, x^t, \xi^t)| \lesssim \langle \lambda(z - x^t) \rangle^{-N} \langle \lambda^{-1}(\zeta - \xi^t) \rangle^{-N'},$$

for sufficiently large $N, N' \in \mathbb{N}$. To show the above bound, we notice that based on (15), the product of the two wavepackets $\overline{g_{z,\zeta}^{A,\lambda}}$ and $g_{x^t,\xi^t}^{A,\lambda}$ can be complemented with a phase factor $e^{i\varphi^A(z,x^t)}$ to make a factor $e^{i\Gamma^B}$ appear. That is,

$$\begin{aligned} &e^{-i\varphi^A(z,x^t)} e^{-i\zeta \cdot (z-x^t)} \overline{g_{z,\zeta}^{A,\lambda}}(y) g_{x^t,\xi^t}^{A,\lambda}(y) \\ &= \lambda^d e^{i(\xi^t - \zeta) \cdot (y-x^t)} e^{-i\varphi^A(y,z)} e^{i\varphi^A(y,x^t)} e^{-i\varphi^A(z,x^t)} g(\lambda(y-z)) g(\lambda(y-x^t)) \\ &= \lambda^d e^{i(\xi^t - \zeta) \cdot (y-x^t)} e^{i\Gamma^B(y,x^t,z)} g(\lambda(y-z)) g(\lambda(y-x^t)). \end{aligned}$$

We can then make use of the derivative bounds (16) of $e^{i\Gamma^B}$, which yield for $\lambda \geq 1$

$$|\partial_{\lambda y}^\alpha e^{i\Gamma^B}(y, x^t, z)| \leq C_{B,\alpha} (\langle \lambda(y-x^t) \rangle + \langle \lambda(y-z) \rangle)^{|\alpha|}, \quad |\alpha| \geq 1.$$

Denote part of the wavepacket product by

$$(53) \quad W_{B,\lambda}(y, x^t, z) := e^{i\Gamma^B(y,x^t,z)} g(\lambda(y-z)) g(\lambda(y-x^t)).$$

By the derivative bounds of $e^{i\Gamma^B}$ above and $g \in \mathcal{S}(\mathbb{R}^d)$, we see that the partial derivatives of $W_{B,\lambda}$ with respect to λy can be bounded for all $N, N' \in \mathbb{N}$ by

$$|\partial_{\lambda y}^\alpha W_{B,\lambda}(y, x^t, z)| \leq C_{N,N',\alpha,B} \langle \lambda(y-z) \rangle^{-N} \langle \lambda(y-x^t) \rangle^{-N'}.$$

Using $\langle \lambda(y-z) \rangle^{-N} \lesssim \langle \lambda(y-x^t) \rangle^N \langle \lambda(z-x^t) \rangle^{-N}$, we further obtain for all $N, N', N'' \in \mathbb{N}$

$$(54) \quad \begin{aligned} &|\partial_{\lambda y}^\alpha W_{B,\lambda}(y, x^t, z)| \\ &\lesssim C_{N,N',N'',\alpha,B} \langle \lambda(z-x^t) \rangle^{-N} \langle \lambda(y-z) \rangle^{-N'} \langle \lambda(y-x^t) \rangle^{-N''}. \end{aligned}$$

We now re-write $G^{A,\lambda}$ with $W_{B,\lambda}$ as

$$(55) \quad \begin{aligned} &G^{A,\lambda}(z, \zeta, x^t, \xi^t) \\ &= e^{i\varphi^A(z,x^t)} e^{i\zeta \cdot (z-x^t)} \int_{\mathbb{R}^d} e^{i(\xi^t - \zeta) \cdot (y-x^t)} W_{B,\lambda}(y, x^t, z) d\lambda y, \end{aligned}$$

and we can obtain a decay factor $\langle \lambda^{-1}(\zeta - \xi^t) \rangle^{-N'}$ by integration by parts repeatedly with respect to the λy -integral, relying on the identity

$$\langle \lambda^{-1}(\zeta - \xi^t) \rangle^{-2} (1 - \Delta_{\lambda y}) e^{i\lambda^{-1}(\xi^t - \zeta) \cdot \lambda y} = e^{i(\xi^t - \zeta) \cdot y}.$$

Now combined with the $W_{B,\lambda}$ bounds (54), we see by integration by parts

$$\langle \lambda(z - x^t) \rangle^N \langle \lambda^{-1}(\zeta - \xi^t) \rangle^{N'} |G^{A,\lambda}(z, \zeta, x^t, \xi^t)| \lesssim C_{B,N,N'}$$

as desired.

For the strong continuity of \tilde{S} , by the fact that $\tilde{S}(t, s)$ is continuous on $M_A^{m,p}(\mathbb{R}^d)$ for fixed t, s , it suffices to show the strong continuity on the dense subspace $\mathcal{S}(\mathbb{R}^d)$ of $M_A^{m,p}(\mathbb{R}^d)$. Here we show the strong continuity for $\tilde{S}(\cdot, s)$, and the strong continuity of $\tilde{S}(t, \cdot)$ is analogous. Since the flow $\chi_B(t, s)$ is continuous in t , and the wavepackets $g_\psi^{A,\lambda}$ are dominated by the Gaussian g , that is

$$|g_\psi^{A,\lambda}(\tau; y, x, \xi)| \leq |g(\lambda(y - x))|,$$

we can use the dominated convergence theorem to see at the limit $t \rightarrow s$, we recover the inversion formula in Proposition 4.5. That is,

$$\begin{aligned} \lim_{t \rightarrow s} \tilde{S}(t, s) u_0(y) &= \int_{\mathbb{R}^{2d} \times \mathbb{R}^d} \lim_{t \rightarrow s} g_\psi^{A,\lambda}(t; y, x, \xi) \overline{g_\psi^{A,\lambda}(s; y', x, \xi)} u_0(y') dy' dx d\xi \\ &= \int_{\mathbb{R}^{2d} \times \mathbb{R}^d} g_{x^s, \xi^s}^{A,\lambda}(y) \overline{g_{x^s, \xi^s}^{A,\lambda}(y')} u_0(y') dy' dx^s d\xi^s \\ &= u_0(y), \end{aligned}$$

so the strong continuity of \tilde{S} holds for all $u_0 \in \mathcal{S}(\mathbb{R}^d)$ and by continuity of $\tilde{S}(t, s)$ and density of $\mathcal{S}(\mathbb{R}^d)$ in $M_A^{m,p}(\mathbb{R}^d)$ it holds on the whole space $M_A^{m,p}(\mathbb{R}^d)$. \square

7. MODULATION SPACE WELL-POSEDNESS

We show the well-posedness of the generalized magnetic Schrödinger equation (9) by constructing its propagator, based on the parametrix construction in equation (49) and Duhamel's principle. For this we need the following definition of a new space and a lemma for the Volterra equation involving the operator K defined in the Proposition 6.4.

Definition 7.1. Define the time-dependent wavepacket transform $\mathcal{T}_{\lambda,t}^A$ on $u \in \mathcal{S}'([0, T] \times \mathbb{R}^d)$ by

$$\tilde{u}(t; x^t, \xi^t) := \mathcal{T}_{\lambda,t}^A u(t; x^t, \xi^t) := \int_{\mathbb{R}^d} \overline{g_{x^t, \xi^t}^{A,\lambda}(y)} u(t; y) dy.$$

For $1 \leq p \leq \infty$ and $m \in \mathbb{R}$, we define the Banach spaces $\mathcal{M}_A^{m,p}(\mathbb{R}^d, L^1([0, T]))$ characterized by the mixed L^p -norm of the transform \tilde{u} on $[0, T] \times \mathbb{R}^{2d}$. That is, we set

$$\begin{aligned} \|u\|_{\mathcal{M}_A^{m,p}(\mathbb{R}^d, L^1([0, T]))} &:= \left(\int_{\mathbb{R}^{2d}} \left(\nu_{\lambda, m}(x, \xi) \int_{[0, T]} |\tilde{u}(t; x^t, \xi^t)| dt \right)^p dx d\xi \right)^{1/p}, \quad 1 \leq p < \infty, \\ \|u\|_{\mathcal{M}_A^{m,\infty}(\mathbb{R}^d, L^1([0, T]))} &:= \operatorname{ess\,sup}_{(x, \xi) \in \mathbb{R}^{2d}} \nu_{\lambda, m}(x, \xi) \int_{[0, T]} |\tilde{u}(t; x^t, \xi^t)| dt, \quad p = \infty, \end{aligned}$$

and define for $1 \leq p \leq \infty$

$$\mathcal{M}_A^{m,p}(\mathbb{R}^d, L^1([0, T])) := \left\{ u \in \mathcal{S}'([0, T] \times \mathbb{R}^d) \mid \|u\|_{\mathcal{M}_A^{m,p}(\mathbb{R}^d, L^1([0, T]))} < \infty \right\}.$$

Lemma 7.2. *Let $T > 0$ be sufficiently small. For every $u_0 \in M_A^{m,p}(\mathbb{R}^d)$ there exists a unique solution $v \in \mathcal{M}_A^{m,p}(\mathbb{R}^d, L^1([0, T]))$ to the Volterra equation*

$$(56) \quad v(t; y) = -K(t, 0)u_0(y) - i \int_0^t K(t, s)v(s; y) ds, \quad y \in \mathbb{R}^d, t > 0.$$

Moreover, for the solution v we have

$$(57) \quad \|v\|_{\mathcal{M}_A^{m,p}(\mathbb{R}^d, L^1([0, T]))} \lesssim C_{d,h,B} \|u_0\|_{M_A^{m,p}(\mathbb{R}^d)}.$$

Proof. In view of Lemma 4.9 and using arguments regarding the weights $\nu_{\lambda, m}$ similar to the ones given in the proof of Proposition 6.4, it suffices to prove the unweighted version with $m = 0$ as long as the relevant phase space kernel has sufficient off-diagonal decay. In the following we show the unweighted case with the spaces $M_A^p(\mathbb{R}^d)$ and $\mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T])) := \mathcal{M}_A^{0,p}(\mathbb{R}^d, L^1([0, T]))$, from which the desired off-diagonal decay of the kernels will become apparent. The necessary modifications for the weighted case are outlined at the end of the proof.

By the Banach fixed point theorem, to show the unique existence of a solution to (56), it suffices to show that the mapping Ψ_{u_0} defined by

$$(58) \quad \Psi_{u_0}(v)(y) := -K(t, 0)u_0(y) - i \int_0^t K(t, s)v(s; y) ds, \quad v \in \mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T])),$$

is a contraction mapping on $\mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T]))$.

We first show that $K(t, 0)u_0$ is in the space $\mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T]))$, by showing $K(t, 0)$ is a bounded linear operator between $M_A^p(\mathbb{R}^d)$ and $\mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T]))$, that is

$$(59) \quad \|K(\cdot, 0)u_0\|_{\mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T]))} \leq C_K \|u_0\|_{M_A^p(\mathbb{R}^d)}, \quad u_0 \in M_A^p(\mathbb{R}^d).$$

The above bound can be obtain by interpolating the following bounds (cf. [15, Section 1.18.7, Theorem 1])

$$(60) \quad \begin{aligned} \|K(\cdot, 0)u_0\|_{\mathcal{M}_A^1(\mathbb{R}^d, L^1([0, T]))} &\leq C'_K \|u_0\|_{M_A^1(\mathbb{R}^d)}, \\ \|K(\cdot, 0)u_0\|_{\mathcal{M}_A^\infty(\mathbb{R}^d, L^1([0, T]))} &\leq C''_K \|u_0\|_{M_A^\infty(\mathbb{R}^d)}. \end{aligned}$$

We begin with the bound for $K(t, 0) : M_A^1(\mathbb{R}^d) \rightarrow \mathcal{M}_A^1(\mathbb{R}^d, L^1([0, T]))$. By definition of K in (48), we have

$$(61) \quad \begin{aligned} &\mathcal{T}_{\lambda, t}^A K(t, 0)u_0(z^t, \zeta^t) \\ &= \int_{\mathbb{R}^d} \overline{g_{z^t, \zeta^t}^{A, \lambda}}(y) \left(\int_{\mathbb{R}^{2d}} (\mathcal{R}_1 + \mathcal{R}_2) g_{x^t, \xi^t}^{A, \lambda}(y) e^{i\psi(t)} \tilde{u}_0(x, \xi) \, dx d\xi \right) dy \\ &= \int_{\mathbb{R}^{2d}} \left(\int_{\mathbb{R}^d} \overline{g_{z^t, \zeta^t}^{A, \lambda}}(y) (\mathcal{R}_1 + \mathcal{R}_2) g_{x^t, \xi^t}^{A, \lambda}(y) \, dy \right) e^{i\psi(t)} \tilde{u}_0(x, \xi) \, dx d\xi. \end{aligned}$$

Denote the integral kernel above as G_K , with

$$(62) \quad G_K(t; x, \xi, z, \zeta) := \int_{\mathbb{R}^d} \overline{g_{z^t, \zeta^t}^{A, \lambda}}(y) (\mathcal{R}_1 + \mathcal{R}_2) g_{x^t, \xi^t}^{A, \lambda}(y) \, dy.$$

We claim that G_K satisfies the bound

$$(63) \quad \begin{aligned} &|G_K(t; x, \xi, z, \zeta)| \\ &\leq C_{N, h, B} (\lambda^{C_N} + \lambda^{-2} \langle x^t \rangle^{-1-\epsilon} |\xi^t|) \langle \lambda(z^t - x^t) \rangle^{-N} \langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^{-N}, \quad N \in \mathbb{N}, \end{aligned}$$

which we will prove in the next section as part of Lemma 7.3. From this, we obtain

$$\begin{aligned} &\|K(t, 0)u_0\|_{\mathcal{M}_A^1(\mathbb{R}^d, L^1([0, T]))} \\ &= \int_{\mathbb{R}^{2d}} \int_{[0, T]} |\mathcal{T}_{\lambda, t}^A K(t, 0)u_0(z^t, \zeta^t)| \, dt \, dz d\zeta \\ &\lesssim \int_{\mathbb{R}^{2d}} \int_{[0, T]} \left[\left(\int_{\mathbb{R}^{2d}} \langle \lambda(z^t - x^t) \rangle^{-d-1} \langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^{-d-1} \, dz^t d\zeta^t \right) \times \right. \\ &\quad \left. \times C_{d, h, B} (\lambda^{C_d} + \lambda^{-2} \langle x^t \rangle^{-1-\epsilon} |\xi^t|) |\tilde{u}_0(x, \xi)| \right] \, dt dx d\xi \\ &\lesssim \int_{\mathbb{R}^{2d}} \left(\int_{[0, T]} C_{d, h, B} (\lambda^{C_d} + \lambda^{-2} \langle x^t \rangle^{-1-\epsilon} |\xi^t|) \, dt \right) |\tilde{u}_0(x, \xi)| \, dx d\xi \\ &\lesssim C_{d, h, B} (T \lambda^{C_d} + \lambda^{-2} C_\epsilon (1 + T)) \int_{\mathbb{R}^{2d}} |\tilde{u}_0(x, \xi)| \, dx d\xi \\ &\lesssim C_{d, h, B, \epsilon} (T \lambda^{C_d} + \lambda^{-2}) \|u_0\|_{M_A^1(\mathbb{R}^d)}, \end{aligned}$$

where for the t -integral we used Lemma 5.3.

For the bound $K(t, 0) : M_A^\infty(\mathbb{R}^d) \rightarrow \mathcal{M}_A^\infty(\mathbb{R}^d, L^1([0, T]))$, we proceed similarly as above, except that we use the other bound for G_K from Lemma 7.3. That is,

$$(64) \quad \begin{aligned} & |G_K(t; x, \xi, z, \zeta)| \\ & \lesssim C_{N,h,B}(\lambda^{C_N} + \lambda^{-2}\langle z^t \rangle^{-1-\epsilon}|\zeta^t|)\langle \lambda(z^t - x^t) \rangle^{-N}\langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^{-N}, \quad N \in \mathbb{N}. \end{aligned}$$

With the above bound and Lemma 5.2, we obtain

$$\begin{aligned} & \|K(t, 0)u_0\|_{\mathcal{M}_A^\infty(\mathbb{R}^d, L^1([0, T]))} \\ & = \operatorname{ess\,sup}_{(z, \zeta) \in \mathbb{R}^{2d}} \int_{[0, T]} |\mathcal{T}_{\lambda, t}^A K(t, 0)u_0(z^t, \zeta^t)| \, dt \\ & \lesssim \operatorname{ess\,sup}_{(z, \zeta) \in \mathbb{R}^{2d}} \int_{[0, T]} \left[C_{d,h,B}(\lambda^{C_d} + \lambda^{-2}\langle z^t \rangle^{-1-\epsilon}|\zeta^t|) \times \right. \\ & \quad \left. \times \int_{\mathbb{R}^{2d}} \langle \lambda(z^t - x^t) \rangle^{-d-1} \langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^{-d-1} |\tilde{u}_0(x, \xi)| \, dx d\xi \right] dt \\ & \lesssim \operatorname{ess\,sup}_{(z, \zeta) \in \mathbb{R}^{2d}} \int_{[0, T]} C_{d,h,B}(\lambda^{C_d} + \lambda^{-2}\langle z^t \rangle^{-1-\epsilon}|\zeta^t|) \, dt \times \\ & \quad \times \operatorname{ess\,sup}_{t \in [0, T]} \left(\operatorname{ess\,sup}_{(x, \xi) \in \mathbb{R}^{2d}} |\tilde{u}_0(x, \xi)| \int_{\mathbb{R}^{2d}} \langle \lambda(z^t - x^t) \rangle^{-d-1} \langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^{-d-1} \, dx^t d\xi^t \right). \end{aligned}$$

We then again use Lemma 5.3 for the t -integral and notice that

$$\int_{\mathbb{R}^{2d}} \langle \lambda(z^t - x^t) \rangle^{-d-1} \langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^{-d-1} \, dx^t d\xi^t \leq C_d$$

uniformly in t , which yields

$$\|K(t, 0)u_0\|_{\mathcal{M}_A^\infty(\mathbb{R}^d, L^1([0, T]))} \lesssim C_{d,h,B,\epsilon}(T\lambda^{C_d} + \lambda^{-2})\|u\|_{M_A^\infty(\mathbb{R}^d)}.$$

With $K(t, 0)u_0 \in \mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T]))$, to show Ψ_{u_0} defined in (58) is a contraction mapping on $\mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T]))$, it suffices to show the map $F : [0, T] \times \mathbb{R}^d \rightarrow [0, T] \times \mathbb{R}^d$ defined as

$$F(v)(t; y) := - \int_0^t K(t, s)v(s; y) \, ds$$

is a contraction mapping on $\mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T]))$. Similar to the bound (59) of $K(t, 0)$, the contraction bound

$$\|F(v)\|_{\mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T]))} \leq C_F \|v\|_{\mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T]))}, \quad C_F \in (0, 1)$$

can be obtain by interpolating the bounds

$$(65) \quad \begin{aligned} & \|F(v)\|_{\mathcal{M}_A^1(\mathbb{R}^d, L^1([0, T]))} \leq C'_F \|v\|_{\mathcal{M}_A^1(\mathbb{R}^d, L^1([0, T]))}, \quad C'_F \in (0, 1), \\ & \|F(v)\|_{\mathcal{M}_A^\infty(\mathbb{R}^d, L^1([0, T]))} \leq C''_F \|v\|_{\mathcal{M}_A^\infty(\mathbb{R}^d, L^1([0, T]))}, \quad C''_F \in (0, 1). \end{aligned}$$

Notice that for the relevant wavepacket transform $\mathcal{T}_{\lambda,t}^A F(v)(t; z^t, \zeta^t)$, we can re-write again with the integral kernel G_K defined in (62)

$$\begin{aligned} \mathcal{T}_{\lambda,t}^A F(v)(t; z^t, \zeta^t) &= \int_{\mathbb{R}^d} \overline{g_{z^t, \zeta^t}^{A, \lambda}}(y) \int_0^t \left(\int_{\mathbb{R}^{2d}} (\mathcal{R}_1 + \mathcal{R}_2) g_{x^t, \xi^t}^{A, \lambda}(y) (e^{i\psi(t) - i\psi(s)} \tilde{v}(s; x^s, \xi^s)) dx d\xi \right) ds dy \\ &= \int_{\mathbb{R}^{2d}} \left(\int_{\mathbb{R}^d} \overline{g_{z^t, \zeta^t}^{A, \lambda}}(y) (\mathcal{R}_1 + \mathcal{R}_2) g_{x^t, \xi^t}^{A, \lambda}(y) dy \right) \left(\int_0^t e^{i\psi(t) - i\psi(s)} \tilde{v}(s; x^s, \xi^s) ds \right) dx d\xi \\ &= \int_{\mathbb{R}^{2d}} G_K(t; x, \xi, z, \zeta) \left(\int_0^t e^{i\psi(t) - i\psi(s)} \tilde{v}(s; x^s, \xi^s) ds \right) dx d\xi. \end{aligned}$$

The above representation is quite similar to (61), where the function $e^{i\psi(t)} \tilde{u}_0(x, \xi)$ is replaced by the slightly more complicated $\int_0^t e^{i\psi(t) - i\psi(s)} \tilde{v}(s; x^s, \xi^s) ds$. Therefore, the proof of the bounds (60) for K can easily be modified to prove the bounds (65) for F , which again relies on the kernel bounds for G_K in Lemma 7.3.

For the contraction bound on $\mathcal{M}_A^1(\mathbb{R}^d, L^1([0, T]))$, we see that

$$\begin{aligned} &\|F(v)\|_{\mathcal{M}_A^1(\mathbb{R}^d, L^1([0, T]))} \\ &\leq \int_{\mathbb{R}^{2d}} \int_{[0, T]} \int_{\mathbb{R}^{2d}} |G_K(t; x, \xi, z, \zeta)| \left(\int_0^t |\tilde{v}(s; x^s, \xi^s)| ds \right) dx d\xi dt dz d\zeta \\ &\lesssim \int_{\mathbb{R}^{2d}} \int_{[0, T]} \left[\left(\int_{\mathbb{R}^{2d}} \langle \lambda(z^t - x^t) \rangle^{-d-1} \langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^{-d-1} dz^t d\zeta^t \right) \times \right. \\ &\quad \left. \times C_{d, h, B} (\lambda^{C_d} + \lambda^{-2} \langle x^t \rangle^{-1-\epsilon} |\xi^t|) \left(\int_0^t |\tilde{v}(s; x^s, \xi^s)| ds \right) \right] dt dx d\xi \\ &\leq \int_{\mathbb{R}^{2d}} \left(\int_{[0, T]} C_{d, h, B} (\lambda^{C_d} + \lambda^{-2} \langle x^t \rangle^{-1-\epsilon} |\xi^t|) dt \right) \operatorname{ess\,sup}_{t \in [0, T]} \left(\int_0^t |\tilde{v}(s; x^s, \xi^s)| ds \right) dx d\xi \\ &\lesssim C_{d, h, B} (T \lambda^{C_d} + \lambda^{-2} C_\epsilon (1 + T)) \int_{\mathbb{R}^{2d}} \left(\int_0^T |\tilde{v}(s; x^s, \xi^s)| ds \right) dx d\xi \\ &\leq C_{d, h, B, \epsilon} (T \lambda^{C_d} + \lambda^{-2}) \|v\|_{\mathcal{M}_A^1(\mathbb{R}^d, L^1([0, T]))}. \end{aligned}$$

For the contraction bound on $\mathcal{M}_A^\infty(\mathbb{R}^d, L^1([0, T]))$, we use

$$\begin{aligned}
& \|F(v)\|_{\mathcal{M}_A^\infty(\mathbb{R}^d, L^1([0, T]))} \\
& \leq \operatorname{ess\,sup}_{(z, \zeta) \in \mathbb{R}^{2d}} \int_{[0, T]} \int_{\mathbb{R}^{2d}} |G_K(t; x, \xi, z, \zeta)| \left(\int_0^t |\tilde{v}(s; x^s, \xi^s)| ds \right) dx d\xi dt \\
& \leq \operatorname{ess\,sup}_{(z, \zeta) \in \mathbb{R}^{2d}} \left[\int_{[0, T]} C_{d, h, B} (\lambda^{C_d} + \lambda^{-2} \langle z^t \rangle^{-1-\epsilon} |\zeta^t|) dt \times \right. \\
& \quad \left. \times \operatorname{ess\,sup}_{t \in [0, T]} \int_{\mathbb{R}^{2d}} \langle \lambda(z^t - x^t) \rangle^{-d-1} \langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^{-d-1} \left(\int_0^t |\tilde{v}(s; x^s, \xi^s)| ds \right) dx d\xi \right] \\
& \leq C_{d, h, B} (T \lambda^{C_d} + \lambda^{-2} C_\epsilon) \operatorname{ess\,sup}_{(z, \zeta) \in \mathbb{R}^{2d}} \operatorname{ess\,sup}_{t \in [0, T]} \left(\int_{\mathbb{R}^{2d}} \langle \lambda(z^t - x^t) \rangle^{-d-1} \langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^{-d-1} dx d\xi \times \right. \\
& \quad \left. \times \operatorname{ess\,sup}_{(x, \xi) \in \mathbb{R}^{2d}} \left(\int_0^t |\tilde{v}(s; x^s, \xi^s)| ds \right) \right) \\
& \leq C_{d, h, B} (T \lambda^{C_d} + \lambda^{-2} C_\epsilon (1 + T)) \operatorname{ess\,sup}_{(x, \xi) \in \mathbb{R}^{2d}} \operatorname{ess\,sup}_{t \in [0, T]} \int_0^t |\tilde{v}(s; x^s, \xi^s)| ds \\
& \leq C_{d, h, B, \epsilon} (T \lambda^{C_d} + \lambda^{-2}) \|v\|_{\mathcal{M}_A^\infty(\mathbb{R}^d, L^1([0, T]))}.
\end{aligned}$$

For given constants $C_{d, h, B}$, C_d , C_ϵ , we can choose sufficiently large λ and small T such that

$$C^* := C_{d, h, B, \epsilon} (T \lambda^{C_d} + \lambda^{-2}) < 1,$$

then by complex interpolation, for $1 \leq p \leq \infty$ the operator F is indeed a contraction on the Banach space $\mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T]))$, and we have the existence and uniqueness of v as the solution to (56) by the Banach fixed point theorem.

Moreover, using equation (56) we have for the solution v

$$\begin{aligned}
\|v\|_{\mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T]))} & \leq \|K(t, 0)u_0\|_{\mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T]))} + \|F(v)\|_{\mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T]))} \\
& \leq C_K \|u_0\|_{M_A^p(\mathbb{R}^d)} + C_F \|v\|_{\mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T]))}.
\end{aligned}$$

Since $C_F \in (0, 1)$, we obtain

$$\|v\|_{\mathcal{M}_A^p(\mathbb{R}^d, L^1([0, T]))} \leq C_K (1 - C_F)^{-1} \|u_0\|_{M_A^p(\mathbb{R}^d)}.$$

To derive the weighted version, it suffices to notice that for the weight $\nu_{\lambda, m}$ we can use the flow property (26) and (27) to obtain

$$\begin{aligned}
\nu_{\lambda, m}(z, \zeta) & \lesssim C_{T, h, B}^m \nu_{\lambda, m}(z^t, \zeta^t) \nu_{\lambda, -m}(x, \xi) \nu_{\lambda, m}(x, \xi) \\
& \lesssim \langle \lambda^{-1}(z^t - x^t) \rangle^m \langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^m \nu_{\lambda, m}(x^t, \xi^t) \nu_{\lambda, -m}(x, \xi) \nu_{\lambda, m}(x, \xi) \\
& \lesssim C_{T, h, B}^{2m} \langle \lambda^{-1}(z^t - x^t) \rangle^m \langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^m \nu_{\lambda, m}(x, \xi).
\end{aligned}$$

Then we can modify the arguments for the unweighted version in the obvious way with the above bound, and combine them with the sufficient off-diagonal decay of G_K as in (63) and (64) to obtain the desired weighted version involving $M_A^{m,p}(\mathbb{R}^d)$ and $\mathcal{M}_A^{m,p}(\mathbb{R}^d, L^1([0, T]))$. \square

We are now ready to prove Theorem 1.5.

Proof of Theorem 1.5. We define the propagator S for $u_0 \in \mathcal{S}(\mathbb{R}^d)$ by

$$S(t)u_0 := \tilde{S}(t, 0)u_0 + i \int_0^t \tilde{S}(t, s)v(s) ds, \quad 0 \leq t \leq T,$$

where v denotes the unique solution to the Volterra equation (56) in Lemma 7.2 and \tilde{S} is defined in (47). Then there holds $S(0) = \text{Id}$, and $u(t) = S(t)u_0$ solves the equation (9), since by direct computation we have

$$(D_t + \text{Op}^A(h))S(t)u_0(y) = 0.$$

Next we show for $1 \leq p \leq \infty, m \geq 0$ and $t \in [0, T]$, the propagator $S(t)$ is bounded on $M_A^{m,p}(\mathbb{R}^d)$. Fix m, p and t . By definition of $S(t)$, we have

$$(66) \quad \|S(t)u_0\|_{M_A^{m,p}(\mathbb{R}^d)} \leq \|\tilde{S}(t, 0)u_0\|_{M_A^{m,p}(\mathbb{R}^d)} + \left\| i \int_0^t \tilde{S}(t, s)v(s, \cdot) ds \right\|_{M_A^{m,p}(\mathbb{R}^d)}.$$

For the terms on the RHS above, we have by Proposition 6.4

$$(67) \quad \|\tilde{S}(t, s)u_0\|_{M_A^{m,p}(\mathbb{R}^d)} \lesssim \|u_0\|_{M_A^{m,p}(\mathbb{R}^d)},$$

so it remains to check the boundedness of the term involving v . Here, we have

$$\left\| i \int_0^t \tilde{S}(t, s)v(s; \cdot) ds \right\|_{M_A^{m,p}(\mathbb{R}^d)}^p = \int_{\mathbb{R}^{2d}} \left| \nu_{\lambda, m}(z, \zeta) \int_{\mathbb{R}^d} \overline{g_{z, \zeta}^{A, \lambda}}(y) \left(\int_0^t \tilde{S}(t, s)v(s; y) ds \right) dy \right|^p dz d\zeta,$$

which depends on the L^p -integrability of the phase space function $\tilde{v}'(t; z, \zeta)$, defined by

$$\tilde{v}'(t; z, \zeta) := \nu_{\lambda, m}(z, \zeta) \int_{\mathbb{R}^d} \overline{g_{z, \zeta}^{A, \lambda}}(y) \left(\int_0^t \tilde{S}(t, s)v(s; y) ds \right) dy.$$

Writing out the action of \tilde{S}

$$\int_0^t \tilde{S}(t, s)v(s; y) ds = \int_0^t \left(\int_{\mathbb{R}^{2d}} e^{i\psi(t) - i\psi(s)} g_{x^t, \xi^t}^{A, \lambda}(y) \tilde{v}(s; x^s, \xi^s) dx d\xi \right) ds,$$

we obtain an expression of $\tilde{v}'(t; z, \zeta)$ with the integral kernel $G^{A,\lambda}$ as defined in (50)

$$\begin{aligned}\tilde{v}'(t; z, \zeta) &= \int_{\mathbb{R}^{2d}} \nu_{\lambda,m}(z, \zeta) \left(\int_{\mathbb{R}^d} \overline{g_{z,\zeta}^{A,\lambda}}(y) g_{x^t, \xi^t}^{A,\lambda}(y) dy \right) \left(\int_0^t e^{i\psi(t)-i\psi(s)} \tilde{v}(s; x^s, \xi^s) ds \right) dx d\xi \\ &= \int_{\mathbb{R}^{2d}} \nu_{\lambda,m}(z, \zeta) G^{A,\lambda}(z, \zeta, x^t, \xi^t) \nu_{\lambda,-m}(x, \xi) \left(\nu_{\lambda,m}(x, \xi) \int_0^t e^{i\psi(t)-i\psi(s)} \tilde{v}(s; x^s, \xi^s) ds \right) dx d\xi.\end{aligned}$$

Since we have proved that $G^{A,\lambda}$ has off-diagonal decays as in (52), together with

$$\nu_{\lambda,m}(z, \zeta) \nu_{\lambda,-m}(x, \xi) \lesssim C_{T,h,B}^m \langle \lambda^{-1}(z - x^t) \rangle^m \langle \lambda^{-1}(\zeta - \xi^t) \rangle^m,$$

we can apply Young's inequality for integral operators to obtain

$$\left\| i \int_0^t \tilde{S}(t, s) v(s; \cdot) ds \right\|_{M_A^{m,p}(\mathbb{R}^d)} = \|\tilde{v}'(t)\|_{L^p(\mathbb{R}^{2d})} \lesssim C_{T,h,B}^m \left\| \nu_{\lambda,m}(x, \xi) \int_0^t e^{i\psi(t)-i\psi(s)} \tilde{v}(s; x^s, \xi^s) ds \right\|_{L^p(\mathbb{R}^{2d})}.$$

The right-hand side above is in fact uniformly bound for $t \in [0, T]$ by

$$\begin{aligned}& \left(\int_{\mathbb{R}^{2d}} \left| \nu_{\lambda,m}(x, \xi) \int_0^t e^{i\psi(t)-i\psi(s)} \tilde{v}(s; x^s, \xi^s) ds \right|^p dx d\xi \right)^{1/p} \\ & \leq \left(\int_{\mathbb{R}^{2d}} \left| \nu_{\lambda,m}(x, \xi) \int_0^T |\tilde{v}(s; x^s, \xi^s)| ds \right|^p dx d\xi \right)^{1/p} \\ & = \|v\|_{\mathcal{M}_A^{m,p}(\mathbb{R}^d, L^1([0, T]))}.\end{aligned}$$

Together with (57) in Lemma 7.2, we now have

$$(68) \quad \left\| i \int_0^t \tilde{S}(t, s) v(s; \cdot) ds \right\|_{M_A^{m,p}(\mathbb{R}^d)}^p \lesssim C_{T,h,B}^m \|v\|_{\mathcal{M}_A^{m,p}(\mathbb{R}^d, L^1([0, T]))}^p \lesssim C_{T,h,B,m} \|u_0\|_{M_A^{m,p}(\mathbb{R}^d)}^p.$$

Plugging (67) and (68) into (66), we obtain the desired bound for $S(t)$

$$\|S(t)u_0\|_{M_A^{m,p}(\mathbb{R}^d)} \lesssim C_{T,h,B,m} \|u_0\|_{M_A^{m,p}(\mathbb{R}^d)}.$$

This estimate implies the uniqueness of the solution $u(t) = S(t)u_0$, and also rules out the possibility of finite-time blow-up, so we can extend the solution globally in t .

In the case of $u_0 \in M_A^{2,p}(\mathbb{R}^d)$, the above bound implies $u(t) = S(t)u_0 \in M_A^{2,p}(\mathbb{R}^d) \subset M_A^p(\mathbb{R}^d)$ for all $t \in [0, T]$. To see $u \in W^{1,1}([0, T], M_A^p(\mathbb{R}^d))$, it suffice to note

$$\|D_t u(t)\|_{M_A^p(\mathbb{R}^d)} = \|\text{Op}^A(h)u(t)\|_{M_A^p(\mathbb{R}^d)} \lesssim C_h \|u(t)\|_{M_A^{2,p}(\mathbb{R}^d)} \lesssim C_{T,h,B} \|u_0\|_{M_A^{2,p}(\mathbb{R}^d)},$$

where the first inequality follows from the boundedness of $\text{Op}^A(h)$ given by Lemma 4.8. \square

7.1. Integral kernel estimates. In this subsection we prove the bounds (63) and (64) for the integral kernel G_K .

Lemma 7.3. *The kernel G_K defined in (62)*

$$G_K(t; x, \xi, z, \zeta) := \int_{\mathbb{R}^d} \overline{g_{z^t, \zeta^t}^{A, \lambda}}(y) (\mathcal{R}_1 + \mathcal{R}_2) g_{x^t, \xi^t}^{A, \lambda}(y) dy, \quad t \in \mathbb{R}, (x, \xi, z, \zeta) \in \mathbb{R}^{4d},$$

can be bounded by, for all $N \in \mathbb{N}$,

$$\begin{aligned} & |G_K(t; x, \xi, z, \zeta)| \\ & \lesssim C_{N, h, B} (\lambda^{C_N} + \lambda^{-2} \langle x^t \rangle^{-1-\epsilon} |\xi^t|) \langle \lambda(z^t - x^t) \rangle^{-N} \langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^{-N}, \end{aligned}$$

or alternatively

$$\begin{aligned} & |G_K(t; x, \xi, z, \zeta)| \\ & \lesssim C'_{N, h, B} (\lambda^{C'_N} + \lambda^{-2} \langle z^t \rangle^{-1-\epsilon} |\zeta^t|) \langle \lambda(z^t - x^t) \rangle^{-N} \langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^{-N}. \end{aligned}$$

Proof. We first note that it suffices to prove the first G_K bound above, since the second bound can be derived from the first one by dominating the factor $\langle x^t \rangle^{-1-\epsilon}$ and $|\xi^t|$ with

$$\begin{aligned} \langle x^t \rangle^{-1-\epsilon} &= \langle x^t - z^t + z^t \rangle^{-1-\epsilon} \\ &\lesssim \langle z^t \rangle^{-1-\epsilon} \langle x^t - z^t \rangle^{1+\epsilon} \\ &\leq \langle z^t \rangle^{-1-\epsilon} \langle \lambda(x^t - z^t) \rangle^2, \end{aligned}$$

and

$$\begin{aligned} |\xi^t| &= |\xi^t - \zeta^t + \zeta^t| \\ &\leq |\xi^t - \zeta^t| + |\zeta^t| \\ &\leq \lambda \langle \lambda^{-1}(\xi^t - \zeta^t) \rangle + |\zeta^t|. \end{aligned}$$

By linearity, we can split $G_K = G_{\mathcal{R}_1} + G_{\mathcal{R}_2}$, where

$$G_{\mathcal{R}_k}(t; x, \xi, z, \zeta) := \int_{\mathbb{R}^d} \overline{g_{z^t, \zeta^t}^{A, \lambda}}(y) \mathcal{R}_k g_{x^t, \xi^t}^{A, \lambda}(y) dy, \quad k \in \{1, 2\}.$$

We start with G_1 , involving the \mathcal{R}_1 factor as defined in (44)

$$\mathcal{R}_1 = \partial_\eta h(t; x^t, \xi^t) \cdot \left(-r_{x^t}^{(1)}(A(y, x^t)) + r_{x^t}^{(1)}(A(x^t, y)) \right).$$

Similar to (55), $G_{\mathcal{R}_1}$ can be re-written with $W_{B, \lambda}$ from (53) as

$$\begin{aligned} & G_{\mathcal{R}_1}(t; x, \xi, z, \zeta) \\ (69) \quad &= e^{i\varphi^A(z^t, x^t)} e^{i\zeta^t \cdot (z^t - x^t)} \partial_\eta h(t; x^t, \xi^t) \int_{\mathbb{R}^d} \left[e^{i\lambda^{-1}(\xi^t - \zeta^t) \cdot \lambda(y - x^t)} \times \right. \\ & \quad \left. \times \left(-r_{x^t}^{(1)}(A(y, x^t)) + r_{x^t}^{(1)}(A(x^t, y)) \right) W_{B, \lambda}(y, x^t, z^t) \right] d\lambda y. \end{aligned}$$

Notice that for the λy -integral, we can do integration by parts to gain decay factors of $\langle \lambda^{-1}(\xi^t - \zeta^t) \rangle^{-2}$, making use of the identity

$$(70) \quad \langle \lambda^{-1}(\xi^t - \zeta^t) \rangle^{-2} (1 - \Delta_{\lambda y}) e^{i\lambda^{-1}(\xi^t - \zeta^t) \cdot \lambda y} = e^{i\lambda^{-1}(\xi^t - \zeta^t) \cdot \lambda y}.$$

In order to perform the integration by parts, we check the $\partial_{\lambda y}$ -derivatives of the integrand. By definition of $r_x^{(1)}(A)$ in (36), we have

$$\begin{aligned} r_{x^t}^{(1)}(A_j(y, x^t)) &= \sum_{m,l=1}^d (y_m - x_m^t)(y_l - x_l^t) \partial_{y_m} \partial_{y_l} A_j(x^t + \theta(y - x^t), x^t) \\ &= \lambda^{-2} \sum_{m,l=1}^d \lambda(y_m - x_m^t) \lambda(y_l - x_l^t) \partial_{y_m} \partial_{y_l} A_j(x^t + \theta(y - x^t), x^t). \end{aligned}$$

So taking $\partial_{\lambda y}$ -derivatives on $r_x^{(1)}(A)$ involves taking $\partial_{\lambda y}$ -derivatives on $\partial_{y_m} \partial_{y_l} A_j$. By (35), it in turn depends on derivatives of B . Denote by $B_{jk}^{(n)}$ the n -th order partial derivatives of B_{jk} , then by the decay assumption (6) for $B_{jk}^{(n)}$ with $n \geq 1$, we have for $0 \leq s \leq 1, |\alpha| \geq 0$ and $\lambda \geq 1$

$$|\partial_{\lambda y}^\alpha B_{jk}^{(n)}(x + s(y - x))| \leq C_{\alpha,B} \langle x + s(y - x) \rangle^{-1-\epsilon} \lesssim C_{\alpha,B} \langle x \rangle^{-1-\epsilon} \langle \lambda(y - x) \rangle^{1+\epsilon}.$$

Plugging this into (35), we can bound

$$|\partial_{\lambda y}^\alpha r_{x^t}^{(1)}(A(y, x^t))| \leq C_{\alpha,B} \lambda^{-2} \langle x^t \rangle^{-1-\epsilon} \langle \lambda(y - x^t) \rangle^{4+\epsilon}, \quad |\alpha| \geq 0,$$

and the same bound holds for $r_{x^t}^{(1)}(A(x^t, y))$. Together with the λy -derivative bounds for $W_{B,\lambda}$ in (54), that is for all $N, N', N'' \in \mathbb{N}$

$$\begin{aligned} &|\partial_{\lambda y}^\alpha W_{B,\lambda}(y, x^t, z^t)| \\ &\lesssim C_{N,N',N'',\alpha,B} \langle \lambda(z^t - x^t) \rangle^{-N} \langle \lambda(y - z^t) \rangle^{-N'} \langle \lambda(y - x^t) \rangle^{-N''}, \end{aligned}$$

we can repeatedly do integration by parts relying on (70) for the λy -integral in (69) to gain the desired decays. So we obtain a bound for $G_{\mathcal{R}_1}$ of the form

$$(71) \quad \begin{aligned} &|G_{\mathcal{R}_1}(t; x, \xi, z, \zeta)| \\ &\leq |\partial_\eta h(t; x^t, \xi^t)| C_{N,B} \lambda^{-2} \langle x^t \rangle^{-1-\epsilon} \langle \lambda(z^t - x^t) \rangle^{-N} \langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^{-N} \times \\ &\quad \times \int_{\mathbb{R}^d} \langle \lambda(y - z^t) \rangle^{-d-1} \langle \lambda(y - x^t) \rangle^{-d-1} d\lambda y, \\ &\lesssim C_{N,h,B} \lambda^{-2} (1 + \langle x^t \rangle^{-1-\epsilon} |\xi^t|) \langle \lambda(z^t - x^t) \rangle^{-N} \langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^{-N}, \quad \forall N \geq 0, \end{aligned}$$

where for the last inequality we used the t -uniform boundedness of $\partial_\eta h$

$$|\partial_\eta h(t; x^t, \xi^t)| \leq C_h (1 + |x^t| + |\xi^t|).$$

For the kernel $G_{\mathcal{R}_2}$, we see that

$$\begin{aligned} G_{\mathcal{R}_2}(t; x, \xi, z, \zeta) &= \int_{\mathbb{R}^d} \overline{g_{z^t, \zeta^t}^{A, \lambda}}(y) \mathcal{R}_2 g_{x^t, \xi^t}^{A, \lambda}(y) dy \\ &= \int_{\mathbb{R}^d} \overline{g_{z^t, \zeta^t}^{A, \lambda}}(y) \text{Op}_{KN}^A \left(r_{x^t, \xi^t}^{(1)}(h) + h_r \right) g_{x^t, \xi^t}^{A, \lambda}(y) dy \\ &=: \int_{\mathbb{R}^d} \overline{g_{z^t, \zeta^t}^{A, \lambda}}(y) \text{Op}_{KN}^A \left(r^*(t; y, \eta, x, \xi) \right) g_{x^t, \xi^t}^{A, \lambda}(y) dy, \end{aligned}$$

which shows that the boundedness of $G_{\mathcal{R}_2}$ relies on the boundedness of the symbol

$$r^* := r_{x^t, \xi^t}^{(1)}(h) + h_r.$$

The symbol h_r is relatively simple, since by definition (38), $h_r(t) \in S^{0, (0)}(\mathbb{R}^{2d})$, so its derivatives are fully bounded, i.e.,

$$|\partial_y^\alpha \partial_\eta^\beta h_r(t; y, \eta)| \leq C_{h, \alpha, \beta}, \quad |\alpha| + |\beta| \geq 0.$$

On the other hand, the symbol $r_{x^t, \xi^t}^{(1)}(h)$ and its derivatives can be bounded by factors of $|y - x^t|$ and $|\eta - \xi^t|$. By definition (40) for $r_{x^t, \xi^t}^{(1)}(h)$, we see that

$$|\partial_y^\alpha \partial_\eta^\beta r_{x^t, \xi^t}^{(1)}(h)(t; y, \eta)| \lesssim \sum_{0 \leq |\gamma_1| + |\gamma_2| \leq 2} C_{h, \alpha, \beta} |y - x^t|^{|\gamma_1|} |\eta - \xi^t|^{|\gamma_2|}.$$

So we bound r^* and its derivatives by

$$(72) \quad |\partial_y^\alpha \partial_\eta^\beta r^*(t; y, \eta, x^t, \xi^t)| \lesssim C_{h, \alpha, \beta} \left(1 + \sum_{0 \leq |\gamma_1| + |\gamma_2| \leq 2} |y - x^t|^{|\gamma_1|} |\eta - \xi^t|^{|\gamma_2|} \right).$$

In the following, the variable t does not play a role, so we suppress it from the notation. We first take a closer look at the pseudo-differentiation

$$\begin{aligned} &\text{Op}_{KN}^A(r^*) g_{x, \xi}^{A, \lambda}(y) \\ &= (2\pi)^{-d} \int_{\mathbb{R}^{2d}} e^{i(y-y') \cdot \eta} e^{-i\varphi^A(y', y)} r^*(y, \eta, x, \xi) g_{x, \xi}^{A, \lambda}(y') dy' d\eta \\ &= (2\pi^{-1} \lambda^{\frac{1}{2}})^d e^{i\varphi^A(y, x)} e^{i(y-x) \cdot \xi} \times \\ &\quad \times \int_{\mathbb{R}^{2d}} e^{i[(y-x) - (y'-x)] \cdot (\eta - \xi)} r^*(y, \eta, x, \xi) e^{i\Gamma^B(y', x, y)} g(\lambda(y' - x)) dy' d\eta \\ &=: (2\pi^{-1} \lambda^{\frac{1}{2}})^d e^{i\varphi^A(y, x)} e^{i(y-x) \cdot \xi} I_{\lambda, B}(r^*, g)(y, x, \xi), \end{aligned}$$

here $I_{\lambda, B}(r^*, g)$ denotes the integral part. We claim $I_{\lambda, B}(r^*, g)$ and its λy -derivatives satisfy

$$(73) \quad |\partial_{\lambda y}^\alpha I_{\lambda, B}(r^*, g)(y, x, \xi)| \lesssim C_{\alpha, B} \lambda^{C_N} \langle \lambda(y - x) \rangle^{-N}, \quad N \in \mathbb{N}.$$

So up to some λ factors, $I_{\lambda,B}(r^*, g)(y, x, \xi)$ can play a similar role as the re-scaled Gaussian $g(\lambda(y-x))$, and in this sense $G_{\mathcal{R}_2}$ is not so different from the simpler kernel $G^{A,\lambda}$ defined in (50). Notice that we can re-write $I_{\lambda,B}(r^*, g)$ as

$$\begin{aligned} & I_{\lambda,B}(r^*, g)(y, x, \xi) \\ &= \int_{\mathbb{R}^d} \left[e^{i\lambda(y-x)\cdot\lambda^{-1}(\eta-\xi)} r^*(y, \eta, x, \xi) \times \right. \\ & \quad \left. \times \left(\int_{\mathbb{R}^d} e^{-i\lambda(y'-x)\cdot\lambda^{-1}(\eta-\xi)} e^{i\Gamma^B(y', x, y)} g(\lambda(y'-x)) d\lambda y' \right) \right] d\lambda^{-1}\eta, \end{aligned}$$

where we substituted $dy'd\eta$ with $d\lambda y' d\lambda^{-1}\eta$. To verify (73), we do λy -differentiation term by term inside the integral $I_{\lambda,B}(r^*, g)$. For the phase term involving y , we see factors that can be dominated by $\langle \lambda^{-1}(\eta - \xi) \rangle$, since

$$(74) \quad |\partial_{\lambda y_j} e^{i\lambda(y-x)\cdot\lambda^{-1}(\eta-\xi)}| = \lambda^{-1} |\eta_j - \xi_j| \leq \langle \lambda^{-1}(\eta - \xi) \rangle.$$

For the λy -differentiation on r^* , similar to (72), by $\lambda \geq 1$ it can be bounded as

$$(75) \quad |\partial_{\lambda y}^\alpha \partial_{\lambda^{-1}\eta}^\beta r^*(y, \eta, x, \xi)| \lesssim C_{h,\alpha,\beta} (\lambda^{|\beta|} + \lambda^{|\beta|+2} \langle \lambda(y-x) \rangle^2 \langle \lambda^{-1}(\eta - \xi) \rangle^2).$$

We note the extra λ^C factors on the RHS above are due to the fact that the η -dependence of r^* is always in the form of $(\eta - \xi)$, and each $\partial_{\lambda^{-1}\eta}$ produces a λ factor

$$\partial_{\lambda^{-1}\eta_j}(\eta_j - \xi_j) \leq \lambda \partial_{\lambda^{-1}\eta_j} \lambda^{-1}(\eta_j - \xi_j) \leq \lambda.$$

For the λy -differentiation on $e^{i\Gamma^B(y', x, y)}$, by the derivative bounds (16) of $e^{i\Gamma^B}$, for $\lambda \geq 1$ and $|\alpha| \geq 1$ we have

$$\begin{aligned} |\partial_{\lambda y}^\alpha e^{i\Gamma^B}(y', x, y)| &\leq C_{\alpha,B} (\langle \lambda(y-y') \rangle + \langle \lambda(y-x) \rangle)^{|\alpha|} \\ &\lesssim C_{\alpha,B} (\langle \lambda(y-x) \rangle \langle \lambda(y'-x) \rangle + \langle \lambda(y-x) \rangle)^{|\alpha|} \\ &\lesssim C_{\alpha,B} [(1 + \langle \lambda(y'-x) \rangle) \langle \lambda(y-x) \rangle]^{|\alpha|}. \end{aligned}$$

Because $g(\lambda(y'-x))$ is Schwartz in $\lambda(y'-x)$, any factor of $\langle \lambda(y'-x) \rangle$ will be harmless. To show the bound (73), it remains to show that we can control factors of $\langle \lambda^{-1}(\eta - \xi) \rangle$ and $\langle \lambda(y-x) \rangle$. Notice we can gain a decaying factor of $\langle \lambda(y-x) \rangle^{-2}$ repeatedly by integration by parts with respect to the $\lambda^{-1}\eta$ -integral using

$$(76) \quad \langle \lambda(y-x) \rangle^{-2} (1 - \Delta_{\lambda^{-1}\eta}) e^{i\lambda(y-x)\cdot\lambda^{-1}\eta} = e^{i\lambda(y-x)\cdot\lambda^{-1}\eta},$$

and gain a decaying factor $\langle \lambda^{-1}(\eta - \xi) \rangle^{-2}$ factors by repeatedly integration by parts with respect to the $\lambda y'$ -integral using

$$(77) \quad \langle \lambda^{-1}(\eta - \xi) \rangle^{-2} (1 - \Delta_{\lambda y'}) e^{i\lambda^{-1}(\eta-\xi)\cdot\lambda y'} = e^{i\lambda^{-1}(\eta-\xi)\cdot\lambda y'}.$$

For the integration by parts of the $\lambda^{-1}\eta$ -integral, we need to check the $\partial_{\lambda^{-1}\eta}$ derivatives of r^* and the $\partial_{\lambda^{-1}\eta}$ derivatives of the $\lambda y'$ -integral. For r^* , we use again the derivative bounds (75). For the $\partial_{\lambda^{-1}\eta}$ derivatives of the λy -integral

$$I_{\lambda y'} := \int_{\mathbb{R}^d} e^{-i\lambda(y'-x)\cdot\lambda^{-1}(\eta-\xi)} e^{i\Gamma^B(y',x,y)} g(\lambda(y'-x)) d\lambda y',$$

each $\lambda^{-1}\eta$ -differentiation only produces a factor that can be dominated by the harmless factor $\langle\lambda(y'-x)\rangle$, that is

$$(78) \quad |\partial_{\lambda^{-1}\eta_j} e^{-i\lambda(y'-x)\cdot\lambda^{-1}(\eta-\xi)}| = \lambda|y'_j - x_j| \leq \langle\lambda(y'-x)\rangle.$$

So by repeated integration by parts relying on (76), we can obtain decay in $\langle\lambda(y-x)\rangle^{-N}$ with the cost of some λ^{CN} factor and a $\langle\lambda^{-1}(\eta-\xi)\rangle^2$ factor from (75). For the integration by parts (77) inside the $I_{\lambda y'}$ integral, we also note by the derivative bounds (16) of $e^{i\Gamma^B}$, for $\lambda \geq 1$ and $|\alpha| \geq 1$ we have

$$|\partial_{\lambda y'}^\alpha e^{i\Gamma^B}(y',x,y)| \lesssim C_{\alpha,B}[(1 + \langle\lambda(y-x)\rangle)\langle\lambda(y'-x)\rangle]^{|\alpha|}.$$

Combined with the fact that $g(\lambda(y'-x))$ is Schwartz in $\lambda(y'-x)$, we obtain

$$(79) \quad \left| \partial_{\lambda y'}^\alpha (e^{i\Gamma^B(y',x,y)} g(\lambda(y'-x))) \right| \lesssim C_{N,\alpha,B} (1 + \langle\lambda(y-x)\rangle^{|\alpha|}) \langle\lambda(y'-x)\rangle^{-N}, \quad N \in \mathbb{N}.$$

This means by integration by parts (77) for the $I_{\lambda y'}$ integral, we obtain sufficient decay of $\langle\lambda^{-1}(\eta-\xi)\rangle^{-N}$ at the cost of some finite power of $\langle\lambda(y-x)\rangle$, which will be dominated by the sufficient decay of $\langle\lambda(y-x)\rangle^{-N}$ we obtained from the integration by parts (76) in the $\lambda^{-1}\eta$ -integral. Then the claimed bound (73) holds by aforementioned integration by parts and bounds (74), (75), (78) and (79).

With the decay property of $I_{\lambda,B}(r^*,g)$ in (73), we define $W'_{B,\lambda}$ similar to $W_{B,\lambda}$ in (53)

$$W'_{B,\lambda}(y, x^t, \xi^t, z^t) := e^{i\Gamma^B(y,x^t,z^t)} g(\lambda(y-z^t)) I_{\lambda,B}(r^*,g)(y, x^t, \xi^t),$$

so that combining (16) and (73) we obtain a decay estimate similar to (54), that is for all $N, N', N'' \in \mathbb{N}$, we have

$$(80) \quad \begin{aligned} & \left| \partial_{\lambda y}^\alpha W'_{B,\lambda}(y, x^t, \xi^t, z^t) \right| \\ & \lesssim C_{N,N',N'',\alpha,B} \lambda^{CN''} \langle\lambda(z^t - x^t)\rangle^{-N} \langle\lambda(y - z^t)\rangle^{-N'} \langle\lambda(y - x^t)\rangle^{-N''}. \end{aligned}$$

Now we can re-write $G_{\mathcal{R}_2}$ with $W'_{B,\lambda}$ as

$$\begin{aligned}
& G_{\mathcal{R}_2}(t; x, \xi, z, \zeta) \\
&= \int_{\mathbb{R}^d} \overline{g_{z^t, \zeta^t}^{A, \lambda}}(y) \lambda^{\frac{d}{2}} e^{i\varphi^A(y, x^t)} e^{i(y-x^t) \cdot \xi^t} I_{\lambda, B}(r^*, g)(y, x^t, \xi^t) dy \\
&= e^{i\varphi^A(z^t, x^t)} e^{i\zeta^t \cdot (z^t - x^t)} \times \\
&\quad \times \int_{\mathbb{R}^d} e^{i(\xi^t - \zeta^t) \cdot (y - x^t)} e^{i\Gamma^B(y, x^t, z^t)} g(\lambda(y - z^t)) I_{\lambda, B}(r^*, g)(y, x^t, \xi^t) d\lambda y \\
&= e^{i\varphi^A(z^t, x^t)} e^{i\zeta^t \cdot (z^t - x^t)} \int_{\mathbb{R}^d} e^{i(\xi^t - \zeta^t) \cdot (y - x^t)} W'_{B, \lambda}(y, x^t, \xi^t, z^t) d\lambda y,
\end{aligned}$$

and integration by parts again based on

$$\langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^{-2} (1 - \Delta_{\lambda y}) e^{i\lambda^{-1}(\zeta - \xi) \cdot \lambda y} = e^{i(\zeta - \xi) \cdot \lambda y}.$$

together with the decay bounds of $W'_{B, \lambda}$ in (80), we obtain for all $N \in \mathbb{N}$

$$|G_{\mathcal{R}_2}(t; x, \xi, z, \zeta)| \lesssim C_{\alpha, B} \lambda^{C_N} \langle \lambda(z^t - x^t) \rangle^{-N} \langle \lambda^{-1}(\zeta^t - \xi^t) \rangle^{-N}.$$

Combining the above bound with the $G_{\mathcal{R}_1}$ bound in (71), we have the desired bound for G_K . \square

8. EXTENSION TO THE CASE OF TIME-DEPENDENT MAGNETIC FIELDS

In this section we consider a time-dependent magnetic field $B(t)$ with components $B_{jk} \in C^1(\mathbb{R}, C^\infty(\mathbb{R}^d))$, satisfying the assumptions in Theorem 1.7. In particular, we recall the decay condition (10) for $B_{jk}(t)$ and $\dot{B}_{jk}(t)$, that is, for some $\epsilon > 0$

$$|\partial^\alpha B_{jk}(t; y)| + |\partial^\alpha \dot{B}_{jk}(t; y)| \leq C_{\alpha, B} \langle y \rangle^{-1-\epsilon}, \quad |\alpha| \geq 1.$$

In principal, by substituting $A(x)$ with $A(t; x)$ constructed from $B(t)$, we can adapt most of the proof for Theorem 1.5 to prove Theorem 1.7. In the following, we only highlight the necessary changes needed beyond obvious substitution and straightforward calculations.

Recalling the construction for the time-independent case, we define analogously the time-dependent vector potentials $A(t)$ and phase function $\varphi^{A(t)}$ by

$$(81) \quad \begin{aligned} A_j(t; y, x) &:= - \sum_{k=1}^d \int_0^1 s(y_k - x_k) B_{jk}(t; x + s(y - x)) ds, \quad y \in \mathbb{R}^d, \\ A_j(t; y) &:= - \sum_{k=1}^d \int_0^1 s y_k B_{jk}(t; sy) ds = A_j(t; y, 0), \\ \varphi^{A(t)}(y, x) &:= (y - x) \cdot \int_0^1 A(t; (1 - s)x + sy, 0) ds. \end{aligned}$$

We can further construct time-dependent magnetic pseudo-differential operators

$$\text{Op}^{A(t)}(h)u(y) := (2\pi)^{-d} \int_{\mathbb{R}^{2d}} e^{i(y-y') \cdot \eta} e^{-i\varphi^{A(t)}(y', y)} h\left(\frac{y+y'}{2}, \eta\right) u(y') dy' d\eta, \quad y \in \mathbb{R}^d,$$

and time-dependent magnetic wavepackets

$$g_{x, \xi}^{A(t), \lambda}(y) := \lambda^{\frac{d}{2}} e^{i\xi \cdot (y-x)} g(\lambda(y-x)) e^{i\varphi^{A(t)}(y, x)}, \quad y \in \mathbb{R}^d,$$

and also accordingly the transform $\mathcal{T}_\lambda^{A(t)}$ and the magnetic modulation spaces $M_{A(t)}^{m, p}$.

With the above constructions, Lemma 6.2 holds with obvious changes since the time dependence does not play a role there. It suffices to substitute the A and B in Lemma 6.2 with the time-dependent ones $A(t)$ and $B(t)$, in particular (43), (44) and (45) should be replaced by

$$(82) \quad \begin{aligned} m^{A(t)}(h)(x, \xi) &:= h(x, \xi) - \partial_\eta h(x, \xi) \cdot (\xi + A(t; x)) \\ \mathcal{R}_1(t)g_{x, \xi}^{A(t), \lambda}(y) &:= \partial_\eta h(x, \xi) \cdot \left(-r_x^{(1)}(A(t; y, x)) + r_x^{(1)}(A(t; x, y))\right) g_{x, \xi}^{A(t), \lambda}(y) \\ \mathcal{R}_2(t)g_{x, \xi}^{A(t), \lambda}(y) &:= \text{Op}_{KN}^{A(t)}\left(r_{x, \xi}^{(1)}(h) + h_r\right) g_{x, \xi}^{A(t), \lambda}(y). \end{aligned}$$

On the other hand, some further modification is required for Lemma 6.3 and Proposition 6.4, because it involves t -derivatives. We define a new vector field $\tilde{H}^{B(t)}$

$$\tilde{H}_j^{B(t)} := \left(\partial_{\xi_j} h(t) \partial_{x_j}, -\partial_{x_j} h(t) \partial_{\xi_j} + \dot{A}_j(t; x) \partial_{\xi_j} + \sum_{k=1}^d B_{kj}(t; x) \partial_{\xi_k} h(t) \partial_{\xi_j} \right), \quad 1 \leq j \leq d,$$

and the associated flow $\chi'_B(h)(t, s) : (x^s, \xi^s) \mapsto (x^t, \xi^t)$ satisfying

$$\begin{cases} \dot{x}_j^t = \partial_{\xi_j} h(t; x^t, \xi^t), \\ \dot{\xi}_j^t = -\partial_{x_j} h(t; x^t, \xi^t) + \dot{A}_j(t; x^t) + \sum_{k=1}^d B_{kj}(t; x^t) \partial_{\xi_k} h(t; x^t, \xi^t). \end{cases}$$

Notice that $\tilde{H}^{B(t)}$ is also divergence-free and by our assumptions of $B(t)$ and construction of $A(t)$, we have

$$|\dot{A}_j(t; x^t)| \lesssim 1 + |x^t|,$$

so it is straightforward to verify that Lemma 5.2 and Lemma 5.3 also hold for $\chi'_B(h)$.

In order to adapt Proposition 6.4 for the time-dependent magnetic potentials, we define

$$(83) \quad \begin{aligned} \psi'(t; x, \xi) &:= \int_0^t -m^{A(\tau)}(h)(\tau; x^\tau, \xi^\tau) d\tau, \\ g_{\psi'}^{A(t), \lambda}(t; y, x, \xi) &:= e^{i\psi'(t)} \chi'_B(h)(t, 0) g_{x, \xi}^{A(t), \lambda}(y). \end{aligned}$$

Lemma 8.1. *The wavepacket $g_{\psi'}^{A(t), \lambda}$ associated to the flow $\chi'_B(h)$ satisfies*

$$(D_t + \text{Op}^{A(t)}(h)) g_{\psi'}^{A(t), \lambda}(t; y, x, \xi) = (\mathcal{R}_1 + \mathcal{R}_2 + \mathcal{R}_3)(t) g_{\psi'}^{A(t), \lambda}(t; y, x, \xi),$$

where \mathcal{R}_1 and \mathcal{R}_2 are as defined in (82), and for some $c \in (0, 1)$,

$$(84) \quad \mathcal{R}_3(t; y, x^t) := \sum_{j=1}^d (y_j - x_j^t) \int_0^1 \partial_x \dot{A}_j(t; x^t + cs(y - x^t)) \cdot s(y - x^t) ds,$$

which satisfies the bound

$$(85) \quad |\mathcal{R}_3(t; y, x^t)| \leq C_B \langle y - x^t \rangle^2, \quad t \in \mathbb{R}.$$

Proof. By the same calculation as in Lemma 6.2, we have

$$\begin{aligned} \text{Op}^{A(t)}(h) g_{\psi'}^{A(t), \lambda}(t; y, x, \xi) &= i \sum_{j=1}^d \left(\partial_{\xi_j} h \partial_{x_j} - (\partial_{x_j} h - \sum_{k=1}^d B_{kj}(t; x) \partial_{\xi_k} h) \partial_{\xi_j} \right) g_{\psi'}^{A(t), \lambda}(t; y, x, \xi) \\ &\quad + (m^{A(t)}(h) + \mathcal{R}_1(t) + \mathcal{R}_2(t)) g_{\psi'}^{A(t), \lambda}(t; y, x, \xi) \end{aligned}$$

with the terms defined in (82). Then it suffices to show that the t -derivative of $g_{\psi'}^{A(t), \lambda}$ is of the form

$$(86) \quad \begin{aligned} D_t g_{\psi'}^{A(t), \lambda}(t; y, x, \xi) &= -i \sum_{j=1}^d \left(\tilde{H}_j^{B(t)} - \dot{A}_j(t; x) \partial_{\xi_j} \right) g_{\psi'}^{A(t), \lambda}(t; y, x, \xi) \\ &\quad - (m^{A(t)}(h) - \mathcal{R}_3(t)) g_{\psi'}^{A(t), \lambda}(t; y, x, \xi), \end{aligned}$$

with \mathcal{R}_3 satisfying the bound (85). We claim

$$D_t e^{i\varphi^{A(t)}(y, x)} = \left((y - x) \cdot \dot{A}(t; x) + \mathcal{R}_3(t; y, x) \right) e^{i\varphi^{A(t)}(y, x)},$$

which combined with the construction of $g_{\psi'}^{A(t), \lambda}$ as in (83) gives (86). Firstly notice

$$\partial_t e^{i\varphi^{A(t)}(y, x)} = i\varphi \dot{A}(t)(y, x) e^{i\varphi^{A(t)}(y, x)}, \quad \varphi \dot{A}(t) := d\varphi^{A(t)}/dt.$$

Recall our construction in (81), from which we get

$$\begin{aligned}
\varphi^{\dot{A}(t)}(y, x) &= (y - x) \cdot \int_0^1 \dot{A}(t; (1 - s)x + sy, 0) ds \\
&= \sum_{j=1}^d (y_j - x_j) \int_0^1 \dot{A}_j(t; x + s(y - x), 0) ds \\
&= \sum_{j=1}^d (y_j - x_j) \int_0^1 \dot{A}_j(t; x) ds \\
&\quad + \sum_{j=1}^d (y_j - x_j) \int_0^1 \partial_x \dot{A}_j(t; x + cs(y - x)) \cdot s(y - x) ds,
\end{aligned}$$

where we used the Taylor expansion of \dot{A}_j at x . With \mathcal{R}_3 as defined in (84), we see that

$$\varphi^{\dot{A}(t)}(y, x) = (y - x) \cdot \dot{A}(t; x) + \mathcal{R}_3(t; y, x).$$

For the first term on the RHS above we use the identity

$$[(y - x) \cdot \dot{A}(t; x)] g_{\psi'}^{A(t), \lambda}(t; y, x, \xi) = -i \dot{A}(t; x) \cdot \partial_\xi g_{\psi'}^{A(t), \lambda}(t; y, x, \xi).$$

For the \mathcal{R}_3 term

$$\mathcal{R}_3(t; y, x) := \sum_{j=1}^d (y_j - x_j) \int_0^1 \partial_x \dot{A}_j(t; x + cs(y - x)) \cdot s(y - x) ds,$$

we see explicitly for the integrand

$$\begin{aligned}
\partial_i \dot{A}_j(t; x + cs(y - x)) &= -\partial_i \left(\sum_{k=1}^d \int_0^1 \theta(x_k + cs(y_k - x_k)) \dot{B}_{jk}(t; \theta(x + cs(y - x))) d\theta \right) \\
&= -\int_0^1 (\theta - cs) \dot{B}_{ji}(t; \theta(x + cs(y - x))) d\theta \\
&\quad - \sum_{k=1}^d \int_0^1 \theta(x_k + cs(y_k - x_k)) \partial_i \dot{B}_{jk}(t; \theta(x + cs(y - x))) d\theta.
\end{aligned}$$

By the boundedness of $B(t)$, $\dot{B}(t)$ and the decay assumption (10), we see that

$$\sup_{x, y} |\partial_i \dot{A}_j(t; x + cs(y - x))| \leq C_B, \quad t \in \mathbb{R}.$$

By the above bound and our definition (84) of \mathcal{R}_3 , we have the bound (85) as desired. \square

With Lemma 8.1, we can now adapt Proposition 6.4 to the case of $A(t)$.

Proposition 8.2. *Let $0 \leq s, t \leq T$. We define the operator $\tilde{S}(t, s)$ by*

$$(87) \quad \tilde{S}(t, s)u_0(y) := \int_{\mathbb{R}^{2d} \times \mathbb{R}^d} g_{\psi'}^{A(t), \lambda}(t; y, x, \xi) \overline{g_{\psi'}^{A(s), \lambda}(s; y', x, \xi)} u_0(y') dy' dx d\xi,$$

for $u_0 \in \mathcal{S}(\mathbb{R}^d)$. We also define the operator $K(t, s)$ by

$$(88) \quad K(t, s)u_0(y) := \int (\mathcal{R}_1 + \mathcal{R}_2 + \mathcal{R}_3)(t) g_{\psi}^{A(t), \lambda}(t; y, x, \xi) \overline{g_{\psi}^{A(s), \lambda}(s; y', x, \xi)} u_0(y') dy' dx d\xi,$$

for $u_0 \in \mathcal{S}(\mathbb{R}^d)$. We then have

$$(89) \quad (D_t + \text{Op}^{A(t)}(h))\tilde{S}(t, s)u_0 = K(t, s)u_0, \quad u_0 \in \mathcal{S}(\mathbb{R}^d).$$

Moreover, for $1 \leq p \leq \infty, m \in \mathbb{R}$ and fixed t, s , the operator $\tilde{S}(t, s)$ is bounded from $M_{A(s)}^{m,p}(\mathbb{R}^d)$ to $M_{A(t)}^{m,p}(\mathbb{R}^d)$.

With the above Proposition, we can prove adapted versions of Lemma 7.3 and Lemma 7.2 by making the obvious changes of substituting A with $A(t)$ to the relevant proofs. The arguments for the time-independent case holds for the adapted case because the time-dependent $\mathcal{R}_1(t), \mathcal{R}_2(t), \mathcal{R}_3(t)$ have bounds that are comparable to the time-independent case. Then eventually Theorem 1.7 can be derived from the adapted lemmas similarly to the derivation of Theorem 1.5.

REFERENCES

- [1] rad Benyi, Karlheinz Grochenig, Kasso A. Okoudjou, and Luke G. Rogers. Unimodular Fourier multipliers for modulation spaces. *J. Funct. Anal.*, 246(2):366–384, 2007.
- [2] rad Benyi and Kasso A. Okoudjou. *Modulation spaces—with applications to pseudodifferential operators and nonlinear Schrodinger equations*. Applied and Numerical Harmonic Analysis. Birkhuser/Springer, New York, 2020.
- [3] Elena Cordero, Karlheinz Grochenig, Fabio Nicola, and Luigi Rodino. Wiener algebras of Fourier integral operators. *J. Math. Pures Appl. (9)*, 99(2):219–233, 2013.
- [4] Elena Cordero, Fabio Nicola, and Luigi Rodino. Schrodinger equations with rough Hamiltonians. *Discrete Contin. Dyn. Syst.*, 35(10):4805–4821, 2015.
- [5] Horia D. Cornean, Bernard Helffer, and Radu Purice. A Beals criterion for magnetic pseudo-differential operators proved with magnetic Gabor frames. *Comm. Partial Differential Equations*, 43(8):1196–1204, 2018.
- [6] Shin-ichi Doi. Smoothness of solutions for Schrodinger equations with unbounded potentials. *Publ. Res. Inst. Math. Sci.*, 41(1):175–221, 2005.
- [7] Karlheinz Grochenig. *Foundations of time-frequency analysis*. Applied and Numerical Harmonic Analysis. Birkhuser Boston, Inc., Boston, MA, 2001.
- [8] Viorel Iftimie, Marius Mantoiu, and Radu Purice. Magnetic pseudodifferential operators. *Publ. Res. Inst. Math. Sci.*, 43(3):585–623, 2007.

- [9] Viorel Iftimie and Radu Purice. Magnetic Fourier integral operators. *J. Pseudo-Differ. Oper. Appl.*, 2(2):141–218, 2011.
- [10] Keiichi Kato and Ryo Muramatsu. Estimates on modulation spaces for Schrödinger operators with time-dependent sub-linear vector potentials. *Hokkaido Math. J.*, 53(1):51–69, 2024.
- [11] John M. Lee. *Introduction to smooth manifolds*, volume 218 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2013.
- [12] Marius Măntoiu and Radu Purice. The magnetic Weyl calculus. *J. Math. Phys.*, 45(4):1394–1417, 2004.
- [13] Marius Măntoiu and Radu Purice. The modulation mapping for magnetic symbols and operators. *Proc. Amer. Math. Soc.*, 138(8):2839–2852, 2010.
- [14] Daniel Tataru. Phase space transforms and microlocal analysis. In *Phase space analysis of partial differential equations. Vol. II*, Pubbl. Cent. Ric. Mat. Ennio Giorgi, pages 505–524. Scuola Norm. Sup., Pisa, 2004.
- [15] H. Triebel. *Interpolation theory, function spaces, differential operators*. VEB Deutscher Verlag der Wissenschaften, Berlin, 1978.
- [16] Kenji Yajima. Schrödinger evolution equations with magnetic fields. *J. Analyse Math.*, 56:29–76, 1991.

DEPARTMENT OF MATHEMATICS, KARLSRUHE INSTITUTE OF TECHNOLOGY, 76131 KARLSRUHE, GERMANY

Email address: `dorothee.frey@kit.edu`

DEPARTMENT OF MATHEMATICS, KARLSRUHE INSTITUTE OF TECHNOLOGY, 76131 KARLSRUHE, GERMANY

Email address: `siliang.weng@kit.edu`